

PRO-ÉTALE COHOMOLOGY

0965

Contents

1. Introduction	1
2. Some topology	2
3. Local isomorphisms	4
4. Ind-Zariski algebra	5
5. Constructing w-local affine schemes	6
6. Identifying local rings versus ind-Zariski	10
7. Ind-étale algebra	13
8. Constructing ind-étale algebras	15
9. Weakly étale versus pro-étale	17
10. Constructing w-contractible covers	18
11. The pro-étale site	20
12. Points of the pro-étale site	29
13. Compact generation	30
14. Derived completion in the constant Noetherian case	30
15. Derived completion on the pro-étale site	31
16. Comparison with the étale site	33
17. Cohomology of a point	37
18. Weakly contractible hypercoverings	38
19. Functoriality of the pro-étale site	40
20. Finite morphisms and pro-étale sites	41
21. Closed immersions and pro-étale sites	41
22. Extension by zero	44
23. Constructible sheaves on the pro-étale site	46
24. Constructible adic sheaves	47
25. A suitable derived category	51
26. Proper base change	55
27. Other chapters	56
References	57

1. Introduction

0966 The material in this chapter and more can be found in the preprint [BS13].

The goal of this chapter is to introduce the pro-étale topology and show how it simplifies the introduction of ℓ -adic cohomology in algebraic geometry.

A brief overview of the history of this material as we have understood it. In [Gro77, Exposés V and VI] Grothendieck et al developed a theory for dealing with ℓ -adic

This is a chapter of the Stacks Project, version 42e2c6f8, compiled on Nov 12, 2017.

sheaves as inverse systems of sheaves of $\mathbf{Z}/\ell^n\mathbf{Z}$ -modules. In his second paper on the Weil conjectures ([Del74]) Deligne introduced a derived category of ℓ -adic sheaves as a certain 2-limit of categories of complexes of sheaves of $\mathbf{Z}/\ell^n\mathbf{Z}$ -modules on the étale site of a scheme X . This approach is used in the paper by Beilinson, Bernstein, and Deligne ([BBD82]) as the basis for their beautiful theory of perverse sheaves. In a paper entitled “Continuous Étale Cohomology” ([Jan88]) Uwe Jannsen discusses an important variant of the cohomology of a ℓ -adic sheaf on a variety over a field. His paper is followed up by a paper of Torsten Ekedahl ([Eke90]) who discusses the adic formalism needed to work comfortably with derived categories defined as limits.

The goal of this chapter is to show that, if we work with the pro-étale site of a scheme, then one can avoid some of the technicalities these authors encountered. This comes at the expense of having to work with non-Noetherian schemes, even when one is only interested in working with ℓ -adic sheaves and cohomology of such on varieties over an algebraically closed field.

2. Some topology

0967 Some preliminaries. We have defined *spectral spaces* and *spectral maps* of spectral spaces in Topology, Section 23. The spectrum of a ring is a spectral space, see Algebra, Lemma 25.2.

0968 **Lemma 2.1.** *Let X be a spectral space. Let $X_0 \subset X$ be the set of closed points. The following are equivalent*

- (1) *Every open covering of X can be refined by a finite disjoint union decomposition $X = \coprod U_i$ with U_i open and closed in X .*
- (2) *The composition $X_0 \rightarrow X \rightarrow \pi_0(X)$ is bijective.*

Moreover, if X_0 is closed in X and every point of X specializes to a unique point of X_0 , then these conditions are satisfied.

Proof. We will use without further mention that X_0 is quasi-compact (Topology, Lemma 12.9) and $\pi_0(X)$ is profinite (Topology, Lemma 23.8). Picture

$$\begin{array}{ccc} X_0 & \longrightarrow & X \\ & \searrow f & \downarrow \pi \\ & & \pi_0(X) \end{array}$$

If (2) holds, the continuous bijective map $f : X_0 \rightarrow \pi_0(X)$ is a homeomorphism by Topology, Lemma 17.8. Given an open covering $X = \bigcup U_i$, we get an open covering $\pi_0(X) = \bigcup f(X_0 \cap U_i)$. By Topology, Lemma 22.3 we can find a finite open covering of the form $\pi_0(X) = \coprod V_j$ which refines this covering. Since $X_0 \rightarrow \pi_0(X)$ is bijective each connected component of X has a unique closed point, whence is equal to the set of points specializing to this closed point. Hence $\pi^{-1}(V_j)$ is the set of points specializing to the points of $f^{-1}(V_j)$. Now, if $f^{-1}(V_j) \subset X_0 \cap U_i \subset U_i$, then it follows that $\pi^{-1}(V_j) \subset U_i$ (because the open set U_i is closed under generalizations). In this way we see that the open covering $X = \coprod \pi^{-1}(V_j)$ refines the covering we started out with. In this way we see that (2) implies (1).

Assume (1). Let $x, y \in X$ be closed points. Then we have the open covering $X = (X \setminus \{x\}) \cup (X \setminus \{y\})$. It follows from (1) that there exists a disjoint union

decomposition $X = U \amalg V$ with U and V open (and closed) and $x \in U$ and $y \in V$. In particular we see that every connected component of X has at most one closed point. By Topology, Lemma 12.8 every connected component (being closed) also does have a closed point. Thus $X_0 \rightarrow \pi_0(X)$ is bijective. In this way we see that (1) implies (2).

Assume X_0 is closed in X and every point specializes to a unique point of X_0 . Then X_0 is a spectral space (Topology, Lemma 23.4) consisting of closed points, hence profinite (Topology, Lemma 23.7). Let $x, y \in X_0$ be distinct. By Topology, Lemma 22.3 we can find a disjoint union decomposition $X_0 = U_0 \amalg V_0$ with U_0 and V_0 open and closed and $x \in U_0$ and $y \in V_0$. Let $U \subset X$, resp. $V \subset X$ be the set of points specializing to U_0 , resp. V_0 . Observe that $X = U \amalg V$. By Topology, Lemma 24.7 we see that U is an intersection of quasi-compact open subsets. Hence U is closed in the constructible topology. Since U is closed under specialization, we see that U is closed by Topology, Lemma 23.5. By symmetry V is closed and hence U and V are both open and closed. This proves that x, y are not in the same connected component of X . In other words, $X_0 \rightarrow \pi_0(X)$ is injective. The map is also surjective by Topology, Lemma 12.8 and the fact that connected components are closed. In this way we see that the final condition implies (2). \square

0969 **Example 2.2.** Let T be a profinite space. Let $t \in T$ be a point and assume that $T \setminus \{t\}$ is not quasi-compact. Let $X = T \times \{0, 1\}$. Consider the topology on X with a subbase given by the sets $U \times \{0, 1\}$ for $U \subset T$ open, $X \setminus \{(t, 0)\}$, and $U \times \{1\}$ for $U \subset T$ open with $t \notin U$. The set of closed points of X is $X_0 = T \times \{0\}$ and $(t, 1)$ is in the closure of X_0 . Moreover, $X_0 \rightarrow \pi_0(X)$ is a bijection. This example shows that conditions (1) and (2) of Lemma 2.1 do not imply the set of closed points is closed.

It turns out it is more convenient to work with spectral spaces which have the slightly stronger property mentioned in the final statement of Lemma 2.1. We give this property a name.

096A **Definition 2.3.** A spectral space X is *w-local* if the set of closed points X_0 is closed and every point of X specializes to a unique closed point. A continuous map $f : X \rightarrow Y$ of *w-local* spaces is *w-local* if it is spectral and maps any closed point of X to a closed point of Y .

We have seen in the proof of Lemma 2.1 that in this case $X_0 \rightarrow \pi_0(X)$ is a homeomorphism and that $X_0 \cong \pi_0(X)$ is a profinite space. Moreover, a connected component of X is exactly the set of points specializing to a given $x \in X_0$.

096B **Lemma 2.4.** *Let X be a w-local spectral space. If $Y \subset X$ is closed, then Y is w-local.*

Proof. The subset $Y_0 \subset Y$ of closed points is closed because $Y_0 = X_0 \cap Y$. Since X is *w-local*, every $y \in Y$ specializes to a unique point of X_0 . This specialization is in Y , and hence also in Y_0 , because $\{y\} \subset Y$. In conclusion, Y is *w-local*. \square

096C **Lemma 2.5.** *Let X be a spectral space. Let*

$$\begin{array}{ccc} Y & \longrightarrow & T \\ \downarrow & & \downarrow \\ X & \longrightarrow & \pi_0(X) \end{array}$$

be a cartesian diagram in the category of topological spaces with T profinite. Then Y is spectral and $T = \pi_0(Y)$. If moreover X is w -local, then Y is w -local, $Y \rightarrow X$ is w -local, and the set of closed points of Y is the inverse image of the set of closed points of X .

Proof. Note that Y is a closed subspace of $X \times T$ as $\pi_0(X)$ is a profinite space hence Hausdorff (use Topology, Lemmas 23.8 and 3.4). Since $X \times T$ is spectral (Topology, Lemma 23.9) it follows that Y is spectral (Topology, Lemma 23.4). Let $Y \rightarrow \pi_0(Y) \rightarrow T$ be the canonical factorization (Topology, Lemma 7.8). It is clear that $\pi_0(Y) \rightarrow T$ is surjective. The fibres of $Y \rightarrow T$ are homeomorphic to the fibres of $X \rightarrow \pi_0(X)$. Hence these fibres are connected. It follows that $\pi_0(Y) \rightarrow T$ is injective. We conclude that $\pi_0(Y) \rightarrow T$ is a homeomorphism by Topology, Lemma 17.8.

Next, assume that X is w -local and let $X_0 \subset X$ be the set of closed points. The inverse image $Y_0 \subset Y$ of X_0 in Y maps bijectively onto T as $X_0 \rightarrow \pi_0(X)$ is a bijection by Lemma 2.1. Moreover, Y_0 is quasi-compact as a closed subset of the spectral space Y . Hence $Y_0 \rightarrow \pi_0(Y) = T$ is a homeomorphism by Topology, Lemma 17.8. It follows that all points of Y_0 are closed in Y . Conversely, if $y \in Y$ is a closed point, then it is closed in the fibre of $Y \rightarrow \pi_0(Y) = T$ and hence its image x in X is closed in the (homeomorphic) fibre of $X \rightarrow \pi_0(X)$. This implies $x \in X_0$ and hence $y \in Y_0$. Thus Y_0 is the collection of closed points of Y and for each $y \in Y_0$ the set of generalizations of y is the fibre of $Y \rightarrow \pi_0(Y)$. The lemma follows. \square

3. Local isomorphisms

096D We start with a definition.

096E **Definition 3.1.** Let $\varphi : A \rightarrow B$ be a ring map.

- (1) We say $A \rightarrow B$ is a *local isomorphism* if for every prime $\mathfrak{q} \subset B$ there exists a $g \in B$, $g \notin \mathfrak{q}$ such that $A \rightarrow B_g$ induces an open immersion $\text{Spec}(B_g) \rightarrow \text{Spec}(A)$.
- (2) We say $A \rightarrow B$ *identifies local rings* if for every prime $\mathfrak{q} \subset B$ the canonical map $A_{\varphi^{-1}(\mathfrak{q})} \rightarrow B_{\mathfrak{q}}$ is an isomorphism.

We list some elementary properties.

096F **Lemma 3.2.** Let $A \rightarrow B$ and $A \rightarrow A'$ be ring maps. Let $B' = B \otimes_A A'$ be the base change of B .

- (1) If $A \rightarrow B$ is a local isomorphism, then $A' \rightarrow B'$ is a local isomorphism.
- (2) If $A \rightarrow B$ identifies local rings, then $A' \rightarrow B'$ identifies local rings.

Proof. Omitted. \square

096G **Lemma 3.3.** Let $A \rightarrow B$ and $B \rightarrow C$ be ring maps.

- (1) If $A \rightarrow B$ and $B \rightarrow C$ are local isomorphisms, then $A \rightarrow C$ is a local isomorphism.
- (2) If $A \rightarrow B$ and $B \rightarrow C$ identify local rings, then $A \rightarrow C$ identifies local rings.

Proof. Omitted. \square

096H **Lemma 3.4.** Let A be a ring. Let $B \rightarrow C$ be an A -algebra homomorphism.

- (1) If $A \rightarrow B$ and $A \rightarrow C$ are local isomorphisms, then $B \rightarrow C$ is a local isomorphism.
- (2) If $A \rightarrow B$ and $A \rightarrow C$ identify local rings, then $B \rightarrow C$ identifies local rings.

Proof. Omitted. □

096I **Lemma 3.5.** *Let $A \rightarrow B$ be a local isomorphism. Then*

- (1) $A \rightarrow B$ is étale,
- (2) $A \rightarrow B$ identifies local rings,
- (3) $A \rightarrow B$ is quasi-finite.

Proof. Omitted. □

096J **Lemma 3.6.** *Let $A \rightarrow B$ be a local isomorphism. Then there exist $n \geq 0$, $g_1, \dots, g_n \in B$, $f_1, \dots, f_n \in A$ such that $(g_1, \dots, g_n) = B$ and $A_{f_i} \cong B_{g_i}$.*

Proof. Omitted. □

096K **Lemma 3.7.** *Let $p : (Y, \mathcal{O}_Y) \rightarrow (X, \mathcal{O}_X)$ and $q : (Z, \mathcal{O}_Z) \rightarrow (X, \mathcal{O}_X)$ be morphisms of locally ringed spaces. If $\mathcal{O}_Y = p^{-1}\mathcal{O}_X$, then*

$Mor_{LRS/(X, \mathcal{O}_X)}((Z, \mathcal{O}_Z), (Y, \mathcal{O}_Y)) \longrightarrow Mor_{Top/X}(Z, Y), \quad (f, f^\sharp) \longmapsto f$

is bijective. Here $LRS/(X, \mathcal{O}_X)$ is the category of locally ringed spaces over X and Top/X is the category of topological spaces over X .

Proof. This is immediate from the definitions. □

096L **Lemma 3.8.** *Let A be a ring. Set $X = \text{Spec}(A)$. The functor*

$$B \longmapsto \text{Spec}(B)$$

from the category of A -algebras B such that $A \rightarrow B$ identifies local rings to the category of topological spaces over X is fully faithful.

Proof. This follows from Lemma 3.7 and the fact that if $A \rightarrow B$ identifies local rings, then the pullback of the structure sheaf of $\text{Spec}(A)$ via $p : \text{Spec}(B) \rightarrow \text{Spec}(A)$ is equal to the structure sheaf of $\text{Spec}(B)$. □

4. Ind-Zariski algebra

096M We start with a definition; please see Remark 6.9 for a comparison with the corresponding definition of the article [BS13].

096N **Definition 4.1.** A ring map $A \rightarrow B$ is said to be *ind-Zariski* if B can be written as a filtered colimit $B = \text{colim } B_i$ with each $A \rightarrow B_i$ a local isomorphism.

An example of an Ind-Zariski map is a localization $A \rightarrow S^{-1}A$, see Algebra, Lemma 9.9. The category of ind-Zariski algebras is closed under several natural operations.

096P **Lemma 4.2.** *Let $A \rightarrow B$ and $A \rightarrow A'$ be ring maps. Let $B' = B \otimes_A A'$ be the base change of B . If $A \rightarrow B$ is ind-Zariski, then $A' \rightarrow B'$ is ind-Zariski.*

Proof. Omitted. □

096Q **Lemma 4.3.** *Let $A \rightarrow B$ and $B \rightarrow C$ be ring maps. If $A \rightarrow B$ and $B \rightarrow C$ are ind-Zariski, then $A \rightarrow C$ is ind-Zariski.*

Proof. Omitted. □

096R **Lemma 4.4.** *Let A be a ring. Let $B \rightarrow C$ be an A -algebra homomorphism. If $A \rightarrow B$ and $A \rightarrow C$ are ind-Zariski, then $B \rightarrow C$ is ind-Zariski.*

Proof. Omitted. □

096S **Lemma 4.5.** *A filtered colimit of ind-Zariski A -algebras is ind-Zariski over A .*

Proof. Omitted. □

096T **Lemma 4.6.** *Let $A \rightarrow B$ be ind-Zariski. Then $A \rightarrow B$ identifies local rings,*

Proof. Omitted. □

5. Constructing w -local affine schemes

096U An affine scheme X is called w -local if its underlying topological space is w -local (Definition 2.3). It turns out given any ring A there is a canonical faithfully flat ind-Zariski ring map $A \rightarrow A_w$ such that $\text{Spec}(A_w)$ is w -local. The key to constructing A_w is the following simple lemma.

096V **Lemma 5.1.** *Let A be a ring. Set $X = \text{Spec}(A)$. Let $Z \subset X$ be a locally closed subscheme which is of the form $D(f) \cap V(I)$ for some $f \in A$ and ideal $I \subset A$. Then*

- (1) *there exists a multiplicative subset $S \subset A$ such that $\text{Spec}(S^{-1}A)$ maps by a homeomorphism to the set of points of X specializing to Z ,*
- (2) *the A -algebra $A_{\tilde{Z}} = S^{-1}A$ depends only on the underlying locally closed subset $Z \subset X$,*
- (3) *Z is a closed subscheme of $\text{Spec}(A_{\tilde{Z}})$,*

If $A \rightarrow A'$ is a ring map and $Z' \subset X' = \text{Spec}(A')$ is a locally closed subscheme of the same form which maps into Z , then there is a unique A -algebra map $A_{\tilde{Z}} \rightarrow (A')_{\tilde{Z}'}$.

Proof. Let $S \subset A$ be the multiplicative set of elements which map to invertible elements of $\Gamma(Z, \mathcal{O}_Z) = (A/I)_f$. If \mathfrak{p} is a prime of A which does not specialize to Z , then \mathfrak{p} generates the unit ideal in $(A/I)_f$. Hence we can write $f^n = g + h$ for some $n \geq 0$, $g \in \mathfrak{p}$, $h \in I$. Then $g \in S$ and we see that \mathfrak{p} is not in the spectrum of $S^{-1}A$. Conversely, if \mathfrak{p} does specialize to Z , say $\mathfrak{p} \subset \mathfrak{q} \supset I$ with $f \notin \mathfrak{q}$, then we see that $S^{-1}A$ maps to $A_{\mathfrak{q}}$ and hence \mathfrak{p} is in the spectrum of $S^{-1}A$. This proves (1).

The isomorphism class of the localization $S^{-1}A$ depends only on the corresponding subset $\text{Spec}(S^{-1}A) \subset \text{Spec}(A)$, whence (2) holds. By construction $S^{-1}A$ maps surjectively onto $(A/I)_f$, hence (3). The final statement follows as the multiplicative subset $S' \subset A'$ corresponding to Z' contains the image of the multiplicative subset S . □

Let A be a ring. Let $E \subset A$ be a finite subset. We get a stratification of $X = \text{Spec}(A)$ into locally closed subschemes by looking at the vanishing behaviour of the elements of E . More precisely, given a disjoint union decomposition $E = E' \amalg E''$ we set

$$096W \quad (5.1.1) \quad Z(E', E'') = \bigcap_{f \in E'} D(f) \cap \bigcap_{f \in E''} V(f) = D\left(\prod_{f \in E'} f\right) \cap V\left(\sum_{f \in E''} fA\right)$$

The points of $Z(E', E'')$ are exactly those $x \in X$ such that $f \in E'$ maps to a nonzero element in $\kappa(x)$ and $f \in E''$ maps to zero in $\kappa(x)$. Thus it is clear that

$$096X \quad (5.1.2) \quad X = \coprod_{E=E' \amalg E''} Z(E', E'')$$

set theoretically. Observe that each stratum is constructible.

096Y **Lemma 5.2.** *Let $X = \text{Spec}(A)$ as above. Given any finite stratification $X = \coprod T_i$ by constructible subsets, there exists a finite subset $E \subset A$ such that the stratification (5.1.2) refines $X = \coprod T_i$.*

Proof. We may write $T_i = \bigcup_j U_{i,j} \cap V_{i,j}^c$ as a finite union for some $U_{i,j}$ and $V_{i,j}$ quasi-compact open in X . Then we may write $U_{i,j} = \bigcup D(f_{i,j,k})$ and $V_{i,j} = \bigcup D(g_{i,j,l})$. Then we set $E = \{f_{i,j,k}\} \cup \{g_{i,j,l}\}$. This does the job, because the stratification (5.1.2) is the one whose strata are labeled by the vanishing pattern of the elements of E which clearly refines the given stratification. \square

We continue the discussion. Given a finite subset $E \subset A$ we set

$$096Z \quad (5.2.1) \quad A_E = \prod_{E=E' \amalg E''} A_{Z(E', E'')}^{\sim}$$

with notation as in Lemma 5.1. This makes sense because (5.1.1) shows that each $Z(E', E'')$ has the correct shape. We take the spectrum of this ring and denote it

$$0970 \quad (5.2.2) \quad X_E = \text{Spec}(A_E) = \prod_{E=E' \amalg E''} X_{E', E''}$$

with $X_{E', E''} = \text{Spec}(A_{Z(E', E'')}^{\sim})$. Note that

$$0971 \quad (5.2.3) \quad Z_E = \prod_{E=E' \amalg E''} Z(E', E'') \longrightarrow X_E$$

is a closed subscheme. By construction the closed subscheme Z_E contains all the closed points of the affine scheme X_E as every point of $X_{E', E''}$ specializes to a point of $Z(E', E'')$.

Let $I(A)$ be the partially ordered set of all finite subsets of A . This is a directed partially ordered set. For $E_1 \subset E_2$ there is a canonical transition map $A_{E_1} \rightarrow A_{E_2}$ of A -algebras. Namely, given a decomposition $E_2 = E'_2 \amalg E''_2$ we set $E'_1 = E_1 \cap E'_2$ and $E''_1 = E_1 \cap E''_2$. Then observe that $Z(E'_1, E''_1) \subset Z(E'_2, E''_2)$ hence a unique A -algebra map $A_{Z(E'_1, E''_1)}^{\sim} \rightarrow A_{Z(E'_2, E''_2)}^{\sim}$ by Lemma 5.1. Using these maps collectively we obtain the desired ring map $A_{E_1} \rightarrow A_{E_2}$. Observe that the corresponding map of affine schemes

$$0972 \quad (5.2.4) \quad X_{E_2} \longrightarrow X_{E_1}$$

maps Z_{E_2} into Z_{E_1} . By uniqueness we obtain a system of A -algebras over $I(A)$ and we set

$$0973 \quad (5.2.5) \quad A_w = \text{colim}_{E \in I(A)} A_E$$

This A -algebra is ind-Zariski and faithfully flat over A . Finally, we set $X_w = \text{Spec}(A_w)$ and endow it with the closed subscheme $Z = \lim_{E \in I(A)} Z_E$. In a formula

$$0974 \quad (5.2.6) \quad X_w = \lim_{E \in I(A)} X_E \supset Z = \lim_{E \in I(A)} Z_E$$

0975 **Lemma 5.3.** *Let $X = \text{Spec}(A)$ be an affine scheme. With $A \rightarrow A_w$, $X_w = \text{Spec}(A_w)$, and $Z \subset X_w$ as above.*

- (1) $A \rightarrow A_w$ is ind-Zariski and faithfully flat,
- (2) $X_w \rightarrow X$ induces a bijection $Z \rightarrow X$,
- (3) Z is the set of closed points of X_w ,
- (4) Z is a reduced scheme, and
- (5) every point of X_w specializes to a unique point of Z .

In particular, X_w is w -local (Definition 2.3).

Proof. The map $A \rightarrow A_w$ is ind-Zariski by construction. For every E the morphism $Z_E \rightarrow X$ is a bijection, hence (2). As $Z \subset X_w$ we conclude $X_w \rightarrow X$ is surjective and $A \rightarrow A_w$ is faithfully flat by Algebra, Lemma 38.16. This proves (1).

Suppose that $y \in X_w$, $y \notin Z$. Then there exists an E such that the image of y in X_E is not contained in Z_E . Then for all $E \subset E'$ also y maps to an element of $X_{E'}$ not contained in $Z_{E'}$. Let $T_{E'} \subset X_{E'}$ be the reduced closed subscheme which is the closure of the image of y . It is clear that $T = \lim_{E \subset E'} T_{E'}$ is the closure of y in X_w . For every $E \subset E'$ the scheme $T_{E'} \cap Z_{E'}$ is nonempty by construction of $X_{E'}$. Hence $\lim T_{E'} \cap Z_{E'}$ is nonempty and we conclude that $T \cap Z$ is nonempty. Thus y is not a closed point. It follows that every closed point of X_w is in Z .

Suppose that $y \in X_w$ specializes to $z, z' \in Z$. We will show that $z = z'$ which will finish the proof of (3) and will imply (5). Let $x, x' \in X$ be the images of z and z' . Since $Z \rightarrow X$ is bijective it suffices to show that $x = x'$. If $x \neq x'$, then there exists an $f \in A$ such that $x \in D(f)$ and $x' \in V(f)$ (or vice versa). Set $E = \{f\}$ so that

$$X_E = \text{Spec}(A_f) \amalg \text{Spec}(A_{\tilde{V}(f)})$$

Then we see that z and z' map x_E and x'_E which are in different parts of the given decomposition of X_E above. But then it is impossible for x_E and x'_E to be specializations of a common point. This is the desired contradiction.

Recall that given a finite subset $E \subset A$ we have Z_E is a disjoint union of the locally closed subschemes $Z(E', E'')$ each isomorphic to the spectrum of $(A/I)_f$ where I is the ideal generated by E'' and f the product of the elements of E' . Any nilpotent element b of $(A/I)_f$ is the class of g/f^n for some $g \in A$. Then setting $E' = E \cup \{g\}$ the reader verifies that b is pulled back to zero under the transition map $Z_{E'} \rightarrow Z_E$ of the system. This proves (4). \square

0976 **Remark 5.4.** Let A be a ring. Let κ be an infinite cardinal bigger or equal than the cardinality of A . Then the cardinality of A_w (Lemma 5.3) is at most κ . Namely, each A_E has cardinality at most κ and the set of finite subsets of A has cardinality at most κ as well. Thus the result follows as $\kappa \otimes \kappa = \kappa$, see Sets, Section 6.

0977 **Lemma 5.5** (Universal property of the construction). *Let A be a ring. Let $A \rightarrow A_w$ be the ring map constructed in Lemma 5.3. For any ring map $A \rightarrow B$ such that $\text{Spec}(B)$ is w -local, there is a unique factorization $A \rightarrow A_w \rightarrow B$ such that $\text{Spec}(B) \rightarrow \text{Spec}(A_w)$ is w -local.*

Proof. Denote $Y = \text{Spec}(B)$ and $Y_0 \subset Y$ the set of closed points. Denote $f : Y \rightarrow X$ the given morphism. Recall that Y_0 is profinite, in particular every constructible subset of Y_0 is open and closed. Let $E \subset A$ be a finite subset. Recall that $A_w = \text{colim } A_E$ and that the set of closed points of $\text{Spec}(A_w)$ is the limit of the closed subsets $Z_E \subset X_E = \text{Spec}(A_E)$. Thus it suffices to show there is a unique factorization $A \rightarrow A_E \rightarrow B$ such that $Y \rightarrow X_E$ maps Y_0 into Z_E . Since $Z_E \rightarrow X = \text{Spec}(A)$ is bijective, and since the strata $Z(E', E'')$ are constructible we see that

$$Y_0 = \coprod f^{-1}(Z(E', E'')) \cap Y_0$$

is a disjoint union decomposition into open and closed subsets. As $Y_0 = \pi_0(Y)$ we obtain a corresponding decomposition of Y into open and closed pieces. Thus

it suffices to construct the factorization in case $f(Y_0) \subset Z(E', E'')$ for some decomposition $E = E' \amalg E''$. In this case $f(Y)$ is contained in the set of points of X specializing to $Z(E', E'')$ which is homeomorphic to $X_{E', E''}$. Thus we obtain a unique continuous map $Y \rightarrow X_{E', E''}$ over X . By Lemma 3.7 this corresponds to a unique morphism of schemes $Y \rightarrow X_{E', E''}$ over X . This finishes the proof. \square

Recall that the spectrum of a ring is profinite if and only if every point is closed. There are in fact a whole slew of equivalent conditions that imply this. See Algebra, Lemma 25.5 or Topology, Lemma 23.7.

0978 **Lemma 5.6.** *Let A be a ring such that $\text{Spec}(A)$ is profinite. Let $A \rightarrow B$ be a ring map. Then $\text{Spec}(B)$ is profinite in each of the following cases:*

- (1) *if $\mathfrak{q}, \mathfrak{q}' \subset B$ lie over the same prime of A , then neither $\mathfrak{q} \subset \mathfrak{q}'$, nor $\mathfrak{q}' \subset \mathfrak{q}$,*
- (2) *$A \rightarrow B$ induces algebraic extensions of residue fields,*
- (3) *$A \rightarrow B$ is a local isomorphism,*
- (4) *$A \rightarrow B$ identifies local rings,*
- (5) *$A \rightarrow B$ is weakly étale,*
- (6) *$A \rightarrow B$ is quasi-finite,*
- (7) *$A \rightarrow B$ is unramified,*
- (8) *$A \rightarrow B$ is étale,*
- (9) *B is a filtered colimit of A -algebras as in (1) – (8),*
- (10) *etc.*

Proof. By the references mentioned above (Algebra, Lemma 25.5 or Topology, Lemma 23.7) there are no specializations between distinct points of $\text{Spec}(A)$ and $\text{Spec}(B)$ is profinite if and only if there are no specializations between distinct points of $\text{Spec}(B)$. These specializations can only happen in the fibres of $\text{Spec}(B) \rightarrow \text{Spec}(A)$. In this way we see that (1) is true.

The assumption in (2) implies all primes of B are maximal by Algebra, Lemma 34.9. Thus (2) holds. If $A \rightarrow B$ is a local isomorphism or identifies local rings, then the residue field extensions are trivial, so (3) and (4) follow from (2). If $A \rightarrow B$ is weakly étale, then More on Algebra, Lemma 87.17 tells us it induces separable algebraic residue field extensions, so (5) follows from (2). If $A \rightarrow B$ is quasi-finite, then the fibres are finite discrete topological spaces. Hence (6) follows from (1). Hence (3) follows from (1). Cases (7) and (8) follow from this as unramified and étale ring map are quasi-finite (Algebra, Lemmas 147.6 and 141.6). If $B = \text{colim } B_i$ is a filtered colimit of A -algebras, then $\text{Spec}(B) = \text{colim } \text{Spec}(B_i)$, hence if each $\text{Spec}(B_i)$ is profinite, so is $\text{Spec}(B)$. This proves (9). \square

0979 **Lemma 5.7.** *Let A be a ring. Let $V(I) \subset \text{Spec}(A)$ be a closed subset which is a profinite topological space. Then there exists an ind-Zariski ring map $A \rightarrow B$ such that $\text{Spec}(B)$ is w -local, the set of closed points is $V(IB)$, and $A/I \cong B/IB$.*

Proof. Let $A \rightarrow A_w$ and $Z \subset Y = \text{Spec}(A_w)$ as in Lemma 5.3. Let $T \subset Z$ be the inverse image of $V(I)$. Then $T \rightarrow V(I)$ is a homeomorphism by Topology, Lemma 17.8. Let $B = (A_w)_{\widetilde{T}}$, see Lemma 5.1. It is clear that B is w -local with closed points $V(IB)$. The ring map $A/I \rightarrow B/IB$ is ind-Zariski and induces a homeomorphism on underlying topological spaces. Hence it is an isomorphism by Lemma 3.8. \square

097A **Lemma 5.8.** *Let A be a ring such that $X = \text{Spec}(A)$ is w -local. Let $I \subset A$ be the radical ideal cutting out the set X_0 of closed points in X . Let $A \rightarrow B$ be a ring map inducing algebraic extensions on residue fields at primes. Then*

- (1) *every point of $Z = V(IB)$ is a closed point of $\text{Spec}(B)$,*
- (2) *there exists an ind-Zariski ring map $B \rightarrow C$ such that*
 - (a) *$B/IB \rightarrow C/IC$ is an isomorphism,*
 - (b) *the space $Y = \text{Spec}(C)$ is w -local,*
 - (c) *the induced map $p : Y \rightarrow X$ is w -local, and*
 - (d) *$p^{-1}(X_0)$ is the set of closed points of Y .*

Proof. By Lemma 5.6 applied to $A/I \rightarrow B/IB$ all points of $Z = V(IB) = \text{Spec}(B/IB)$ are closed, in fact $\text{Spec}(B/IB)$ is a profinite space. To finish the proof we apply Lemma 5.7 to $IB \subset B$. \square

6. Identifying local rings versus ind-Zariski

097B An ind-Zariski ring map $A \rightarrow B$ identifies local rings (Lemma 4.6). The converse does not hold (Examples, Section 38). However, it turns out that there is a kind of structure theorem for ring maps which identify local rings in terms of ind-Zariski ring maps, see Proposition 6.6.

Let A be a ring. Let $X = \text{Spec}(A)$. The space of connected components $\pi_0(X)$ is a profinite space by Topology, Lemma 23.8 (and Algebra, Lemma 25.2).

097C **Lemma 6.1.** *Let A be a ring. Let $X = \text{Spec}(A)$. Let $T \subset \pi_0(X)$ be a closed subset. There exists a surjective ind-Zariski ring map $A \rightarrow B$ such that $\text{Spec}(B) \rightarrow \text{Spec}(A)$ induces a homeomorphism of $\text{Spec}(B)$ with the inverse image of T in X .*

Proof. Let $Z \subset X$ be the inverse image of T . Then Z is the intersection $Z = \bigcap Z_\alpha$ of the open and closed subsets of X containing Z , see Topology, Lemma 12.12. For each α we have $Z_\alpha = \text{Spec}(A_\alpha)$ where $A \rightarrow A_\alpha$ is a local isomorphism (a localization at an idempotent). Setting $B = \text{colim } A_\alpha$ proves the lemma. \square

097D **Lemma 6.2.** *Let A be a ring and let $X = \text{Spec}(A)$. Let T be a profinite space and let $T \rightarrow \pi_0(X)$ be a continuous map. There exists an ind-Zariski ring map $A \rightarrow B$ such that with $Y = \text{Spec}(B)$ the diagram*

$$\begin{array}{ccc} Y & \longrightarrow & \pi_0(Y) \\ \downarrow & & \downarrow \\ X & \longrightarrow & \pi_0(X) \end{array}$$

is cartesian in the category of topological spaces and such that $\pi_0(Y) = T$ as spaces over $\pi_0(X)$.

Proof. Namely, write $T = \lim T_i$ as the limit of an inverse system finite discrete spaces over a directed set (see Topology, Lemma 22.2). For each i let $Z_i = \text{Im}(T \rightarrow \pi_0(X) \times T_i)$. This is a closed subset. Observe that $X \times T_i$ is the spectrum of $A_i = \prod_{t \in T_i} A$ and that $A \rightarrow A_i$ is a local isomorphism. By Lemma 6.1 we see that $Z_i \subset \pi_0(X \times T_i) = \pi_0(X) \times T_i$ corresponds to a surjection $A_i \rightarrow B_i$ which is ind-Zariski such that $\text{Spec}(B_i) = X \times_{\pi_0(X)} Z_i$ as subsets of $X \times T_i$. The transition maps $T_i \rightarrow T_{i'}$ induce maps $Z_i \rightarrow Z_{i'}$ and $X \times_{\pi_0(X)} Z_i \rightarrow X \times_{\pi_0(X)} Z_{i'}$. Hence ring maps $B_{i'} \rightarrow B_i$ (Lemmas 3.8 and 4.6). Set $B = \text{colim } B_i$. Because $T = \lim Z_i$ we

have $X \times_{\pi_0(X)} T = \lim X \times_{\pi_0(X)} Z_i$ and hence $Y = \text{Spec}(B) = \lim \text{Spec}(B_i)$ fits into the cartesian diagram

$$\begin{array}{ccc} Y & \longrightarrow & T \\ \downarrow & & \downarrow \\ X & \longrightarrow & \pi_0(X) \end{array}$$

of topological spaces. By Lemma 2.5 we conclude that $T = \pi_0(Y)$. \square

09BJ **Example 6.3.** Let k be a field. Let T be a profinite topological space. There exists an ind-Zariski ring map $k \rightarrow A$ such that $\text{Spec}(A)$ is homeomorphic to T . Namely, just apply Lemma 6.2 to $T \rightarrow \pi_0(\text{Spec}(k)) = \{*\}$. In fact, in this case we have

$$A = \text{colim Map}(T_i, k)$$

whenever we write $T = \lim T_i$ as a filtered limit with each T_i finite.

097E **Lemma 6.4.** *Let $A \rightarrow B$ be ring map such that*

- (1) $A \rightarrow B$ identifies local rings,
- (2) the topological spaces $\text{Spec}(B)$, $\text{Spec}(A)$ are w -local,
- (3) $\text{Spec}(B) \rightarrow \text{Spec}(A)$ is w -local, and
- (4) $\pi_0(\text{Spec}(B)) \rightarrow \pi_0(\text{Spec}(A))$ is bijective.

Then $A \rightarrow B$ is an isomorphism

Proof. Let $X_0 \subset X = \text{Spec}(A)$ and $Y_0 \subset Y = \text{Spec}(B)$ be the sets of closed points. By assumption Y_0 maps into X_0 and the induced map $Y_0 \rightarrow X_0$ is a bijection. As a space $\text{Spec}(A)$ is the disjoint union of the spectra of the local rings of A at closed points. Similarly for B . Hence $X \rightarrow Y$ is a bijection. Since $A \rightarrow B$ is flat we have going down (Algebra, Lemma 38.18). Thus Algebra, Lemma 40.11 shows for any prime $\mathfrak{q} \subset B$ lying over $\mathfrak{p} \subset A$ we have $B_{\mathfrak{q}} = B_{\mathfrak{p}}$. Since $B_{\mathfrak{q}} = A_{\mathfrak{p}}$ by assumption, we see that $A_{\mathfrak{p}} = B_{\mathfrak{p}}$ for all primes \mathfrak{p} of A . Thus $A = B$ by Algebra, Lemma 23.1. \square

097F **Lemma 6.5.** *Let $A \rightarrow B$ be ring map such that*

- (1) $A \rightarrow B$ identifies local rings,
- (2) the topological spaces $\text{Spec}(B)$, $\text{Spec}(A)$ are w -local, and
- (3) $\text{Spec}(B) \rightarrow \text{Spec}(A)$ is w -local.

Then $A \rightarrow B$ is ind-Zariski.

Proof. Set $X = \text{Spec}(A)$ and $Y = \text{Spec}(B)$. Let $X_0 \subset X$ and $Y_0 \subset Y$ be the set of closed points. Let $A \rightarrow A'$ be the ind-Zariski morphism of affine schemes such that with $X' = \text{Spec}(A')$ the diagram

$$\begin{array}{ccc} X' & \longrightarrow & \pi_0(X') \\ \downarrow & & \downarrow \\ X & \longrightarrow & \pi_0(X) \end{array}$$

is cartesian in the category of topological spaces and such that $\pi_0(X') = \pi_0(Y)$ as spaces over $\pi_0(X)$, see Lemma 6.2. By Lemma 2.5 we see that X' is w -local and the set of closed points $X'_0 \subset X'$ is the inverse image of X_0 .

We obtain a continuous map $Y \rightarrow X'$ of underlying topological spaces over X identifying $\pi_0(Y)$ with $\pi_0(X')$. By Lemma 3.8 (and Lemma 4.6) this corresponds

to a morphism of affine schemes $Y \rightarrow X'$ over X . Since $Y \rightarrow X$ maps Y_0 into X_0 we see that $Y \rightarrow X'$ maps Y_0 into X'_0 , i.e., $Y \rightarrow X'$ is w-local. By Lemma 6.4 we see that $Y \cong X'$ and we win. \square

The following proposition is a warm up for the type of result we will prove later.

097G **Proposition 6.6.** *Let $A \rightarrow B$ be a ring map which identifies local rings. Then there exists a faithfully flat, ind-Zariski ring map $B \rightarrow B'$ such that $A \rightarrow B'$ is ind-Zariski.*

Proof. Let $A \rightarrow A_w$, resp. $B \rightarrow B_w$ be the faithfully flat, ind-Zariski ring map constructed in Lemma 5.3 for A , resp. B . Since $\text{Spec}(B_w)$ is w-local, there exists a unique factorization $A \rightarrow A_w \rightarrow B_w$ such that $\text{Spec}(B_w) \rightarrow \text{Spec}(A_w)$ is w-local by Lemma 5.5. Note that $A_w \rightarrow B_w$ identifies local rings, see Lemma 3.4. By Lemma 6.5 this means $A_w \rightarrow B_w$ is ind-Zariski. Since $B \rightarrow B_w$ is faithfully flat, ind-Zariski (Lemma 5.3) and the composition $A \rightarrow B \rightarrow B_w$ is ind-Zariski (Lemma 4.3) the proposition is proved. \square

The proposition above allows us to characterize the affine, weakly contractible objects in the pro-Zariski site of an affine scheme.

09AZ **Lemma 6.7.** *Let A be a ring. The following are equivalent*

- (1) *every faithfully flat ring map $A \rightarrow B$ identifying local rings has a section,*
- (2) *every faithfully flat ind-Zariski ring map $A \rightarrow B$ has a section, and*
- (3) *A satisfies*
 - (a) *$\text{Spec}(A)$ is w-local, and*
 - (b) *$\pi_0(\text{Spec}(A))$ is extremally disconnected.*

Proof. The equivalence of (1) and (2) follows immediately from Proposition 6.6.

Assume (3)(a) and (3)(b). Let $A \rightarrow B$ be faithfully flat and ind-Zariski. We will use without further mention the fact that a flat map $A \rightarrow B$ is faithfully flat if and only if every closed point of $\text{Spec}(A)$ is in the image of $\text{Spec}(B) \rightarrow \text{Spec}(A)$. We will show that $A \rightarrow B$ has a section.

Let $I \subset A$ be an ideal such that $V(I) \subset \text{Spec}(A)$ is the set of closed points of $\text{Spec}(A)$. We may replace B by the ring C constructed in Lemma 5.8 for $A \rightarrow B$ and $I \subset A$. Thus we may assume $\text{Spec}(B)$ is w-local such that the set of closed points of $\text{Spec}(B)$ is $V(IB)$.

Assume $\text{Spec}(B)$ is w-local and the set of closed points of $\text{Spec}(B)$ is $V(IB)$. Choose a continuous section to the surjective continuous map $V(IB) \rightarrow V(I)$. This is possible as $V(I) \cong \pi_0(\text{Spec}(A))$ is extremally disconnected, see Topology, Proposition 26.6. The image is a closed subspace $T \subset \pi_0(\text{Spec}(B)) \cong V(JB)$ mapping homeomorphically onto $\pi_0(A)$. Replacing B by the ind-Zariski quotient ring constructed in Lemma 6.1 we see that we may assume $\pi_0(\text{Spec}(B)) \rightarrow \pi_0(\text{Spec}(A))$ is bijective. At this point $A \rightarrow B$ is an isomorphism by Lemma 6.4.

Assume (1) or equivalently (2). Let $A \rightarrow A_w$ be the ring map constructed in Lemma 5.3. By (1) there is a section $A_w \rightarrow A$. Thus $\text{Spec}(A)$ is homeomorphic to a closed subset of $\text{Spec}(A_w)$. By Lemma 2.4 we see (3)(a) holds. Finally, let $T \rightarrow \pi_0(A)$ be a surjective map with T an extremally disconnected, quasi-compact, Hausdorff topological space (Topology, Lemma 26.9). Choose $A \rightarrow B$ as in Lemma 6.2 adapted to $T \rightarrow \pi_0(\text{Spec}(A))$. By (1) there is a section $B \rightarrow A$. Thus we

see that $T = \pi_0(\mathrm{Spec}(B)) \rightarrow \pi_0(\mathrm{Spec}(A))$ has a section. A formal categorical argument, using Topology, Proposition 26.6, implies that $\pi_0(\mathrm{Spec}(A))$ is extremally disconnected. \square

09B0 **Lemma 6.8.** *Let A be a ring. There exists a faithfully flat, ind-Zariski ring map $A \rightarrow B$ such that B satisfies the equivalent conditions of Lemma 6.7.*

Proof. We first apply Lemma 5.3 to see that we may assume that $\mathrm{Spec}(A)$ is w-local. Choose an extremally disconnected space T and a surjective continuous map $T \rightarrow \pi_0(\mathrm{Spec}(A))$, see Topology, Lemma 26.9. Note that T is profinite. Apply Lemma 6.2 to find an ind-Zariski ring map $A \rightarrow B$ such that $\pi_0(\mathrm{Spec}(B)) \rightarrow \pi_0(\mathrm{Spec}(A))$ realizes $T \rightarrow \pi_0(\mathrm{Spec}(A))$ and such that

$$\begin{array}{ccc} \mathrm{Spec}(B) & \longrightarrow & \pi_0(\mathrm{Spec}(B)) \\ \downarrow & & \downarrow \\ \mathrm{Spec}(A) & \longrightarrow & \pi_0(\mathrm{Spec}(A)) \end{array}$$

is cartesian in the category of topological spaces. Note that $\mathrm{Spec}(B)$ is w-local, that $\mathrm{Spec}(B) \rightarrow \mathrm{Spec}(A)$ is w-local, and that the set of closed points of $\mathrm{Spec}(B)$ is the inverse image of the set of closed points of $\mathrm{Spec}(A)$, see Lemma 2.5. Thus condition (3) of Lemma 6.7 holds for B . \square

0A0D **Remark 6.9.** In each of Lemmas 6.1, 6.2, Proposition 6.6, and Lemma 6.8 we find an ind-Zariski ring map with some properties. In the paper [BS13] the authors use the notion of an ind-(Zariski localization) which is a filtered colimit of finite products of principal localizations. It is possible to replace ind-Zariski by ind-(Zariski localization) in each of the results listed above. However, we do not need this and the notion of an ind-Zariski homomorphism of rings as defined here has slightly better formal properties. Moreover, the notion of an ind-Zariski ring map is the natural analogue of the notion of an ind-étale ring map defined in the next section.

7. Ind-étale algebra

097H We start with a definition.

097I **Definition 7.1.** A ring map $A \rightarrow B$ is said to be *ind-étale* if B can be written as a filtered colimit of étale A -algebras.

The category of ind-étale algebras is closed under a number of natural operations.

097J **Lemma 7.2.** *Let $A \rightarrow B$ and $A \rightarrow A'$ be ring maps. Let $B' = B \otimes_A A'$ be the base change of B . If $A \rightarrow B$ is ind-étale, then $A' \rightarrow B'$ is ind-étale.*

Proof. This is Algebra, Lemma 149.1. \square

097K **Lemma 7.3.** *Let $A \rightarrow B$ and $B \rightarrow C$ be ring maps. If $A \rightarrow B$ and $B \rightarrow C$ are ind-étale, then $A \rightarrow C$ is ind-étale.*

Proof. This is Algebra, Lemma 149.2. \square

097L **Lemma 7.4.** *A filtered colimit of ind-étale A -algebras is ind-étale over A .*

Proof. This is Algebra, Lemma 149.3. \square

097M **Lemma 7.5.** *Let A be a ring. Let $B \rightarrow C$ be an A -algebra map of ind-étale A -algebras. Then C is an ind-étale B -algebra.*

Proof. This is Algebra, Lemma 149.4. □

097N **Lemma 7.6.** *Let $A \rightarrow B$ be ind-étale. Then $A \rightarrow B$ is weakly étale (More on Algebra, Definition 87.1).*

Proof. This follows from More on Algebra, Lemma 87.14. □

097P **Lemma 7.7.** *Let A be a ring and let $I \subset A$ be an ideal. The base change functor*

$$\text{ind-étale } A\text{-algebras} \longrightarrow \text{ind-étale } A/I\text{-algebras}, \quad C \longmapsto C/IC$$

has a fully faithful right adjoint v . In particular, given an ind-étale A/I -algebra \overline{C} there exists an ind-étale A -algebra $C = v(\overline{C})$ such that $\overline{C} = C/IC$.

Proof. Let \overline{C} be an ind-étale A/I -algebra. Consider the category \mathcal{C} of factorizations $A \rightarrow B \rightarrow \overline{C}$ where $A \rightarrow B$ is étale. (We ignore some set theoretical issues in this proof.) We will show that this category is directed and that $C = \text{colim}_{\mathcal{C}} B$ is an ind-étale A -algebra such that $\overline{C} = C/IC$.

We first prove that \mathcal{C} is directed (Categories, Definition 19.1). The category is nonempty as $A \rightarrow A \rightarrow \overline{C}$ is an object. Suppose that $A \rightarrow B \rightarrow \overline{C}$ and $A \rightarrow B' \rightarrow \overline{C}$ are two objects of \mathcal{C} . Then $A \rightarrow B \otimes_A B' \rightarrow \overline{C}$ is another (use Algebra, Lemma 141.3). Suppose that $f, g : B \rightarrow B'$ are two maps between objects $A \rightarrow B \rightarrow \overline{C}$ and $A \rightarrow B' \rightarrow \overline{C}$ of \mathcal{C} . Then a coequalizer is $A \rightarrow B' \otimes_{f, B, g} B' \rightarrow \overline{C}$. This is an object of \mathcal{C} by Algebra, Lemmas 141.3 and 141.8. Thus the category \mathcal{C} is directed.

Write $\overline{C} = \text{colim } \overline{B}_i$ as a filtered colimit with \overline{B}_i étale over A/I . For every i there exists $A \rightarrow B_i$ étale with $\overline{B}_i = B_i/IB_i$, see Algebra, Lemma 141.10. Thus $C \rightarrow \overline{C}$ is surjective. Since $C/IC \rightarrow \overline{C}$ is ind-étale (Lemma 7.5) we see that it is flat. Hence \overline{C} is a localization of C/IC at some multiplicative subset $S \subset C/IC$ (Algebra, Lemma 107.2). Take an $f \in C$ mapping to an element of $S \subset C/IC$. Choose $A \rightarrow B \rightarrow \overline{C}$ in \mathcal{C} and $g \in B$ mapping to f in the colimit. Then we see that $A \rightarrow B_g \rightarrow \overline{C}$ is an object of \mathcal{C} as well. Thus f is an invertible element of C . It follows that $C/IC = \overline{C}$.

Next, we claim that for an ind-étale algebra D over A we have

$$\text{Mor}_A(D, C) = \text{Mor}_{A/I}(D/ID, \overline{C})$$

Namely, let $D/ID \rightarrow \overline{C}$ be an A/I -algebra map. Write $D = \text{colim}_{i \in I} D_i$ as a colimit over a directed set I with D_i étale over A . By choice of \mathcal{C} we obtain a transformation $I \rightarrow \mathcal{C}$ and hence a map $D \rightarrow C$ compatible with maps to \overline{C} . Whence the claim.

It follows that the functor v defined by the rule

$$\overline{C} \longmapsto v(\overline{C}) = \text{colim}_{A \rightarrow B \rightarrow \overline{C}} B$$

is a right adjoint to the base change functor u as required by the lemma. The functor v is fully faithful because $u \circ v = \text{id}$ by construction, see Categories, Lemma 24.3. □

8. Constructing ind-étale algebras

097Q Let A be a ring. Recall that any étale ring map $A \rightarrow B$ is isomorphic to a standard smooth ring map of relative dimension 0. Such a ring map is of the form

$$A \longrightarrow A[x_1, \dots, x_n]/(f_1, \dots, f_n)$$

where the determinant of the $n \times n$ -matrix with entries $\partial f_i / \partial x_j$ is invertible in the quotient ring. See Algebra, Lemma 141.2.

Let $S(A)$ be the set of all *faithfully flat*¹ standard smooth A -algebras of relative dimension 0. Let $I(A)$ be the partially ordered (by inclusion) set of finite subsets E of $S(A)$. Note that $I(A)$ is a directed partially ordered set. For $E = \{A \rightarrow B_1, \dots, A \rightarrow B_n\}$ set

$$B_E = B_1 \otimes_A \dots \otimes_A B_n$$

Observe that B_E is a faithfully flat étale A -algebra. For $E \subset E'$, there is a canonical transition map $B_E \rightarrow B_{E'}$ of étale A -algebras. Namely, say $E = \{A \rightarrow B_1, \dots, A \rightarrow B_n\}$ and $E' = \{A \rightarrow B_1, \dots, A \rightarrow B_{n+m}\}$ then $B_E \rightarrow B_{E'}$ sends $b_1 \otimes \dots \otimes b_n$ to the element $b_1 \otimes \dots \otimes b_n \otimes 1 \otimes \dots \otimes 1$ of $B_{E'}$. This construction defines a system of faithfully flat étale A -algebras over $I(A)$ and we set

$$T(A) = \operatorname{colim}_{E \in I(A)} B_E$$

Observe that $T(A)$ is a faithfully flat ind-étale A -algebra (Algebra, Lemma 38.20). By construction given any faithfully flat étale A -algebra B there is a (non-unique) A -algebra map $B \rightarrow T(A)$. Namely, pick some $(A \rightarrow B_0) \in S(A)$ and an isomorphism $B \cong B_0$. Then the canonical coprojection

$$B \rightarrow B_0 \rightarrow T(A) = \operatorname{colim}_{E \in I(A)} B_E$$

is the desired map.

097R **Lemma 8.1.** *Given a ring A there exists a faithfully flat ind-étale A -algebra C such that every faithfully flat étale ring map $C \rightarrow B$ has a section.*

Proof. Set $T^1(A) = T(A)$ and $T^{n+1}(A) = T(T^n(A))$. Let

$$C = \operatorname{colim} T^n(A)$$

This algebra is faithfully flat over each $T^n(A)$ and in particular over A , see Algebra, Lemma 38.20. Moreover, C is ind-étale over A by Lemma 7.4. If $C \rightarrow B$ is étale, then there exists an n and an étale ring map $T^n(A) \rightarrow B'$ such that $B = C \otimes_{T^n(A)} B'$, see Algebra, Lemma 141.3. If $C \rightarrow B$ is faithfully flat, then $\operatorname{Spec}(B) \rightarrow \operatorname{Spec}(C) \rightarrow \operatorname{Spec}(T^n(A))$ is surjective, hence $\operatorname{Spec}(B') \rightarrow \operatorname{Spec}(T^n(A))$ is surjective. In other words, $T^n(A) \rightarrow B'$ is faithfully flat. By our construction, there is a $T^n(A)$ -algebra map $B' \rightarrow T^{n+1}(A)$. This induces a C -algebra map $B \rightarrow C$ which finishes the proof. \square

097S **Remark 8.2.** Let A be a ring. Let κ be an infinite cardinal bigger or equal than the cardinality of A . Then the cardinality of $T(A)$ is at most κ . Namely, each B_E has cardinality at most κ and the index set $I(A)$ has cardinality at most κ as well. Thus the result follows as $\kappa \otimes \kappa = \kappa$, see Sets, Section 6. It follows that the ring constructed in the proof of Lemma 8.1 has cardinality at most κ as well.

¹In the presence of flatness, e.g., for smooth or étale ring maps, this just means that the induced map on spectra is surjective. See Algebra, Lemma 38.16.

097T **Remark 8.3.** The construction $A \mapsto T(A)$ is functorial in the following sense: If $A \rightarrow A'$ is a ring map, then we can construct a commutative diagram

$$\begin{array}{ccc} A & \longrightarrow & T(A) \\ \downarrow & & \downarrow \\ A' & \longrightarrow & T(A') \end{array}$$

Namely, given $(A \rightarrow A[x_1, \dots, x_n]/(f_1, \dots, f_n))$ in $S(A)$ we can use the ring map $\varphi : A \rightarrow A'$ to obtain a corresponding element $(A' \rightarrow A'[x_1, \dots, x_n]/(f_1^\varphi, \dots, f_n^\varphi))$ of $S(A')$ where f^φ means the polynomial obtained by applying φ to the coefficients of the polynomial f . Moreover, there is a commutative diagram

$$\begin{array}{ccc} A & \longrightarrow & A[x_1, \dots, x_n]/(f_1, \dots, f_n) \\ \downarrow & & \downarrow \\ A' & \longrightarrow & A'[x_1, \dots, x_n]/(f_1^\varphi, \dots, f_n^\varphi) \end{array}$$

which is a in the category of rings. For $E \subset S(A)$ finite, set $E' = \varphi(E)$ and define $B_E \rightarrow B_{E'}$ in the obvious manner. Taking the colimit gives the desired map $T(A) \rightarrow T(A')$, see Categories, Lemma 14.7.

097U **Lemma 8.4.** *Let A be a ring such that every faithfully flat étale ring map $A \rightarrow B$ has a section. Then the same is true for every quotient ring A/I .*

Proof. Omitted. □

097V **Lemma 8.5.** *Let A be a ring such that every faithfully flat étale ring map $A \rightarrow B$ has a section. Then every local ring of A at a maximal ideal is strictly henselian.*

Proof. Let \mathfrak{m} be a maximal ideal of A . Let $A \rightarrow B$ be an étale ring map and let $\mathfrak{q} \subset B$ be a prime lying over \mathfrak{m} . By the description of the strict henselization $A_{\mathfrak{m}}^{sh}$ in Algebra, Lemma 150.13 it suffices to show that $A_{\mathfrak{m}} = B_{\mathfrak{q}}$. Note that there are finitely many primes $\mathfrak{q} = \mathfrak{q}_1, \mathfrak{q}_2, \dots, \mathfrak{q}_n$ lying over \mathfrak{m} and there are no specializations between them as an étale ring map is quasi-finite, see Algebra, Lemma 141.6. Thus \mathfrak{q}_i is a maximal ideal and we can find $g \in \mathfrak{q}_2 \cap \dots \cap \mathfrak{q}_n, g \notin \mathfrak{q}$ (Algebra, Lemma 14.2). After replacing B by B_g we see that \mathfrak{q} is the only prime of B lying over \mathfrak{m} . The image $U \subset \text{Spec}(A)$ of $\text{Spec}(B) \rightarrow \text{Spec}(A)$ is open (Algebra, Proposition 40.8). Thus the complement $\text{Spec}(A) \setminus U$ is closed and we can find $f \in A, f \notin \mathfrak{p}$ such that $\text{Spec}(A) = U \cup D(f)$. The ring map $A \rightarrow B \times A_f$ is faithfully flat and étale, hence has a section $\sigma : B \times A_f \rightarrow A$ by assumption on A . Observe that σ is étale, hence flat as a map between étale A -algebras (Algebra, Lemma 141.8). Since \mathfrak{q} is the only prime of $B \times A_f$ lying over A we find that $A_{\mathfrak{p}} \rightarrow B_{\mathfrak{q}}$ has a section which is also flat. Thus $A_{\mathfrak{p}} \rightarrow B_{\mathfrak{q}} \rightarrow A_{\mathfrak{p}}$ are flat local ring maps whose composition is the identity. Since a flat local homomorphism of local rings is injective we conclude these maps are isomorphisms as desired. □

097W **Lemma 8.6.** *Let A be a ring such that every faithfully flat étale ring map $A \rightarrow B$ has a section. Let $Z \subset \text{Spec}(A)$ be a closed subscheme of the form $D(f) \cap V(I)$ and let $A \rightarrow A_{\tilde{Z}}$ be as constructed in Lemma 5.1. Then every faithfully flat étale ring map $A_{\tilde{Z}} \rightarrow C$ has a section.*

Proof. There exists an étale ring map $A \rightarrow B'$ such that $C = B' \otimes_A \tilde{A}_{\tilde{Z}}$ as $\tilde{A}_{\tilde{Z}}$ -algebras. The image $U' \subset \text{Spec}(A)$ of $\text{Spec}(B') \rightarrow \text{Spec}(A)$ is open and contains $V(I)$, hence we can find $f \in I$ such that $\text{Spec}(A) = U' \cup D(f)$. Then $A \rightarrow B' \times A_f$ is étale and faithfully flat. By assumption there is a section $B' \times A_f \rightarrow A$. Localizing we obtain the desired section $C \rightarrow \tilde{A}_{\tilde{Z}}$. \square

097X **Lemma 8.7.** *Let $A \rightarrow B$ be a ring map inducing algebraic extensions on residue fields. There exists a commutative diagram*

$$\begin{array}{ccc} B & \longrightarrow & D \\ \uparrow & & \uparrow \\ A & \longrightarrow & C \end{array}$$

with the following properties:

- (1) $A \rightarrow C$ is faithfully flat and ind-étale,
- (2) $B \rightarrow D$ is faithfully flat and ind-étale,
- (3) $\text{Spec}(C)$ is w -local,
- (4) $\text{Spec}(D)$ is w -local,
- (5) $\text{Spec}(D) \rightarrow \text{Spec}(C)$ is w -local,
- (6) the set of closed points of $\text{Spec}(D)$ is the inverse image of the set of closed points of $\text{Spec}(C)$,
- (7) the set of closed points of $\text{Spec}(C)$ surjects onto $\text{Spec}(A)$,
- (8) the set of closed points of $\text{Spec}(D)$ surjects onto $\text{Spec}(B)$,
- (9) for $\mathfrak{m} \subset C$ maximal the local ring $C_{\mathfrak{m}}$ is strictly henselian.

Proof. There is a faithfully flat, ind-Zariski ring map $A \rightarrow A'$ such that $\text{Spec}(A')$ is w -local and such that the set of closed points of $\text{Spec}(A')$ maps onto $\text{Spec}(A)$, see Lemma 5.3. Let $I \subset A'$ be the ideal such that $V(I)$ is the set of closed points of $\text{Spec}(A')$. Choose $A' \rightarrow C'$ as in Lemma 8.1. Note that the local rings $C'_{\mathfrak{m}'}$ at maximal ideals $\mathfrak{m}' \subset C'$ are strictly henselian by Lemma 8.5. We apply Lemma 5.8 to $A' \rightarrow C'$ and $I \subset A'$ to get $C' \rightarrow C$ with $C'/IC' \cong C/IC$. Note that since $A' \rightarrow C'$ is faithfully flat, $\text{Spec}(C'/IC')$ surjects onto the set of closed points of A' and in particular onto $\text{Spec}(A)$. Moreover, as $V(IC) \subset \text{Spec}(C)$ is the set of closed points of C and $C' \rightarrow C$ is ind-Zariski (and identifies local rings) we obtain properties (1), (3), (7), and (9).

Denote $J \subset C$ the ideal such that $V(J)$ is the set of closed points of $\text{Spec}(C)$. Set $D' = B \otimes_A C$. The ring map $C \rightarrow D'$ induces algebraic residue field extensions. Keep in mind that since $V(J) \rightarrow \text{Spec}(A)$ is surjective the map $T = V(JD) \rightarrow \text{Spec}(B)$ is surjective too. Apply Lemma 5.8 to $C \rightarrow D'$ and $J \subset C$ to get $D' \rightarrow D$ with $D'/JD' \cong D/JD$. All of the remaining properties given in the lemma are immediate from the results of Lemma 5.8. \square

9. Weakly étale versus pro-étale

097Y Recall that a ring homomorphism $A \rightarrow B$ is *weakly étale* if $A \rightarrow B$ is flat and $B \otimes_A B \rightarrow B$ is flat. We have proved some properties of such ring maps in More on Algebra, Section 87. In particular, if $A \rightarrow B$ is a local homomorphism, and A is a strictly henselian local rings, then $A = B$, see More on Algebra, Theorem 87.25. Using this theorem and the work we've done above we obtain the following structure theorem for weakly étale ring maps.

097Z **Proposition 9.1.** *Let $A \rightarrow B$ be a weakly étale ring map. Then there exists a faithfully flat, ind-étale ring map $B \rightarrow B'$ such that $A \rightarrow B'$ is ind-étale.*

Proof. The ring map $A \rightarrow B$ induces (separable) algebraic extensions of residue fields, see More on Algebra, Lemma 87.17. Thus we may apply Lemma 8.7 and choose a diagram

$$\begin{array}{ccc} B & \longrightarrow & D \\ \uparrow & & \uparrow \\ A & \longrightarrow & C \end{array}$$

with the properties as listed in the lemma. Note that $C \rightarrow D$ is weakly étale by More on Algebra, Lemma 87.11. Pick a maximal ideal $\mathfrak{m} \subset D$. By construction this lies over a maximal ideal $\mathfrak{m}' \subset C$. By More on Algebra, Theorem 87.25 the ring map $C_{\mathfrak{m}'} \rightarrow D_{\mathfrak{m}}$ is an isomorphism. As every point of $\text{Spec}(C)$ specializes to a closed point we conclude that $C \rightarrow D$ identifies local rings. Thus Proposition 6.6 applies to the ring map $C \rightarrow D$. Pick $D \rightarrow D'$ faithfully flat and ind-Zariski such that $C \rightarrow D'$ is ind-Zariski. Then $B \rightarrow D'$ is a solution to the problem posed in the proposition. \square

10. Constructing w-contractible covers

0980 In this section we construct w-contractible covers of affine schemes.

0981 **Definition 10.1.** Let A be a ring. We say A is *w-contractible* if every faithfully flat weakly-étale ring map $A \rightarrow B$ has a section.

We remark that by Proposition 9.1 an equivalent definition would be to ask that every faithfully flat, ind-étale ring map $A \rightarrow B$ has a section. Here is a key observation that will allow us to construct w-contractible rings.

0982 **Lemma 10.2.** *Let A be a ring. The following are equivalent*

- (1) A is w-contractible,
- (2) every faithfully flat, ind-étale ring map $A \rightarrow B$ has a section, and
- (3) A satisfies
 - (a) $\text{Spec}(A)$ is w-local,
 - (b) $\pi_0(\text{Spec}(A))$ is extremally disconnected, and
 - (c) for every maximal ideal $\mathfrak{m} \subset A$ the local ring $A_{\mathfrak{m}}$ is strictly henselian.

Proof. The equivalence of (1) and (2) follows immediately from Proposition 9.1.

Assume (3)(a), (3)(b), and (3)(c). Let $A \rightarrow B$ be faithfully flat and ind-étale. We will use without further mention the fact that a flat map $A \rightarrow B$ is faithfully flat if and only if every closed point of $\text{Spec}(A)$ is in the image of $\text{Spec}(B) \rightarrow \text{Spec}(A)$. We will show that $A \rightarrow B$ has a section.

Let $I \subset A$ be an ideal such that $V(I) \subset \text{Spec}(A)$ is the set of closed points of $\text{Spec}(A)$. We may replace B by the ring C constructed in Lemma 5.8 for $A \rightarrow B$ and $I \subset A$. Thus we may assume $\text{Spec}(B)$ is w-local such that the set of closed points of $\text{Spec}(B)$ is $V(IB)$. In this case $A \rightarrow B$ identifies local rings by condition (3)(c) as it suffices to check this at maximal ideals of B which lie over maximal ideals of A . Thus $A \rightarrow B$ has a section by Lemma 6.7.

Assume (1) or equivalently (2). We have (3)(c) by Lemma 8.5. Properties (3)(a) and (3)(b) follow from Lemma 6.7. \square

0983 **Proposition 10.3.** *For every ring A there exists a faithfully flat, ind-étale ring map $A \rightarrow D$ such that D is w -contractible.*

Proof. Applying Lemma 8.7 to $\text{id}_A : A \rightarrow A$ we find a faithfully flat, ind-étale ring map $A \rightarrow C$ such that C is w -local and such that every local ring at a maximal ideal of C is strictly henselian. Choose an extremally disconnected space T and a surjective continuous map $T \rightarrow \pi_0(\text{Spec}(C))$, see Topology, Lemma 26.9. Note that T is profinite. Apply Lemma 6.2 to find an ind-Zariski ring map $C \rightarrow D$ such that $\pi_0(\text{Spec}(D)) \rightarrow \pi_0(\text{Spec}(C))$ realizes $T \rightarrow \pi_0(\text{Spec}(C))$ and such that

$$\begin{array}{ccc} \text{Spec}(D) & \longrightarrow & \pi_0(\text{Spec}(D)) \\ \downarrow & & \downarrow \\ \text{Spec}(C) & \longrightarrow & \pi_0(\text{Spec}(C)) \end{array}$$

is cartesian in the category of topological spaces. Note that $\text{Spec}(D)$ is w -local, that $\text{Spec}(D) \rightarrow \text{Spec}(C)$ is w -local, and that the set of closed points of $\text{Spec}(D)$ is the inverse image of the set of closed points of $\text{Spec}(C)$, see Lemma 2.5. Thus it is still true that the local rings of D at its maximal ideals are strictly henselian (as they are isomorphic to the local rings at the corresponding maximal ideals of C). It follows from Lemma 10.2 that D is w -contractible. \square

0984 **Remark 10.4.** Let A be a ring. Let κ be an infinite cardinal bigger or equal than the cardinality of A . Then the cardinality of the ring D constructed in Proposition 10.3 is at most

$$\kappa^{2^{2^{2^\kappa}}}.$$

Namely, the ring map $A \rightarrow D$ is constructed as a composition

$$A \rightarrow A_w = A' \rightarrow C' \rightarrow C \rightarrow D.$$

Here the first three steps of the construction are carried out in the first paragraph of the proof of Lemma 8.7. For the first step we have $|A_w| \leq \kappa$ by Remark 5.4. We have $|C'| \leq \kappa$ by Remark 8.2. Then $|C| \leq \kappa$ because C is a localization of $(C')_w$ (it is constructed from C' by an application of Lemma 5.7 in the proof of Lemma 5.8). Thus C has at most 2^κ maximal ideals. Finally, the ring map $C \rightarrow D$ identifies local rings and the cardinality of the set of maximal ideals of D is at most $2^{2^{2^\kappa}}$ by Topology, Remark 26.10. Since $D \subset \prod_{\mathfrak{m} \subset D} D_{\mathfrak{m}}$ we see that D has at most the size displayed above.

0985 **Lemma 10.5.** *Let $A \rightarrow B$ be a quasi-finite and finitely presented ring map. If the residue fields of A are separably algebraically closed and $\text{Spec}(A)$ is extremally disconnected, then $\text{Spec}(B)$ is extremally disconnected.*

Proof. Set $X = \text{Spec}(A)$ and $Y = \text{Spec}(B)$. Choose a finite partition $X = \coprod X_i$ and $X'_i \rightarrow X_i$ as in Étale Cohomology, Lemma 71.3. Because X is extremally disconnected, every constructible locally closed subset is open and closed, hence we see that X is topologically the disjoint union of the strata X_i . Thus we may replace X by the X_i and assume there exists a surjective finite locally free morphism $X' \rightarrow X$ such that $(X' \times_X Y)_{\text{red}}$ is isomorphic to a finite disjoint union of copies

of X'_{red} . Picture

$$\begin{array}{ccc} \coprod_{i=1, \dots, r} X' & \longrightarrow & Y \\ \downarrow & & \downarrow \\ X' & \longrightarrow & X \end{array}$$

The assumption on the residue fields of A implies that this diagram is a fibre product diagram on underlying sets of points (details omitted). Since X is extremally disconnected and X' is Hausdorff (Lemma 5.6), the continuous map $X' \rightarrow X$ has a continuous section σ . Then $\coprod_{i=1, \dots, r} \sigma(X) \rightarrow Y$ is a bijective continuous map. By Topology, Lemma 17.8 we see that it is a homeomorphism and the proof is done. \square

0986 **Lemma 10.6.** *Let $A \rightarrow B$ be a finite and finitely presented ring map. If A is w-contractible, so is B .*

Proof. We will use the criterion of Lemma 10.2. Set $X = \text{Spec}(A)$ and $Y = \text{Spec}(B)$. As $Y \rightarrow X$ is a finite morphism, we see that the set of closed points Y_0 of Y is the inverse image of the set of closed points X_0 of X . Moreover, every point of Y specializes to a unique point of Y_0 as (a) this is true for X and (b) the map $X \rightarrow Y$ is separated. For every $y \in Y_0$ with image $x \in X_0$ we see that $\mathcal{O}_{Y,y}$ is strictly henselian by Algebra, Lemma 148.4 applied to $\mathcal{O}_{X,x} \rightarrow B \otimes_A \mathcal{O}_{X,x}$. It remains to show that Y_0 is extremally disconnected. To do this we look at $X_0 \times_X Y \rightarrow X_0$ where $X_0 \subset X$ is the reduced induced scheme structure. Note that the underlying topological space of $X_0 \times_X Y$ agrees with Y_0 . Now the desired result follows from Lemma 10.5. \square

0987 **Lemma 10.7.** *Let A be a ring. Let $Z \subset \text{Spec}(A)$ be a closed subset of the form $Z = V(f_1, \dots, f_r)$. Set $B = A_{\tilde{Z}}$, see Lemma 5.1. If A is w-contractible, so is B .*

Proof. Let $A_{\tilde{Z}} \rightarrow B$ be a weakly étale faithfully flat ring map. Consider the ring map

$$A \longrightarrow A_{f_1} \times \dots \times A_{f_r} \times B$$

this is faithful flat and weakly étale. If A is w-contractible, then there is a section σ . Consider the morphism

$$\text{Spec}(A_{\tilde{Z}}) \rightarrow \text{Spec}(A) \xrightarrow{\text{Spec}(\sigma)} \coprod \text{Spec}(A_{f_i}) \amalg \text{Spec}(B)$$

Every point of $Z \subset \text{Spec}(A_{\tilde{Z}})$ maps into the component $\text{Spec}(B)$. Since every point of $\text{Spec}(A_{\tilde{Z}})$ specializes to a point of Z we find a morphism $\text{Spec}(A_{\tilde{Z}}) \rightarrow \text{Spec}(B)$ as desired. \square

11. The pro-étale site

0988 The (small) pro-étale site of a scheme has some remarkable properties. In particular, it has enough w-contractible objects which implies a number of useful consequences for the derived category of abelian sheaves and for inverse systems of sheaves. Thus it is well adapted to deal with some of the intricacies of working with ℓ -adic sheaves.

On the other hand, the pro-étale topology is a bit like the fpqc topology (see Topologies, Section 9) in that the topos of sheaves on the small pro-étale site of a scheme depends on the choice of the underlying category of schemes. Thus we cannot speak of *the* pro-étale topos of a scheme. However, it will be true that the

cohomology groups of a sheaf are unchanged if we enlarge our underlying category of schemes.

Another curiosity is that we define pro-étale coverings using weakly étale morphisms of schemes, see More on Morphisms, Section 53. The reason is that, on the one hand, it is somewhat awkward to define the notion of a pro-étale morphism of schemes, and on the other, Proposition 9.1 assures us that we obtain the same sheaves with the definition that follows.

0989 **Definition 11.1.** Let T be a scheme. A *pro-étale covering* of T is a family of morphisms $\{f_i : T_i \rightarrow T\}_{i \in I}$ of schemes such that each f_i is weakly-étale and such that for every affine open $U \subset T$ there exists $n \geq 0$, a map $a : \{1, \dots, n\} \rightarrow I$ and affine opens $V_j \subset T_{a(j)}$, $j = 1, \dots, n$ with $\bigcup_{j=1}^n f_{a(j)}(V_j) = U$.

To be sure this condition implies that $T = \bigcup f_i(T_i)$. Here is a lemma that will allow us to recognize pro-étale coverings. It will also allow us to reduce many lemmas about pro-étale coverings to the corresponding results for fpqc coverings.

098A **Lemma 11.2.** *Let T be a scheme. Let $\{f_i : T_i \rightarrow T\}_{i \in I}$ be a family of morphisms of schemes with target T . The following are equivalent*

- (1) $\{f_i : T_i \rightarrow T\}_{i \in I}$ is a pro-étale covering,
- (2) each f_i is weakly étale and $\{f_i : T_i \rightarrow T\}_{i \in I}$ is an fpqc covering,
- (3) each f_i is weakly étale and for every affine open $U \subset T$ there exist quasi-compact opens $U_i \subset T_i$ which are almost all empty, such that $U = \bigcup f_i(U_i)$,
- (4) each f_i is weakly étale and there exists an affine open covering $T = \bigcup_{\alpha \in A} U_\alpha$ and for each $\alpha \in A$ there exist $i_{\alpha,1}, \dots, i_{\alpha,n(\alpha)} \in I$ and quasi-compact opens $U_{\alpha,j} \subset T_{i_{\alpha,j}}$ such that $U_\alpha = \bigcup_{j=1, \dots, n(\alpha)} f_{i_{\alpha,j}}(U_{\alpha,j})$.

If T is quasi-separated, these are also equivalent to

- (5) each f_i is weakly étale, and for every $t \in T$ there exist $i_1, \dots, i_n \in I$ and quasi-compact opens $U_j \subset T_{i_j}$ such that $\bigcup_{j=1, \dots, n} f_{i_j}(U_j)$ is a (not necessarily open) neighbourhood of t in T .

Proof. The equivalence of (1) and (2) is immediate from the definitions. Hence the lemma follows from Topologies, Lemma 9.2. \square

098B **Lemma 11.3.** *Any étale covering and any Zariski covering is a pro-étale covering.*

Proof. This follows from the corresponding result for fpqc coverings (Topologies, Lemma 9.6), Lemma 11.2, and the fact that an étale morphism is a weakly étale morphism, see More on Morphisms, Lemma 53.9. \square

098C **Lemma 11.4.** *Let T be a scheme.*

- (1) If $T' \rightarrow T$ is an isomorphism then $\{T' \rightarrow T\}$ is a pro-étale covering of T .
- (2) If $\{T_i \rightarrow T\}_{i \in I}$ is a pro-étale covering and for each i we have a pro-étale covering $\{T_{ij} \rightarrow T_i\}_{j \in J_i}$, then $\{T_{ij} \rightarrow T\}_{i \in I, j \in J_i}$ is a pro-étale covering.
- (3) If $\{T_i \rightarrow T\}_{i \in I}$ is a pro-étale covering and $T' \rightarrow T$ is a morphism of schemes then $\{T' \times_T T_i \rightarrow T'\}_{i \in I}$ is a pro-étale covering.

Proof. This follows from the fact that composition and base changes of weakly étale morphisms are weakly étale (More on Morphisms, Lemmas 53.5 and 53.6), Lemma 11.2, and the corresponding results for fpqc coverings, see Topologies, Lemma 9.7. \square

098D **Lemma 11.5.** *Let T be an affine scheme. Let $\{T_i \rightarrow T\}_{i \in I}$ be a pro-étale covering of T . Then there exists a pro-étale covering $\{U_j \rightarrow T\}_{j=1, \dots, n}$ which is a refinement of $\{T_i \rightarrow T\}_{i \in I}$ such that each U_j is an affine scheme. Moreover, we may choose each U_j to be open affine in one of the T_i .*

Proof. This follows directly from the definition. □

Thus we define the corresponding standard coverings of affines as follows.

098E **Definition 11.6.** Let T be an affine scheme. A *standard pro-étale covering* of T is a family $\{f_i : T_i \rightarrow T\}_{i=1, \dots, n}$ where each T_j is affine, each f_i is weakly étale, and $T = \bigcup f_i(T_i)$.

We interrupt the discussion for an explanation of the notion of w-contractible rings in terms of pro-étale coverings.

098F **Lemma 11.7.** *Let $T = \text{Spec}(A)$ be an affine scheme. The following are equivalent*

- (1) *A is w-contractible, and*
- (2) *every pro-étale covering of T can be refined by a Zariski covering of the form $T = \coprod_{i=1, \dots, n} U_i$.*

Proof. Assume A is w-contractible. By Lemma 11.5 it suffices to prove we can refine every standard pro-étale covering $\{f_i : T_i \rightarrow T\}_{i=1, \dots, n}$ by a Zariski covering of T . The morphism $\coprod T_i \rightarrow T$ is a surjective weakly étale morphism of affine schemes. Hence by Definition 10.1 there exists a morphism $\sigma : T \rightarrow \coprod T_i$ over T . Then the Zariski covering $T = \coprod \sigma^{-1}(T_i)$ refines $\{f_i : T_i \rightarrow T\}$.

Conversely, assume (2). If $A \rightarrow B$ is faithfully flat and weakly étale, then $\{\text{Spec}(B) \rightarrow T\}$ is a pro-étale covering. Hence there exists a Zariski covering $T = \coprod U_i$ and morphisms $U_i \rightarrow \text{Spec}(B)$ over T . Since $T = \coprod U_i$ we obtain $T \rightarrow \text{Spec}(B)$, i.e., an A -algebra map $B \rightarrow A$. This means A is w-contractible. □

We follow the general outline given in Topologies, Section 2 for constructing the big pro-étale site we will be working with. However, because we need a bit larger rings to accommodate for the size of certain constructions we modify the constructions slightly.

098G **Definition 11.8.** A *big pro-étale site* is any site $Sch_{\text{pro-étale}}$ as in Sites, Definition 6.2 constructed as follows:

- (1) Choose any set of schemes S_0 , and any set of pro-étale coverings Cov_0 among these schemes.
- (2) Change the function *Bound* of Sets, Equation (9.1.1) into

$$\text{Bound}(\kappa) = \max\{\kappa^{2^{2^{2^\kappa}}}, \kappa^{\aleph_0}, \kappa^+\}.$$

- (3) As underlying category take any category Sch_α constructed as in Sets, Lemma 9.2 starting with the set S_0 and the function *Bound*.
- (4) Choose any set of coverings as in Sets, Lemma 11.1 starting with the category Sch_α and the class of pro-étale coverings, and the set Cov_0 chosen above.

See the remarks following Topologies, Definition 3.5 for motivation and explanation regarding the definition of big sites.

Before we continue with the introduction of the big and small pro-étale sites of a scheme, let us point out that (1) our category contains many weakly contractible objects, and (2) the topology on a big pro-étale site $Sch_{pro\text{-}\acute{e}tale}$ is in some sense induced from the pro-étale topology on the category of all schemes.

098H **Lemma 11.9.** *Let $Sch_{pro\text{-}\acute{e}tale}$ be a big pro-étale site as in Definition 11.8. Let $T = \text{Spec}(A)$ be an affine object of $Sch_{pro\text{-}\acute{e}tale}$. If A is w-contractible, then T is a weakly contractible (Sites, Definition 39.2) object of $Sch_{pro\text{-}\acute{e}tale}$.*

Proof. Let $\mathcal{F} \rightarrow \mathcal{G}$ be a surjection of sheaves on $Sch_{pro\text{-}\acute{e}tale}$. Let $s \in \mathcal{G}(T)$. We have to show that s is in the image of $\mathcal{F}(T) \rightarrow \mathcal{G}(T)$. We can find a covering $\{T_i \rightarrow T\}$ of $Sch_{pro\text{-}\acute{e}tale}$ such that s lifts to a section of \mathcal{F} over T_i (Sites, Definition 11.1). By Lemma 11.7 we can refine $\{T_i \rightarrow T\}$ by a Zariski covering of the form $T = \coprod_{j=1, \dots, m} V_j$. Hence we get $t_j \in \mathcal{F}(U_j)$ mapping to $s|_{U_j}$. Since Zariski coverings are coverings in $Sch_{pro\text{-}\acute{e}tale}$ (Lemma 11.3) we conclude that $\mathcal{F}(T) = \prod \mathcal{F}(U_j)$. Thus, taking $t = (t_1, \dots, t_m) \in \mathcal{F}(T)$ is a section mapping to s . \square

098I **Lemma 11.10.** *Let $Sch_{pro\text{-}\acute{e}tale}$ be a big pro-étale site as in Definition 11.8. For every object T of $Sch_{pro\text{-}\acute{e}tale}$ there exists a covering $\{T_i \rightarrow T\}$ in $Sch_{pro\text{-}\acute{e}tale}$ with each T_i affine and the spectrum of a w-contractible ring. In particular, T_i is weakly contractible in $Sch_{pro\text{-}\acute{e}tale}$.*

Proof. For those readers who do not care about set-theoretical issues this lemma is a trivial consequence of Lemma 11.9 and Proposition 10.3. Here are the details. Choose an affine open covering $T = \bigcup U_i$. Write $U_i = \text{Spec}(A_i)$. Choose faithfully flat, ind-étale ring maps $A_i \rightarrow D_i$ such that D_i is w-contractible as in Proposition 10.3. The family of morphisms $\{\text{Spec}(D_i) \rightarrow T\}$ is a pro-étale covering. If we can show that $\text{Spec}(D_i)$ is isomorphic to an object, say T_i , of $Sch_{pro\text{-}\acute{e}tale}$, then $\{T_i \rightarrow T\}$ will be combinatorially equivalent to a covering of $Sch_{pro\text{-}\acute{e}tale}$ by the construction of $Sch_{pro\text{-}\acute{e}tale}$ in Definition 11.8 and more precisely the application of Sets, Lemma 11.1 in the last step. To prove $\text{Spec}(D_i)$ is isomorphic to an object of $Sch_{pro\text{-}\acute{e}tale}$, it suffices to prove that $|D_i| \leq \text{Bound}(\text{size}(T))$ by the construction of $Sch_{pro\text{-}\acute{e}tale}$ in Definition 11.8 and more precisely the application of Sets, Lemma 9.2 in step (3). Since $|A_i| \leq \text{size}(U_i) \leq \text{size}(T)$ by Sets, Lemmas 9.4 and 9.7 we get $|D_i| \leq \kappa^{2^{2^{\kappa}}}$ where $\kappa = \text{size}(T)$ by Remark 10.4. Thus by our choice of the function Bound in Definition 11.8 we win. \square

098J **Lemma 11.11.** *Let $Sch_{pro\text{-}\acute{e}tale}$ be a big pro-étale site as in Definition 11.8. Let $T \in \text{Ob}(Sch_{pro\text{-}\acute{e}tale})$. Let $\{T_i \rightarrow T\}_{i \in I}$ be an arbitrary pro-étale covering of T . There exists a covering $\{U_j \rightarrow T\}_{j \in J}$ of T in the site $Sch_{pro\text{-}\acute{e}tale}$ which refines $\{T_i \rightarrow T\}_{i \in I}$.*

Proof. Namely, we first let $\{V_k \rightarrow T\}$ be a covering as in Lemma 11.10. Then the pro-étale coverings $\{T_i \times_T V_k \rightarrow V_k\}$ can be refined by a finite disjoint open covering $V_k = V_{k,1} \amalg \dots \amalg V_{k,n_k}$, see Lemma 11.7. Then $\{V_{k,i} \rightarrow T\}$ is a covering of $Sch_{pro\text{-}\acute{e}tale}$ which refines $\{T_i \rightarrow T\}_{i \in I}$. \square

098K **Definition 11.12.** Let S be a scheme. Let $Sch_{pro\text{-}\acute{e}tale}$ be a big pro-étale site containing S .

- (1) The big pro-étale site of S , denoted $(Sch/S)_{pro\text{-}\acute{e}tale}$, is the site $Sch_{pro\text{-}\acute{e}tale}/S$ introduced in Sites, Section 24.

- (2) The *small pro-étale site of S* , which we denote $S_{\text{pro-étale}}$, is the full subcategory of $(\text{Sch}/S)_{\text{pro-étale}}$ whose objects are those U/S such that $U \rightarrow S$ is weakly étale. A covering of $S_{\text{pro-étale}}$ is any covering $\{U_i \rightarrow U\}$ of $(\text{Sch}/S)_{\text{pro-étale}}$ with $U \in \text{Ob}(S_{\text{pro-étale}})$.
- (3) The *big affine pro-étale site of S* , denoted $(\text{Aff}/S)_{\text{pro-étale}}$, is the full subcategory of $(\text{Sch}/S)_{\text{pro-étale}}$ whose objects are affine U/S . A covering of $(\text{Aff}/S)_{\text{pro-étale}}$ is any covering $\{U_i \rightarrow U\}$ of $(\text{Sch}/S)_{\text{pro-étale}}$ which is a standard pro-étale covering.

It is not completely clear that the small pro-étale site and the big affine pro-étale site are sites. We check this now.

098L **Lemma 11.13.** *Let S be a scheme. Let $\text{Sch}_{\text{pro-étale}}$ be a big pro-étale site containing S . Both $S_{\text{pro-étale}}$ and $(\text{Aff}/S)_{\text{pro-étale}}$ are sites.*

Proof. Let us show that $S_{\text{pro-étale}}$ is a site. It is a category with a given set of families of morphisms with fixed target. Thus we have to show properties (1), (2) and (3) of Sites, Definition 6.2. Since $(\text{Sch}/S)_{\text{pro-étale}}$ is a site, it suffices to prove that given any covering $\{U_i \rightarrow U\}$ of $(\text{Sch}/S)_{\text{pro-étale}}$ with $U \in \text{Ob}(S_{\text{pro-étale}})$ we also have $U_i \in \text{Ob}(S_{\text{pro-étale}})$. This follows from the definitions as the composition of weakly étale morphisms is weakly étale.

To show that $(\text{Aff}/S)_{\text{pro-étale}}$ is a site, reasoning as above, it suffices to show that the collection of standard pro-étale coverings of affines satisfies properties (1), (2) and (3) of Sites, Definition 6.2. This follows from Lemma 11.2 and the corresponding result for standard fpqc coverings (Topologies, Lemma 9.10). \square

098M **Lemma 11.14.** *Let S be a scheme. Let $\text{Sch}_{\text{pro-étale}}$ be a big pro-étale site containing S . Let Sch be the category of all schemes.*

- (1) *The categories $\text{Sch}_{\text{pro-étale}}$, $(\text{Sch}/S)_{\text{pro-étale}}$, $S_{\text{pro-étale}}$, and $(\text{Aff}/S)_{\text{pro-étale}}$ have fibre products agreeing with fibre products in Sch .*
- (2) *The categories $\text{Sch}_{\text{pro-étale}}$, $(\text{Sch}/S)_{\text{pro-étale}}$, $S_{\text{pro-étale}}$ have equalizers agreeing with equalizers in Sch .*
- (3) *The categories $(\text{Sch}/S)_{\text{pro-étale}}$, and $S_{\text{pro-étale}}$ both have a final object, namely S/S .*
- (4) *The category $\text{Sch}_{\text{pro-étale}}$ has a final object agreeing with the final object of Sch , namely $\text{Spec}(\mathbf{Z})$.*

Proof. The category $\text{Sch}_{\text{pro-étale}}$ contains $\text{Spec}(\mathbf{Z})$ and is closed under products and fibre products by construction, see Sets, Lemma 9.9. Suppose we have $U \rightarrow S$, $V \rightarrow U$, $W \rightarrow U$ morphisms of schemes with $U, V, W \in \text{Ob}(\text{Sch}_{\text{pro-étale}})$. The fibre product $V \times_U W$ in $\text{Sch}_{\text{pro-étale}}$ is a fibre product in Sch and is the fibre product of V/S with W/S over U/S in the category of all schemes over S , and hence also a fibre product in $(\text{Sch}/S)_{\text{pro-étale}}$. This proves the result for $(\text{Sch}/S)_{\text{pro-étale}}$. If $U \rightarrow S$, $V \rightarrow U$ and $W \rightarrow U$ are weakly étale then so is $V \times_U W \rightarrow S$ (see More on Morphisms, Section 53) and hence we get fibre products for $S_{\text{pro-étale}}$. If U, V, W are affine, so is $V \times_U W$ and hence we get fibre products for $(\text{Aff}/S)_{\text{pro-étale}}$.

Let $a, b : U \rightarrow V$ be two morphisms in $\text{Sch}_{\text{pro-étale}}$. In this case the equalizer of a and b (in the category of schemes) is

$$V \times_{\Delta_{V/\text{Spec}(\mathbf{Z}), V \times_{\text{Spec}(\mathbf{Z})} V, (a, b)}} (U \times_{\text{Spec}(\mathbf{Z})} U)$$

which is an object of $Sch_{pro\text{-}\acute{e}tale}$ by what we saw above. Thus $Sch_{pro\text{-}\acute{e}tale}$ has equalizers. If a and b are morphisms over S , then the equalizer (in the category of schemes) is also given by

$$V \times_{\Delta_{V/S}, V \times_S V, (a,b)} (U \times_S U)$$

hence we see that $(Sch/S)_{pro\text{-}\acute{e}tale}$ has equalizers. Moreover, if U and V are weakly étale over S , then so is the equalizer above as a fibre product of schemes weakly étale over S . Thus $S_{pro\text{-}\acute{e}tale}$ has equalizers. The statements on final objects is clear. \square

Next, we check that the big affine pro-étale site defines the same topos as the big pro-étale site.

098N **Lemma 11.15.** *Let S be a scheme. Let $Sch_{pro\text{-}\acute{e}tale}$ be a big pro-étale site containing S . The functor $(Aff/S)_{pro\text{-}\acute{e}tale} \rightarrow (Sch/S)_{pro\text{-}\acute{e}tale}$ is a special cocontinuous functor. Hence it induces an equivalence of topoi from $Sh((Aff/S)_{pro\text{-}\acute{e}tale})$ to $Sh((Sch/S)_{pro\text{-}\acute{e}tale})$.*

Proof. The notion of a special cocontinuous functor is introduced in Sites, Definition 28.2. Thus we have to verify assumptions (1) – (5) of Sites, Lemma 28.1. Denote the inclusion functor $u : (Aff/S)_{pro\text{-}\acute{e}tale} \rightarrow (Sch/S)_{pro\text{-}\acute{e}tale}$. Being cocontinuous just means that any pro-étale covering of T/S , T affine, can be refined by a standard pro-étale covering of T . This is the content of Lemma 11.5. Hence (1) holds. We see u is continuous simply because a standard pro-étale covering is a pro-étale covering. Hence (2) holds. Parts (3) and (4) follow immediately from the fact that u is fully faithful. And finally condition (5) follows from the fact that every scheme has an affine open covering. \square

098P **Lemma 11.16.** *Let $Sch_{pro\text{-}\acute{e}tale}$ be a big pro-étale site. Let $f : T \rightarrow S$ be a morphism in $Sch_{pro\text{-}\acute{e}tale}$. The functor $T_{pro\text{-}\acute{e}tale} \rightarrow (Sch/S)_{pro\text{-}\acute{e}tale}$ is cocontinuous and induces a morphism of topoi*

$$i_f : Sh(T_{pro\text{-}\acute{e}tale}) \longrightarrow Sh((Sch/S)_{pro\text{-}\acute{e}tale})$$

For a sheaf \mathcal{G} on $(Sch/S)_{pro\text{-}\acute{e}tale}$ we have the formula $(i_f^{-1}\mathcal{G})(U/T) = \mathcal{G}(U/S)$. The functor i_f^{-1} also has a left adjoint $i_{f,!}$ which commutes with fibre products and equalizers.

Proof. Denote the functor $u : T_{pro\text{-}\acute{e}tale} \rightarrow (Sch/S)_{pro\text{-}\acute{e}tale}$. In other words, given a weakly étale morphism $j : U \rightarrow T$ corresponding to an object of $T_{pro\text{-}\acute{e}tale}$ we set $u(U \rightarrow T) = (f \circ j : U \rightarrow S)$. This functor commutes with fibre products, see Lemma 11.14. Moreover, $T_{pro\text{-}\acute{e}tale}$ has equalizers and u commutes with them by Lemma 11.14. It is clearly cocontinuous. It is also continuous as u transforms coverings to coverings and commutes with fibre products. Hence the lemma follows from Sites, Lemmas 20.5 and 20.6. \square

098Q **Lemma 11.17.** *Let S be a scheme. Let $Sch_{pro\text{-}\acute{e}tale}$ be a big pro-étale site containing S . The inclusion functor $S_{pro\text{-}\acute{e}tale} \rightarrow (Sch/S)_{pro\text{-}\acute{e}tale}$ satisfies the hypotheses of Sites, Lemma 20.8 and hence induces a morphism of sites*

$$\pi_S : (Sch/S)_{pro\text{-}\acute{e}tale} \longrightarrow S_{pro\text{-}\acute{e}tale}$$

and a morphism of topoi

$$i_S : Sh(S_{pro\text{-}\acute{e}tale}) \longrightarrow Sh((Sch/S)_{pro\text{-}\acute{e}tale})$$

such that $\pi_S \circ i_S = id$. Moreover, $i_S = i_{id_S}$ with i_{id_S} as in Lemma 11.16. In particular the functor $i_S^{-1} = \pi_{S,*}$ is described by the rule $i_S^{-1}(\mathcal{G})(U/S) = \mathcal{G}(U/S)$.

Proof. In this case the functor $u : S_{pro\text{-}\acute{e}tale} \rightarrow (Sch/S)_{pro\text{-}\acute{e}tale}$, in addition to the properties seen in the proof of Lemma 11.16 above, also is fully faithful and transforms the final object into the final object. The lemma follows from Sites, Lemma 20.8. \square

098R **Definition 11.18.** In the situation of Lemma 11.17 the functor $i_S^{-1} = \pi_{S,*}$ is often called the *restriction to the small pro-étale site*, and for a sheaf \mathcal{F} on the big pro-étale site we denote $\mathcal{F}|_{S_{pro\text{-}\acute{e}tale}}$ this restriction.

With this notation in place we have for a sheaf \mathcal{F} on the big site and a sheaf \mathcal{G} on the big site that

$$\begin{aligned} Mor_{Sh(S_{pro\text{-}\acute{e}tale})}(\mathcal{F}|_{S_{pro\text{-}\acute{e}tale}}, \mathcal{G}) &= Mor_{Sh((Sch/S)_{pro\text{-}\acute{e}tale})}(\mathcal{F}, i_{S,*}\mathcal{G}) \\ Mor_{Sh(S_{pro\text{-}\acute{e}tale})}(\mathcal{G}, \mathcal{F}|_{S_{pro\text{-}\acute{e}tale}}) &= Mor_{Sh((Sch/S)_{pro\text{-}\acute{e}tale})}(\pi_S^{-1}\mathcal{G}, \mathcal{F}) \end{aligned}$$

Moreover, we have $(i_{S,*}\mathcal{G})|_{S_{pro\text{-}\acute{e}tale}} = \mathcal{G}$ and we have $(\pi_S^{-1}\mathcal{G})|_{S_{pro\text{-}\acute{e}tale}} = \mathcal{G}$.

098S **Lemma 11.19.** Let $Sch_{pro\text{-}\acute{e}tale}$ be a big pro-étale site. Let $f : T \rightarrow S$ be a morphism in $Sch_{pro\text{-}\acute{e}tale}$. The functor

$$u : (Sch/T)_{pro\text{-}\acute{e}tale} \longrightarrow (Sch/S)_{pro\text{-}\acute{e}tale}, \quad V/T \longmapsto V/S$$

is cocontinuous, and has a continuous right adjoint

$$v : (Sch/S)_{pro\text{-}\acute{e}tale} \longrightarrow (Sch/T)_{pro\text{-}\acute{e}tale}, \quad (U \rightarrow S) \longmapsto (U \times_S T \rightarrow T).$$

They induce the same morphism of topoi

$$f_{big} : Sh((Sch/T)_{pro\text{-}\acute{e}tale}) \longrightarrow Sh((Sch/S)_{pro\text{-}\acute{e}tale})$$

We have $f_{big}^{-1}(\mathcal{G})(U/T) = \mathcal{G}(U/S)$. We have $f_{big,*}(\mathcal{F})(U/S) = \mathcal{F}(U \times_S T/T)$. Also, f_{big}^{-1} has a left adjoint $f_{big}!$ which commutes with fibre products and equalizers.

Proof. The functor u is cocontinuous, continuous, and commutes with fibre products and equalizers (details omitted; compare with proof of Lemma 11.16). Hence Sites, Lemmas 20.5 and 20.6 apply and we deduce the formula for f_{big}^{-1} and the existence of $f_{big}!$. Moreover, the functor v is a right adjoint because given U/T and V/S we have $Mor_S(u(U), V) = Mor_T(U, V \times_S T)$ as desired. Thus we may apply Sites, Lemmas 21.1 and 21.2 to get the formula for $f_{big,*}$. \square

098T **Lemma 11.20.** Let $Sch_{pro\text{-}\acute{e}tale}$ be a big pro-étale site. Let $f : T \rightarrow S$ be a morphism in $Sch_{pro\text{-}\acute{e}tale}$.

- (1) We have $i_f = f_{big} \circ i_T$ with i_f as in Lemma 11.16 and i_T as in Lemma 11.17.
- (2) The functor $S_{pro\text{-}\acute{e}tale} \rightarrow T_{pro\text{-}\acute{e}tale}$, $(U \rightarrow S) \mapsto (U \times_S T \rightarrow T)$ is continuous and induces a morphism of topoi

$$f_{small} : Sh(T_{pro\text{-}\acute{e}tale}) \longrightarrow Sh(S_{pro\text{-}\acute{e}tale}).$$

We have $f_{small,*}(\mathcal{F})(U/S) = \mathcal{F}(U \times_S T/T)$.

(3) We have a commutative diagram of morphisms of sites

$$\begin{array}{ccc} T_{\text{pro-étale}} & \xleftarrow{\pi_T} & (\text{Sch}/T)_{\text{pro-étale}} \\ f_{\text{small}} \downarrow & & \downarrow f_{\text{big}} \\ S_{\text{pro-étale}} & \xleftarrow{\pi_S} & (\text{Sch}/S)_{\text{pro-étale}} \end{array}$$

so that $f_{\text{small}} \circ \pi_T = \pi_S \circ f_{\text{big}}$ as morphisms of topoi.

(4) We have $f_{\text{small}} = \pi_S \circ f_{\text{big}} \circ i_T = \pi_S \circ i_f$.

Proof. The equality $i_f = f_{\text{big}} \circ i_T$ follows from the equality $i_f^{-1} = i_T^{-1} \circ f_{\text{big}}^{-1}$ which is clear from the descriptions of these functors above. Thus we see (1).

The functor $u : S_{\text{pro-étale}} \rightarrow T_{\text{pro-étale}}$, $u(U \rightarrow S) = (U \times_S T \rightarrow T)$ transforms coverings into coverings and commutes with fibre products, see Lemmas 11.4 and 11.14. Moreover, both $S_{\text{pro-étale}}$, $T_{\text{pro-étale}}$ have final objects, namely S/S and T/T and $u(S/S) = T/T$. Hence by Sites, Proposition 14.6 the functor u corresponds to a morphism of sites $T_{\text{pro-étale}} \rightarrow S_{\text{pro-étale}}$. This in turn gives rise to the morphism of topoi, see Sites, Lemma 15.2. The description of the pushforward is clear from these references.

Part (3) follows because π_S and π_T are given by the inclusion functors and f_{small} and f_{big} by the base change functors $U \mapsto U \times_S T$.

Statement (4) follows from (3) by precomposing with i_T . \square

In the situation of the lemma, using the terminology of Definition 11.18 we have: for \mathcal{F} a sheaf on the big pro-étale site of T

$$(f_{\text{big},*} \mathcal{F})|_{S_{\text{pro-étale}}} = f_{\text{small},*}(\mathcal{F}|_{T_{\text{pro-étale}}}),$$

This equality is clear from the commutativity of the diagram of sites of the lemma, since restriction to the small pro-étale site of T , resp. S is given by $\pi_{T,*}$, resp. $\pi_{S,*}$. A similar formula involving pullbacks and restrictions is false.

098U **Lemma 11.21.** *Given schemes X, Y, Z in $\text{Sch}_{\text{pro-étale}}$ and morphisms $f : X \rightarrow Y, g : Y \rightarrow Z$ we have $g_{\text{big}} \circ f_{\text{big}} = (g \circ f)_{\text{big}}$ and $g_{\text{small}} \circ f_{\text{small}} = (g \circ f)_{\text{small}}$.*

Proof. This follows from the simple description of pushforward and pullback for the functors on the big sites from Lemma 11.19. For the functors on the small sites this follows from the description of the pushforward functors in Lemma 11.20. \square

We can think about a sheaf on the big pro-étale site of S as a collection of sheaves on the small pro-étale site on schemes over S .

098V **Lemma 11.22.** *Let S be a scheme contained in a big pro-étale site $\text{Sch}_{\text{pro-étale}}$. A sheaf \mathcal{F} on the big pro-étale site $(\text{Sch}/S)_{\text{pro-étale}}$ is given by the following data:*

- (1) for every $T/S \in \text{Ob}((\text{Sch}/S)_{\text{pro-étale}})$ a sheaf \mathcal{F}_T on $T_{\text{pro-étale}}$,
- (2) for every $f : T' \rightarrow T$ in $(\text{Sch}/S)_{\text{pro-étale}}$ a map $c_f : f_{\text{small}}^{-1} \mathcal{F}_T \rightarrow \mathcal{F}_{T'}$.

These data are subject to the following conditions:

- (a) given any $f : T' \rightarrow T$ and $g : T'' \rightarrow T'$ in $(\text{Sch}/S)_{\text{pro-étale}}$ the composition $g_{\text{small}}^{-1} c_f \circ c_g$ is equal to $c_{f \circ g}$, and
- (b) if $f : T' \rightarrow T$ in $(\text{Sch}/S)_{\text{pro-étale}}$ is weakly étale then c_f is an isomorphism.

Proof. Identical to the proof of Topologies, Lemma 4.19. \square

098W **Lemma 11.23.** *Let S be a scheme. Let $S_{\text{affine,pro-étale}}$ denote the full subcategory of $S_{\text{pro-étale}}$ consisting of affine objects. A covering of $S_{\text{affine,pro-étale}}$ will be a standard étale covering, see Definition 11.6. Then restriction*

$$\mathcal{F} \longmapsto \mathcal{F}|_{S_{\text{affine,étale}}}$$

defines an equivalence of topoi $Sh(S_{\text{pro-étale}}) \cong Sh(S_{\text{affine,pro-étale}})$.

Proof. This you can show directly from the definitions, and is a good exercise. But it also follows immediately from Sites, Lemma 28.1 by checking that the inclusion functor $S_{\text{affine,pro-étale}} \rightarrow S_{\text{pro-étale}}$ is a special cocontinuous functor (see Sites, Definition 28.2). \square

098X **Lemma 11.24.** *Let S be an affine scheme. Let S_{app} denote the full subcategory of $S_{\text{pro-étale}}$ consisting of affine objects U such that $\mathcal{O}(S) \rightarrow \mathcal{O}(U)$ is ind-étale. A covering of S_{app} will be a standard pro-étale covering, see Definition 11.6. Then restriction*

$$\mathcal{F} \longmapsto \mathcal{F}|_{S_{\text{app}}}$$

defines an equivalence of topoi $Sh(S_{\text{pro-étale}}) \cong Sh(S_{\text{app}})$.

Proof. By Lemma 11.23 we may replace $S_{\text{pro-étale}}$ by $S_{\text{affine,pro-étale}}$. The lemma follows from Sites, Lemma 28.1 by checking that the inclusion functor $S_{\text{app}} \rightarrow S_{\text{affine,pro-étale}}$ is a special cocontinuous functor, see Sites, Definition 28.2. The conditions of Sites, Lemma 28.1 follow immediately from the definition and the facts (a) any object U of $S_{\text{affine,pro-étale}}$ has a covering $\{V \rightarrow U\}$ with V ind-étale over X (Proposition 9.1) and (b) the functor u is fully faithful. \square

Next we show that cohomology of sheaves is independent of the choice of a partial universe. Namely, the functor g_* of the lemma below is an embedding of pro-étale topoi which does not change cohomology.

098Y **Lemma 11.25.** *Let S be a scheme. Let $S_{\text{pro-étale}} \subset S'_{\text{pro-étale}}$ be two small pro-étale sites of S as constructed in Definition 11.12. Then the inclusion functor satisfies the assumptions of Sites, Lemma 20.8. Hence there exist morphisms of topoi*

$$Sh(S_{\text{pro-étale}}) \xrightarrow{g} Sh(S'_{\text{pro-étale}}) \xrightarrow{f} Sh(S_{\text{pro-étale}})$$

whose composition is isomorphic to the identity and with $f_ = g^{-1}$. Moreover,*

- (1) *for $\mathcal{F}' \in Ab(S'_{\text{pro-étale}})$ we have $H^p(S'_{\text{pro-étale}}, \mathcal{F}') = H^p(S_{\text{pro-étale}}, g^{-1}\mathcal{F}')$,*
- (2) *for $\mathcal{F} \in Ab(S_{\text{pro-étale}})$ we have*

$$H^p(S_{\text{pro-étale}}, \mathcal{F}) = H^p(S'_{\text{pro-étale}}, g_*\mathcal{F}) = H^p(S'_{\text{pro-étale}}, f^{-1}\mathcal{F}).$$

Proof. The inclusion functor is fully faithful and continuous. We have seen that $S_{\text{pro-étale}}$ and $S'_{\text{pro-étale}}$ have fibre products and final objects and that our functor commutes with these (Lemma 11.14). It follows from Lemma 11.11 that the inclusion functor is cocontinuous. Hence the existence of f and g follows from Sites, Lemma 20.8. The equality in (1) is Cohomology on Sites, Lemma 8.2. Part (2) follows from (1) as $\mathcal{F} = g^{-1}g_*\mathcal{F} = g^{-1}f^{-1}\mathcal{F}$. \square

098Z **Lemma 11.26.** *Let S be a scheme. The topology on each of the pro-étale sites $S_{\text{pro-étale}}$, $(Sch/S)_{\text{pro-étale}}$, $S_{\text{affine,pro-étale}}$, and $(Aff/S)_{\text{pro-étale}}$ is subcanonical.*

Proof. Combine Lemma 11.2 and Descent, Lemma 10.3. \square

0990 **Lemma 11.27.** *Let S be a scheme. The pro-étale sites $S_{pro\text{-}\acute{e}tale}$, $(Sch/S)_{pro\text{-}\acute{e}tale}$, $S_{affine,pro\text{-}\acute{e}tale}$, and $(Aff/S)_{pro\text{-}\acute{e}tale}$ and if S is affine S_{app} have enough quasi-compact, weakly contractible objects, see Sites, Definition 39.2.*

Proof. Follows immediately from Lemma 11.10. □

12. Points of the pro-étale site

0991 We first apply Deligne’s criterion to show that there are enough points.

0992 **Lemma 12.1.** *Let S be a scheme. The pro-étale sites $S_{pro\text{-}\acute{e}tale}$, $(Sch/S)_{pro\text{-}\acute{e}tale}$, $S_{affine,pro\text{-}\acute{e}tale}$, and $(Aff/S)_{pro\text{-}\acute{e}tale}$ have enough points.*

Proof. The big topos is equivalent to the topos defined by $(Aff/S)_{pro\text{-}\acute{e}tale}$, see Lemma 11.15. The topos of sheaves on $S_{pro\text{-}\acute{e}tale}$ is equivalent to the topos associated to $S_{affine,pro\text{-}\acute{e}tale}$, see Lemma 11.23. The result for the sites $(Aff/S)_{pro\text{-}\acute{e}tale}$ and $S_{affine,pro\text{-}\acute{e}tale}$ follows immediately from Deligne’s result Sites, Lemma 38.4. □

Let S be a scheme. Let $\bar{s} : \text{Spec}(k) \rightarrow S$ be a geometric point. We define a *pro-étale neighbourhood* of \bar{s} to be a commutative diagram

$$\begin{array}{ccc} \text{Spec}(k) & \xrightarrow{\bar{u}} & U \\ & \searrow \bar{s} & \downarrow \\ & & S \end{array}$$

with $U \rightarrow S$ weakly étale. In exactly the same manner as in Étale Cohomology, Lemma 29.4 one shows that the category of pro-étale neighbourhoods of \bar{s} is cofiltered. Moreover, if (U, \bar{u}) is a pro-étale neighbourhood, and if $\{U_i \rightarrow U\}$ is a pro-étale covering, then there exists an i and a lift of \bar{u} to a geometric point \bar{u}_i of U_i . For \mathcal{F} in $Sh(S_{pro\text{-}\acute{e}tale})$ define the *stalk of \mathcal{F} at \bar{s}* by the formula

$$\mathcal{F}_{\bar{s}} = \text{colim}_{(U, \bar{u})} \mathcal{F}(U)$$

where the colimit is over all pro-étale neighbourhoods (U, \bar{u}) of \bar{s} with $U \in \text{Ob}(S_{pro\text{-}\acute{e}tale})$. A formal argument using the facts above shows the functor $\mathcal{F} \mapsto \mathcal{F}_{\bar{s}}$ defines a point of the topos $Sh(S_{pro\text{-}\acute{e}tale})$: it is an exact functor which commutes with arbitrary colimits. In fact, this functor has another description.

0993 **Lemma 12.2.** *In the situation above the scheme $\text{Spec}(\mathcal{O}_{S, \bar{s}}^{sh})$ is an object of $X_{pro\text{-}\acute{e}tale}$ and there is a canonical isomorphism*

$$\mathcal{F}(\text{Spec}(\mathcal{O}_{S, \bar{s}}^{sh})) = \mathcal{F}_{\bar{s}}$$

functorial in \mathcal{F} .

Proof. The first statement is clear from the construction of the strict henselization as a filtered colimit of étale algebras over S , or by the characterization of weakly étale morphisms of More on Morphisms, Lemma 53.11. The second statement follows as by Olivier’s theorem (More on Algebra, Theorem 87.25) the scheme $\text{Spec}(\mathcal{O}_{S, \bar{s}}^{sh})$ is an initial object of the category of pro-étale neighbourhoods of \bar{s} . □

Contrary to the situation with the étale topos of S it is not true that every point of $Sh(S_{pro\text{-}\acute{e}tale})$ is of this form, and it is not true that the collection of points associated to geometric points is conservative. Namely, suppose that $S = \text{Spec}(k)$

where k is an algebraically closed field. Let A be an abelian group. Consider the sheaf \mathcal{F} on $S_{pro\text{-}\acute{e}tale}$ defined by the rule

$$\mathcal{F}(U) = \frac{\{\text{functions } U \rightarrow A\}}{\{\text{locally constant functions}\}}$$

Then $\mathcal{F}(U) = 0$ if $U = S = \text{Spec}(k)$ but in general \mathcal{F} is not zero. Namely, $S_{pro\text{-}\acute{e}tale}$ contains quasi-compact objects with infinitely many points. For example, let $E = \lim E_n$ be an inverse limit of finite sets with surjective transition maps, e.g., $E = \lim \mathbf{Z}/n\mathbf{Z}$. The scheme $\text{Spec}(\text{colim Map}(E_n, k))$ is an object of $S_{pro\text{-}\acute{e}tale}$ because $\text{colim Map}(E_n, k)$ is weakly étale (even ind-Zariski) over k . Thus \mathcal{F} is a nonzero abelian sheaf whose stalk at the unique geometric point of S is zero.

The solution is to use the existence of quasi-compact, weakly contractible objects. First, there are enough quasi-compact, weakly contractible objects by Lemma 11.27. Second, if $W \in \text{Ob}(S_{pro\text{-}\acute{e}tale})$ is quasi-compact, weakly contractible, then the functor

$$\text{Sh}(S_{pro\text{-}\acute{e}tale}) \longrightarrow \text{Sets}, \quad \mathcal{F} \longmapsto \mathcal{F}(W)$$

is an exact functor $\text{Sh}(S_{pro\text{-}\acute{e}tale}) \rightarrow \text{Sets}$ which commutes with all limits. The functor

$$\text{Ab}(S_{pro\text{-}\acute{e}tale}) \longrightarrow \text{Ab}, \quad \mathcal{F} \longmapsto \mathcal{F}(W)$$

is exact and commutes with direct sums (as W is quasi-compact, see Sites, Lemma 17.5), hence commutes with all limits and colimits. Moreover, we can check exactness of a complex of abelian sheaves by evaluation at the quasi-compact, weakly contractible objects of $S_{pro\text{-}\acute{e}tale}$, see Cohomology on Sites, Proposition 41.2.

13. Compact generation

0994 Let S be a scheme. The site $S_{pro\text{-}\acute{e}tale}$ has enough quasi-compact, weakly contractible objects U . For any sheaf of rings \mathcal{A} on $S_{pro\text{-}\acute{e}tale}$ the corresponding objects $j_{U!}\mathcal{A}_U$ are compact objects of the derived category $D(\mathcal{A})$, see Cohomology on Sites, Lemma 42.5. Since every complex of \mathcal{A} -modules is quasi-isomorphic to a complex whose terms are direct sums of the modules $j_{U!}\mathcal{A}_U$ (details omitted). Thus we see that $D(\mathcal{A})$ is generated by its compact objects.

The same argument works for the big pro-étale site of S .

14. Derived completion in the constant Noetherian case

099L We continue the discussion started in Local Cohomology, Section 12; we assume the reader has read at least some of that section. Let \mathcal{C} be a site. Let Λ be a Noetherian ring and let $I \subset \Lambda$ be an ideal. Recall from Modules on Sites, Lemma 41.4 that

$$\underline{\Lambda}^\wedge = \lim \underline{\Lambda/I^n}$$

is a flat $\underline{\Lambda}$ -algebra and that the map $\underline{\Lambda} \rightarrow \underline{\Lambda}^\wedge$ identifies quotients by I . Hence Local Cohomology, Lemma 12.17 tells us that

$$D_{comp}(\mathcal{C}, \Lambda) = D_{comp}(\mathcal{C}, \underline{\Lambda}^\wedge)$$

In particular the cohomology sheaves $H^i(K)$ of an object K of $D_{comp}(\mathcal{C}, \Lambda)$ are sheaves of $\underline{\Lambda}^\wedge$ -modules. For notational convenience we often work with $D_{comp}(\mathcal{C}, \Lambda)$.

099M **Lemma 14.1.** *Let \mathcal{C} be a site. Let Λ be a Noetherian ring and let $I \subset \Lambda$ be an ideal. The left adjoint to the inclusion functor $D_{\text{comp}}(\mathcal{C}, \Lambda) \rightarrow D(\mathcal{C}, \Lambda)$ of Local Cohomology, Proposition 12.12 sends K to*

$$K^\wedge = R\lim(K \otimes_\Lambda^{\mathbf{L}} \underline{\Lambda/I^n})$$

In particular, K is derived complete if and only if $K = R\lim(K \otimes_\Lambda^{\mathbf{L}} \underline{\Lambda/I^n})$.

Proof. Choose generators f_1, \dots, f_r of I . By Local Cohomology, Lemma 12.9 we have

$$K^\wedge = R\lim(K \otimes_\Lambda^{\mathbf{L}} K_n)$$

where $K_n = K(\Lambda, f_1^n, \dots, f_r^n)$. In More on Algebra, Lemma 82.1 we have seen that the pro-systems $\{K_n\}$ and $\{\Lambda/I^n\}$ of $D(\Lambda)$ are isomorphic. Thus the lemma follows. \square

099N **Lemma 14.2.** *Let Λ be a Noetherian ring. Let $I \subset \Lambda$ be an ideal. Let $f : \text{Sh}(\mathcal{D}) \rightarrow \text{Sh}(\mathcal{C})$ be a morphism of topoi. Then*

- (1) Rf_* sends $D_{\text{comp}}(\mathcal{D}, \Lambda)$ into $D_{\text{comp}}(\mathcal{C}, \Lambda)$,
- (2) the map $Rf_* : D_{\text{comp}}(\mathcal{D}, \Lambda) \rightarrow D_{\text{comp}}(\mathcal{C}, \Lambda)$ has a left adjoint $Lf_{\text{comp}}^* : D_{\text{comp}}(\mathcal{C}, \Lambda) \rightarrow D_{\text{comp}}(\mathcal{D}, \Lambda)$ which is Lf^* followed by derived completion,
- (3) Rf_* commutes with derived completion,
- (4) for K in $D_{\text{comp}}(\mathcal{D}, \Lambda)$ we have $Rf_*K = R\lim Rf_*(K \otimes_\Lambda^{\mathbf{L}} \underline{\Lambda/I^n})$.
- (5) for M in $D_{\text{comp}}(\mathcal{C}, \Lambda)$ we have $Lf_{\text{comp}}^*M = R\lim Lf^*(M \otimes_\Lambda^{\mathbf{L}} \underline{\Lambda/I^n})$.

Proof. We have seen (1) and (2) in Local Cohomology, Lemma 12.18. Part (3) follows from Local Cohomology, Lemma 12.19. For (4) let K be derived complete. Then

$$Rf_*K = Rf_*(R\lim K \otimes_\Lambda^{\mathbf{L}} \underline{\Lambda/I^n}) = R\lim Rf_*(K \otimes_\Lambda^{\mathbf{L}} \underline{\Lambda/I^n})$$

the first equality by Lemma 14.1 and the second because Rf_* commutes with $R\lim$ (Cohomology on Sites, Lemma 22.3). This proves (4). To prove (5), by Lemma 14.1 we have

$$Lf_{\text{comp}}^*M = R\lim(Lf^*M \otimes_\Lambda^{\mathbf{L}} \underline{\Lambda/I^n})$$

Since Lf^* commutes with derived tensor product by Cohomology on Sites, Lemma 19.4 and since $Lf^*\underline{\Lambda/I^n} = \underline{\Lambda/I^n}$ we get (5). \square

15. Derived completion on the pro-étale site

099P Let \mathcal{C} be a site. Let Λ be a Noetherian ring. Let $I \subset \Lambda$ be an ideal. Although the general theory (see Local Cohomology, Section 12 and Section 14) concerning $D_{\text{comp}}(\mathcal{C}, \Lambda)$ is quite satisfactory it is somewhat useless as it is hard to explicitly give examples of derived complete complexes. We know that

- (1) every object M of $D(\mathcal{C}, \Lambda/I^n)$ restricts to a derived complete object of $D(\mathcal{C}, \Lambda)$, and
- (2) for every $K \in D(\mathcal{C}, \Lambda)$ the derived completion $K^\wedge = R\lim(K \otimes_\Lambda^{\mathbf{L}} \underline{\Lambda/I^n})$ is derived complete.

The first type of objects are trivially complete and perhaps not interesting. The problem with (2) is that derived completion in general is somewhat mysterious, even in case $K = \underline{\Lambda}$. Namely, by definition of homotopy limits there is a distinguished triangle

$$R\lim(\underline{\Lambda/I^n}) \rightarrow \prod \underline{\Lambda/I^n} \rightarrow \prod \underline{\Lambda/I^n} \rightarrow R\lim(\underline{\Lambda/I^n})[1]$$

in $D(\mathcal{C}, \Lambda)$ where the products are in $D(\mathcal{C}, \Lambda)$. These are computed by taking products of injective resolutions (Injectives, Lemma 13.4), so we see that the sheaf $H^p(\varprojlim \Lambda/I^n)$ is the sheafification of the presheaf

$$U \mapsto \prod H^p(U, \Lambda/I^n).$$

As an explicit example, if $X = \text{Spec}(\mathbf{C}[t, t^{-1}])$, $\mathcal{C} = X_{\text{étale}}$, $\Lambda = \mathbf{Z}$, $I = (2)$, and $p = 1$, then we get the sheafification of the presheaf

$$U \mapsto \prod H^1(U_{\text{étale}}, \mathbf{Z}/2^n \mathbf{Z})$$

for U étale over X . Note that $H^1(X_{\text{étale}}, \mathbf{Z}/m\mathbf{Z})$ is cyclic of order m with generator α_m given by the finite étale $\mathbf{Z}/m\mathbf{Z}$ -covering given by the equation $t = s^m$ (see Étale Cohomology, Section 6). Then the section

$$\alpha = (\alpha_{2^n}) \in \prod H^1(X_{\text{étale}}, \mathbf{Z}/2^n \mathbf{Z})$$

of the presheaf above does not restrict to zero on any nonempty étale scheme over X , whence the sheaf associated to the presheaf is not zero.

However, on the pro-étale site this phenomenon does not occur. The reason is that we have enough (quasi-compact) weakly contractible objects. In the following proposition we collect some results about derived completion in the Noetherian constant case for sites having enough weakly contractible objects (see Sites, Definition 39.2).

099Q **Proposition 15.1.** *Let \mathcal{C} be a site. Assume \mathcal{C} has enough weakly contractible objects. Let Λ be a Noetherian ring. Let $I \subset \Lambda$ be an ideal.*

- (1) *The category of derived complete sheaves Λ -modules is a weak Serre subcategory of $\text{Mod}(\mathcal{C}, \Lambda)$.*
- (2) *A sheaf \mathcal{F} of Λ -modules satisfies $\mathcal{F} = \lim \mathcal{F}/I^n \mathcal{F}$ if and only if \mathcal{F} is derived complete and $\bigcap I^n \mathcal{F} = 0$.*
- (3) *The sheaf $\underline{\Lambda}^\wedge$ is derived complete.*
- (4) *If $\dots \rightarrow \mathcal{F}_3 \rightarrow \mathcal{F}_2 \rightarrow \mathcal{F}_1$ is an inverse system of derived complete sheaves of Λ -modules, then $\lim \mathcal{F}_n$ is derived complete.*
- (5) *An object $K \in D(\mathcal{C}, \Lambda)$ is derived complete if and only if each cohomology sheaf $H^p(K)$ is derived complete.*
- (6) *An object $K \in D_{\text{comp}}(\mathcal{C}, \Lambda)$ is bounded above if and only if $K \otimes_{\Lambda}^{\mathbf{L}} \underline{\Lambda}/I$ is bounded above.*
- (7) *An object $K \in D_{\text{comp}}(\mathcal{C}, \Lambda)$ is bounded if $K \otimes_{\Lambda}^{\mathbf{L}} \underline{\Lambda}/I$ has finite tor dimension.*

Proof. Let $\mathcal{B} \subset \text{Ob}(\mathcal{C})$ be a subset such that every $U \in \mathcal{B}$ is weakly contractible and every object of \mathcal{C} has a covering by elements of \mathcal{B} . We will use the results of Cohomology on Sites, Lemma 41.1 and Proposition 41.2 without further mention.

Recall that $R\lim$ commutes with $R\Gamma(U, -)$, see Injectives, Lemma 13.6. Let $f \in I$. Recall that $T(K, f)$ is the homotopy limit of the system

$$\dots K \xrightarrow{f} K \xrightarrow{f} K$$

in $D(\mathcal{C}, \Lambda)$. Thus

$$R\Gamma(U, T(K, f)) = T(R\Gamma(U, K), f).$$

Since we can test isomorphisms of maps between objects of $D(\mathcal{C}, \Lambda)$ by evaluating at $U \in \mathcal{B}$ we conclude an object K of $D(\mathcal{C}, \Lambda)$ is derived complete if and only if for every $U \in \mathcal{B}$ the object $R\Gamma(U, K)$ is derived complete as an object of $D(\Lambda)$.

The remark above implies that items (1), (5) follow from the corresponding results for modules over rings, see More on Algebra, Lemmas 80.1 and 80.6. In the same way (2) can be deduced from More on Algebra, Proposition 80.5 as $(I^n \mathcal{F})(U) = I^n \cdot \mathcal{F}(U)$ for $U \in \mathcal{B}$ (by exactness of evaluating at U).

Proof of (4). The homotopy limit $R\lim \mathcal{F}_n$ is in $D_{comp}(X, \Lambda)$ (see discussion following Local Cohomology, Definition 12.4). By part (5) just proved we conclude that $\lim \mathcal{F}_n = H^0(R\lim \mathcal{F}_n)$ is derived complete. Part (3) is a special case of (4).

Proof of (6) and (7). Follows from Lemma 14.1 and Cohomology on Sites, Lemma 38.8 and the computation of homotopy limits in Cohomology on Sites, Proposition 41.2. \square

16. Comparison with the étale site

099R Let X be a scheme. With suitable choices of sites (as in Topologies, Remark 10.1) the functor $u : X_{\acute{e}tale} \rightarrow X_{pro\text{-}\acute{e}tale}$ sending U/X to U/X defines a morphism of sites

$$\epsilon : X_{pro\text{-}\acute{e}tale} \longrightarrow X_{\acute{e}tale}$$

This follows from Sites, Proposition 14.6. A fundamental fact about this comparison morphism is the following.

099S **Lemma 16.1.** *Let X be a scheme. Let $Y = \lim Y_i$ be an inverse limit of quasi-compact and quasi-separated schemes étale over X with affine transition morphisms. For any sheaf \mathcal{F} on $X_{\acute{e}tale}$ we have $\epsilon^{-1}\mathcal{F}(Y) = \text{colim } \mathcal{F}(Y_i)$.*

Proof. Let $\mathcal{F} = h_U$ be a representable sheaf on $X_{\acute{e}tale}$ with U an object of $X_{\acute{e}tale}$. In this case $\epsilon^{-1}h_U = h_{u(U)}$ where $u(U)$ is U viewed as an object of $X_{pro\text{-}\acute{e}tale}$ (Sites, Lemma 13.5). Then

$$\begin{aligned} (\epsilon^{-1}h_U)(Y) &= h_{u(U)}(Y) \\ &= \text{Mor}_X(Y, U) \\ &= \text{colim } \text{Mor}_X(Y_i, U) \\ &= \text{colim } h_{u(U)}(Y_i) \\ &= \text{colim}(\epsilon^{-1}h_U)(Y_i) \end{aligned}$$

Here the only nonformal equality is the 3rd which holds by Limits, Proposition 6.1. Hence the lemma holds for every representable sheaf. Since every sheaf is a coequalizer of a map of coproducts of representable sheaves (Sites, Lemma 12.5) we obtain the result in general. \square

099T **Lemma 16.2.** *Let X be a scheme. For every sheaf \mathcal{F} on $X_{\acute{e}tale}$ the adjunction map $\mathcal{F} \rightarrow \epsilon_*\epsilon^{-1}\mathcal{F}$ is an isomorphism.*

Proof. Suppose that U is a quasi-compact and quasi-separated scheme étale over X . Then

$$\epsilon_*\epsilon^{-1}\mathcal{F}(U) = \epsilon^{-1}\mathcal{F}(U) = \mathcal{F}(U)$$

the second equality by (a special case of) Lemma 16.1. Since every object of $X_{\acute{e}tale}$ has a covering by quasi-compact and quasi-separated objects we conclude. \square

099U **Lemma 16.3.** *Let X be an affine scheme. For injective abelian sheaf \mathcal{I} on $X_{\acute{e}tale}$ we have $H^p(X_{pro\text{-}\acute{e}tale}, \epsilon^{-1}\mathcal{I}) = 0$ for $p > 0$.*

Proof. We are going to use Cohomology on Sites, Lemma 11.9 to prove this. The idea is simple: We show that every standard pro-étale covering of X is a limit of coverings in $X_{\acute{e}tale}$. If this holds then Lemma 16.1 will kick in to show the Čech cohomology groups of $\epsilon^{-1}\mathcal{I}$ are colimits of those of \mathcal{I} which are zero in positive degree.

Here are the details. Let $\mathcal{B} \subset \text{Ob}(X_{pro\text{-}\acute{e}tale})$ be the set of affine schemes U over X such that $\mathcal{O}(X) \rightarrow \mathcal{O}(U)$ is ind-étale. Let Cov be the set of pro-étale coverings $\{U_i \rightarrow U\}_{i=1, \dots, n}$ with $U, U_i \in \mathcal{B}$ such that $\mathcal{O}(U) \rightarrow \mathcal{O}(U_i)$ is ind-étale for $i = 1, \dots, n$. Properties (1) and (2) of Cohomology on Sites, Lemma 11.9 hold for \mathcal{B} and Cov by Proposition 9.1 (it also follows from Lemma 11.10).

To check condition (3) suppose that $\{U_i \rightarrow U\}_{i=1, \dots, n}$ is an element of Cov . Then we can write $U_i = \lim_{a \in A_i} U_{i,a}$ with $U_{i,a} \rightarrow U$ étale and $U_{i,a}$ affine. Next we write $U = \lim_{b \in B} U_b$ with U_b affine and $U_b \rightarrow U$ étale. By Limits, Lemma 10.1 for each i and $a \in A_i$ we can choose a $b(i, a) \in B$ and for all $b \geq b(i, a)$ an affine scheme $U_{i,a,b}$ étale over U_b such that $U_{i,a} = \lim_{b \geq b(i,a)} U_{i,a,b}$ ². Moreover, any transition map $U_{i,a} \rightarrow U_{i,a'}$ comes from an essentially unique morphism $U_{i,a,b} \rightarrow U_{i,a',b}$ for b large enough (by the same reference). Finally, given $a_1 \in A_1, \dots, a_n \in A_n$ the morphism $U_{1,a_1} \amalg \dots \amalg U_{n,a_n} \rightarrow U$ is surjective, hence for b large enough the map $U_{1,a_1,b} \amalg \dots \amalg U_{n,a_n,b} \rightarrow U_b$ is surjective by Limits, Lemma 8.13. Let \mathcal{D} be the category of coverings $\{U_{i,a_i,b} \rightarrow U_b\}_{i=1, \dots, n}$ so obtained. This category is cofiltered. We claim that, given $i_0, \dots, i_p \in \{1, \dots, n\}$ we have

$$U_{i_0} \times_U U_{i_1} \times_U \dots \times_U U_{i_p} = \lim_{\mathcal{D}} U_{i_0, a_{i_0}, b} \times_{U_b} U_{i_1, a_{i_1}, b} \times_{U_b} \dots \times_{U_b} U_{i_p, a_{i_p}, b}$$

This is clear from the fact that it holds for $p = -1$ (i.e., $U = \lim_{\mathcal{D}} U_b$) and for $p = 0$ (i.e., $U_i = \lim_{\mathcal{D}} U_{i,a_i,b}$) and the fact that fibre products commute with limits. Then finally it follows from Lemma 16.1 that

$$\check{C}^\bullet(\{U_i \rightarrow U\}, \epsilon^{-1}\mathcal{I}) = \text{colim}_{\mathcal{D} \circ \text{opp}} \check{C}^\bullet(\{U_{i,a_i,b} \rightarrow U_b\}, \mathcal{I})$$

Since each of the Čech complexes on the right hand side is acyclic in positive degrees (Cohomology on Sites, Lemma 11.2) it follows that the one on the left is too. This prove condition (3) of Cohomology on Sites, Lemma 11.9. Since $X \in \mathcal{B}$ the lemma follows. \square

099V **Lemma 16.4.** *Let X be a scheme. For an abelian sheaf \mathcal{F} on $X_{\acute{e}tale}$ we have $R\epsilon_* (\epsilon^{-1}\mathcal{F}) = \mathcal{F}$.*

Proof. Let \mathcal{I} be an injective abelian sheaf on $X_{\acute{e}tale}$. Recall that $R^q \epsilon_* (\epsilon^{-1}\mathcal{I})$ is the sheaf associated to $U \mapsto H^q(U_{pro\text{-}\acute{e}tale}, \epsilon^{-1}\mathcal{I})$, see Cohomology on Sites, Lemma 8.4. By Lemma 16.3 we see that this is zero for $q > 0$ and U affine and étale over X . Since every object of $X_{\acute{e}tale}$ has a covering by affine objects, it follows that $R^q \epsilon_* (\epsilon^{-1}\mathcal{I}) = 0$ for $q > 0$. Combined with Lemma 16.2 we conclude that $R\epsilon_* \epsilon^{-1}\mathcal{I} = \mathcal{I}$ for every injective abelian sheaf. Since every abelian sheaf has a resolution by injective sheaves, the result follows. (Hint: use Leray acyclicity theorem – Derived Categories, Lemma 17.7.) \square

²To be sure, we pick $U_{i,a,b} = U_b \times_{U_{b(i,a)}} U_{i,a,b(i,a)}$ although this isn't necessary for what follows.

099W **Lemma 16.5.** *Let X be a scheme. For an abelian sheaf \mathcal{F} on $X_{\acute{e}tale}$ we have*

$$H^i(X_{\acute{e}tale}, \mathcal{F}) = H^i(X_{pro\text{-}\acute{e}tale}, \epsilon^{-1}\mathcal{F})$$

for all i .

Proof. Immediate consequence of Lemma 16.4 and the Leray spectral sequence (Cohomology on Sites, Lemma 15.6). \square

099X **Lemma 16.6.** *Let X be a scheme. Let \mathcal{G} be a sheaf of (possibly noncommutative) groups on $X_{\acute{e}tale}$. We have*

$$H^1(X_{\acute{e}tale}, \mathcal{G}) = H^1(X_{pro\text{-}\acute{e}tale}, \epsilon^{-1}\mathcal{G})$$

where H^1 is defined as the set of isomorphism classes of torsors (see Cohomology on Sites, Section 5).

Proof. Since the functor ϵ^{-1} is fully faithful by Lemma 16.2 it is clear that the map $H^1(X_{\acute{e}tale}, \mathcal{G}) \rightarrow H^1(X_{pro\text{-}\acute{e}tale}, \epsilon^{-1}\mathcal{G})$ is injective. To show surjectivity it suffices to show that any $\epsilon^{-1}\mathcal{G}$ -torsor \mathcal{F} is étale locally trivial. To do this we may assume that X is affine. Thus we reduce to proving surjectivity for X affine.

Choose a covering $\{U \rightarrow X\}$ with (a) U affine, (b) $\mathcal{O}(X) \rightarrow \mathcal{O}(U)$ ind-étale, and (c) $\mathcal{F}(U)$ nonempty. We can do this by Proposition 9.1 and the fact that standard pro-étale coverings of X are cofinal among all pro-étale coverings of X (Lemma 11.5). Write $U = \lim U_i$ as a limit of affine schemes étale over X . Pick $s \in \mathcal{F}(U)$. Let $g \in \epsilon^{-1}\mathcal{G}(U \times_X U)$ be the unique section such that $g \cdot \text{pr}_1^*s = \text{pr}_2^*s$ in $\mathcal{F}(U \times_X U)$. Then g satisfies the cocycle condition

$$\text{pr}_{12}^*g \cdot \text{pr}_{23}^*g = \text{pr}_{13}^*g$$

in $\epsilon^{-1}\mathcal{G}(U \times_X U \times_X U)$. By Lemma 16.1 we have

$$\epsilon^{-1}\mathcal{G}(U \times_X U) = \text{colim } \mathcal{G}(U_i \times_X U_i)$$

and

$$\epsilon^{-1}\mathcal{G}(U \times_X U \times_X U) = \text{colim } \mathcal{G}(U_i \times_X U_i \times_X U_i)$$

hence we can find an i and an element $g_i \in \mathcal{G}(U_i)$ mapping to g satisfying the cocycle condition. The cocycle g_i then defines a torsor for \mathcal{G} on $X_{\acute{e}tale}$ whose pullback is isomorphic to \mathcal{F} by construction. Some details omitted (namely, the relationship between torsors and 1-cocycles which should be added to the chapter on cohomology on sites). \square

09B1 **Lemma 16.7.** *Let X be a scheme. Let Λ be a ring.*

- (1) *The essential image of $\epsilon^{-1} : \text{Mod}(X_{\acute{e}tale}, \Lambda) \rightarrow \text{Mod}(X_{pro\text{-}\acute{e}tale}, \Lambda)$ is a weak Serre subcategory \mathcal{C} .*
- (2) *The functor ϵ^{-1} defines an equivalence of categories of $D^+(X_{\acute{e}tale}, \Lambda)$ with $D_{\mathcal{C}}^+(X_{pro\text{-}\acute{e}tale}, \Lambda)$.*

Proof. To prove (1) we will prove conditions (1) – (4) of Homology, Lemma 9.3. Since ϵ^{-1} is fully faithful (Lemma 16.2) and exact, everything is clear except for condition (4). However, if

$$0 \rightarrow \epsilon^{-1}\mathcal{F}_1 \rightarrow \mathcal{G} \rightarrow \epsilon^{-1}\mathcal{F}_2 \rightarrow 0$$

is a short exact sequence of sheaves of Λ -modules on $X_{pro\text{-}\acute{e}tale}$, then we get

$$0 \rightarrow \epsilon_*\epsilon^{-1}\mathcal{F}_1 \rightarrow \epsilon_*\mathcal{G} \rightarrow \epsilon_*\epsilon^{-1}\mathcal{F}_2 \rightarrow R^1\epsilon_*\epsilon^{-1}\mathcal{F}_1$$

which by Lemma 16.4 is the same as a short exact sequence

$$0 \rightarrow \mathcal{F}_1 \rightarrow \epsilon_* \mathcal{G} \rightarrow \mathcal{F}_2 \rightarrow 0$$

Pulling back we find that $\mathcal{G} = \epsilon^{-1} \epsilon_* \mathcal{G}$. This proves (1).

By (1) and the discussion in Derived Categories, Section 13 we obtain a strictly full, saturated, triangulated subcategory $D_{\mathcal{C}}(X_{\text{pro-étale}}, \Lambda)$. It is clear that ϵ^{-1} maps $D(X_{\text{étale}}, \Lambda)$ into $D_{\mathcal{C}}(X_{\text{pro-étale}}, \Lambda)$. If M is in $D^+(X_{\text{étale}}, \Lambda)$, then Lemma 16.4 shows that $M \rightarrow R\epsilon_* \epsilon^{-1} M$ is an isomorphism. If K is in $D_{\mathcal{C}}^+(X_{\text{pro-étale}}, \Lambda)$, then the spectral sequence

$$R^q \epsilon_* H^p(K) \Rightarrow H^{p+q}(R\epsilon_* K)$$

and the vanishing in Lemma 16.4 shows that $H^p(R\epsilon_* K) = R\epsilon_* H^p(K)$. Since ϵ is a flat morphism of ringed sites (ringed by the constant sheaf $\underline{\Lambda}$) we see that $\epsilon^{-1} R\epsilon_* K$ has cohomology sheaves $\epsilon^{-1} R\epsilon_* H^p(K)$. Since we've assumed $H^p(K)$ is in \mathcal{C} we conclude by Lemma 16.4 once more that $\epsilon^{-1} R\epsilon_* K \rightarrow K$ is an isomorphism. In this way we see that ϵ^{-1} and $R\epsilon_*$ are quasi-inverse functors proving (2). \square

Let Λ be a ring. In Modules on Sites, Section 42 we have defined the notion of a locally constant sheaf of Λ -modules on a site. If M is a Λ -module, then \underline{M} is of finite presentation as a sheaf of $\underline{\Lambda}$ -modules if and only if M is a finitely presented Λ -module, see Modules on Sites, Lemma 41.5.

099Y **Lemma 16.8.** *Let X be a scheme. Let Λ be a ring. The functor ϵ^{-1} defines an equivalence of categories*

$$\left\{ \begin{array}{l} \text{locally constant sheaves} \\ \text{of } \Lambda\text{-modules on } X_{\text{étale}} \\ \text{of finite presentation} \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{l} \text{locally constant sheaves} \\ \text{of } \Lambda\text{-modules on } X_{\text{pro-étale}} \\ \text{of finite presentation} \end{array} \right\}$$

Proof. Let \mathcal{F} be a locally constant sheaf of Λ -modules on $X_{\text{pro-étale}}$ of finite presentation. Choose a pro-étale covering $\{U_i \rightarrow X\}$ such that $\mathcal{F}|_{U_i}$ is constant, say $\mathcal{F}|_{U_i} \cong \underline{M}_i|_{U_i}$. Observe that $U_i \times_X U_j$ is empty if M_i is not isomorphic to M_j . For each Λ -module M let $I_M = \{i \in I \mid M_i \cong M\}$. As pro-étale coverings are fpqc coverings and by Descent, Lemma 10.2 we see that $U_M = \bigcup_{i \in I_M} \text{Im}(U_i \rightarrow X)$ is an open subset of X . Then $X = \coprod U_M$ is a disjoint open covering of X . We may replace X by U_M for some M and assume that $M_i = M$ for all i .

Consider the sheaf $\mathcal{I} = \text{Isom}(\underline{M}, \mathcal{F})$. This sheaf is a torsor for $\mathcal{G} = \text{Isom}(\underline{M}, \underline{M})$. By Modules on Sites, Lemma 42.4 we have $\mathcal{G} = \underline{G}$ where $G = \text{Isom}_{\Lambda}(M, M)$. Since torsors for the étale topology and the pro-étale topology agree by Lemma 16.6 it follows that \mathcal{I} has sections étale locally on X . Thus \mathcal{F} is étale locally a constant sheaf which is what we had to show. \square

099Z **Lemma 16.9.** *Let X be a scheme. Let Λ be a Noetherian ring. Let $D_{\text{flc}}(X_{\text{étale}}, \Lambda)$, resp. $D_{\text{flc}}(X_{\text{pro-étale}}, \Lambda)$ be the full subcategory of $D(X_{\text{étale}}, \Lambda)$, resp. $D(X_{\text{pro-étale}}, \Lambda)$ consisting of those complexes whose cohomology sheaves are locally constant sheaves of Λ -modules of finite type. Then*

$$\epsilon^{-1} : D_{\text{flc}}^+(X_{\text{étale}}, \Lambda) \longrightarrow D_{\text{flc}}^+(X_{\text{pro-étale}}, \Lambda)$$

is an equivalence of categories.

Proof. The categories $D_{flc}(X_{\acute{e}tale}, \Lambda)$ and $D_{flc}(X_{pro\text{-}\acute{e}tale}, \Lambda)$ are strictly full, saturated, triangulated subcategories of $D(X_{\acute{e}tale}, \Lambda)$ and $D(X_{pro\text{-}\acute{e}tale}, \Lambda)$ by Modules on Sites, Lemma 42.5 and Derived Categories, Section 13. The statement of the lemma follows by combining Lemmas 16.7 and 16.8. \square

09B2 **Lemma 16.10.** *Let X be a scheme. Let Λ be a Noetherian ring. Let K be an object of $D(X_{pro\text{-}\acute{e}tale}, \Lambda)$. Set $K_n = K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n$. If K_1 is*

- (1) *in the essential image of $\epsilon^{-1} : D(X_{\acute{e}tale}, \Lambda/I) \rightarrow D(X_{pro\text{-}\acute{e}tale}, \Lambda/I)$, and*
- (2) *has tor amplitude in $[a, \infty)$ for some $a \in \mathbf{Z}$,*

then (1) and (2) hold for K_n as an object of $D(X_{pro\text{-}\acute{e}tale}, \Lambda/I^n)$.

Proof. For assertion (2) this follows from the more general Cohomology on Sites, Lemma 38.8. The second assertion follows from the fact that the essential image of ϵ^{-1} is a triangulated subcategory of $D^+(X_{pro\text{-}\acute{e}tale}, \Lambda/I^n)$ (Lemma 16.7), the distinguished triangles

$$K \otimes_{\Lambda}^{\mathbf{L}} I^n/I^{n+1} \rightarrow K_{n+1} \rightarrow K_n \rightarrow K \otimes_{\Lambda}^{\mathbf{L}} I^n/I^{n+1}[1]$$

and the isomorphism

$$K \otimes_{\Lambda}^{\mathbf{L}} I^n/I^{n+1} = K_1 \otimes_{\Lambda/I}^{\mathbf{L}} I^n/I^{n+1}$$

\square

17. Cohomology of a point

09B3 Let Λ be a Noetherian ring complete with respect to an ideal $I \subset \Lambda$. Let k be a field. In this section we “compute”

$$H^i(\mathrm{Spec}(k)_{pro\text{-}\acute{e}tale}, \underline{\Lambda}^{\wedge})$$

where $\underline{\Lambda}^{\wedge} = \lim_m \underline{\Lambda}/I^m$ as before. Let k^{sep} be a separable algebraic closure of k . Then

$$\mathcal{U} = \{\mathrm{Spec}(k^{sep}) \rightarrow \mathrm{Spec}(k)\}$$

is a pro-étale covering of $\mathrm{Spec}(k)$. We will use the Čech to cohomology spectral sequence with respect to this covering. Set $U_0 = \mathrm{Spec}(k^{sep})$ and

$$\begin{aligned} U_n &= \mathrm{Spec}(k^{sep}) \times_{\mathrm{Spec}(k)} \mathrm{Spec}(k^{sep}) \times_{\mathrm{Spec}(k)} \dots \times_{\mathrm{Spec}(k)} \mathrm{Spec}(k^{sep}) \\ &= \mathrm{Spec}(k^{sep} \otimes_k k^{sep} \otimes_k \dots \otimes_k k^{sep}) \end{aligned}$$

($n+1$ factors). Note that the underlying topological space $|U_0|$ of U_0 is a singleton and for $n \geq 1$ we have

$$|U_n| = G \times \dots \times G \quad (n \text{ factors})$$

as profinite spaces where $G = \mathrm{Gal}(k^{sep}/k)$. Namely, every point of U_n has residue field k^{sep} and we identify $(\sigma_1, \dots, \sigma_n)$ with the point corresponding to the surjection

$$k^{sep} \otimes_k k^{sep} \otimes_k \dots \otimes_k k^{sep} \longrightarrow k^{sep}, \quad \lambda_0 \otimes \lambda_1 \otimes \dots \otimes \lambda_n \longmapsto \lambda_0 \sigma_1(\lambda_1) \dots \sigma_n(\lambda_n)$$

Then we compute

$$\begin{aligned} R\Gamma((U_n)_{pro\text{-}\acute{e}tale}, \underline{\Lambda}^{\wedge}) &= R\lim_m R\Gamma((U_n)_{pro\text{-}\acute{e}tale}, \underline{\Lambda}/I^m) \\ &= R\lim_m R\Gamma((U_n)_{\acute{e}tale}, \underline{\Lambda}/I^m) \\ &= \lim_m H^0(U_n, \underline{\Lambda}/I^m) \\ &= \mathrm{Maps}_{cont}(G \times \dots \times G, \Lambda) \end{aligned}$$

The first equality because $R\Gamma$ commutes with derived limits and as Λ^\wedge is the derived limit of the sheaves $\underline{\Lambda/I^m}$ by Proposition 15.1. The second equality by Lemma 16.5. The third equality by Étale Cohomology, Lemma 54.7. The fourth equality uses Étale Cohomology, Remark 23.2 to identify sections of the constant sheaf $\underline{\Lambda/I^m}$. Then it uses the fact that Λ is complete with respect to I and hence equal to $\lim_m \Lambda/I^m$ as a topological space, to see that $\lim_m \text{Map}_{cont}(G, \Lambda/I^m) = \text{Map}_{cont}(G, \Lambda)$ and similarly for higher powers of G . At this point Cohomology on Sites, Lemmas 11.3 and 11.7 tell us that

$$\Lambda \rightarrow \text{Maps}_{cont}(G, \Lambda) \rightarrow \text{Maps}_{cont}(G \times G, \Lambda) \rightarrow \dots$$

computes the pro-étale cohomology. In other words, we see that

$$H^i(\text{Spec}(k)_{pro\text{-}\acute{e}tale}, \underline{\Lambda}^\wedge) = H_{cont}^i(G, \Lambda)$$

where the right hand side is continuous group cohomology as defined by Tate in [Tat76], see Étale Cohomology, Section 57. Of course, this is as it should be.

09B4 **Lemma 17.1.** *Let k be a field. Let $G = \text{Gal}(k^{sep}/k)$ be its absolute Galois group. Further,*

- (1) *let M be a profinite abelian group with a continuous G -action, or*
- (2) *let Λ be a Noetherian ring and $I \subset \Lambda$ an ideal and let M be an I -adically complete Λ -module with continuous G -action.*

Then there is a canonical sheaf \underline{M}^\wedge on $\text{Spec}(k)_{pro\text{-}\acute{e}tale}$ associated to M such that

$$H^i(\text{Spec}(k), \underline{M}^\wedge) = H_{cont}^i(G, M)$$

as abelian groups or Λ -modules.

Proof. Proof in case (2). Set $M_n = M/I^n M$. Then $M = \lim M_n$ as M is assumed I -adically complete. Since the action of G is continuous we get continuous actions of G on M_n . By Étale Cohomology, Theorem 55.3 this action corresponds to a (locally constant) sheaf \underline{M}_n of Λ/I^n -modules on $\text{Spec}(k)_{\acute{e}tale}$. Pull back to $\text{Spec}(k)_{pro\text{-}\acute{e}tale}$ by the comparison morphism ϵ and take the limit

$$\underline{M}^\wedge = \lim \epsilon^{-1} \underline{M}_n$$

to get the sheaf promised in the lemma. Exactly the same argument as given in the introduction of this section gives the comparison with Tate's continuous Galois cohomology. \square

18. Weakly contractible hypercoverings

09A0 Let X be a scheme. For every object $U \in \text{Ob}(X_{pro\text{-}\acute{e}tale})$ there exists a covering $\{V \rightarrow U\}$ of $X_{pro\text{-}\acute{e}tale}$ with V weakly contractible. This follows from Lemma 11.10 and the elementary fact that a disjoint union of weakly contractible objects in $X_{pro\text{-}\acute{e}tale}$ is weakly contractible (discussion of set theoretic issues omitted). This observation leads to the existence of hypercoverings made up out weakly contractible objects.

09A1 **Lemma 18.1.** *Let X be a scheme.*

- (1) *For every object U of $X_{pro\text{-}\acute{e}tale}$ there exists a hypercovering K of U in $X_{pro\text{-}\acute{e}tale}$ such that each term K_n consists of a single weakly contractible object of $X_{pro\text{-}\acute{e}tale}$ covering U .*

- (2) For every quasi-compact and quasi-separated object U of $X_{\text{pro-étale}}$ there exists a hypercovering K of U in $X_{\text{pro-étale}}$ such that each term K_n consists of a single affine and weakly contractible object of $X_{\text{pro-étale}}$ covering U .

Proof. Let $\mathcal{B} \subset \text{Ob}(X_{\text{pro-étale}})$ be the set of weakly contractible objects of $X_{\text{pro-étale}}$. Every object of $X_{\text{pro-étale}}$ has a covering by an element of \mathcal{B} by Lemma 11.10 and the elementary fact that a disjoint union of weakly contractible objects is weakly contractible. We apply Hypercoverings, Lemma 12.6 to get a hypercovering K of U such that $K_n = \{U_{n,i}\}_{i \in I_n}$ with I_n finite and $U_{n,i}$ weakly contractible. Then we can replace K by the hypercovering of U given by $\{U_n\}$ in degree n where $U_n = \coprod_{i \in I_n} U_{n,i}$. This is allowed by Hypercoverings, Remark 12.9.

Let $X_{\text{qcqs,pro-étale}} \subset X_{\text{pro-étale}}$ be the full subcategory consisting of quasi-compact and quasi-separated objects. A covering of $X_{\text{qcqs,pro-étale}}$ will be a finite pro-étale covering. Then $X_{\text{qcqs,pro-étale}}$ is a site, has fibre products, and the inclusion functor $X_{\text{qcqs,pro-étale}} \rightarrow X_{\text{pro-étale}}$ is continuous and commutes with fibre products. In particular, if K is a hypercovering of an object U in $X_{\text{qcqs,pro-étale}}$ then K is a hypercovering of U in $X_{\text{pro-étale}}$ by Hypercoverings, Lemma 12.5. Let $\mathcal{B} \subset \text{Ob}(X_{\text{qcqs,pro-étale}})$ be the set of affine and weakly contractible objects. By Lemma 11.10 and the fact that finite unions of affines are affine, for every object U of $X_{\text{qcqs,pro-étale}}$ there exists a covering $\{V \rightarrow U\}$ of $X_{\text{qcqs,pro-étale}}$ with $V \in \mathcal{B}$. By Hypercoverings, Lemma 12.6 we get a hypercovering K of U such that $K_n = \{U_{n,i}\}_{i \in I_n}$ with I_n finite and $U_{n,i}$ affine and weakly contractible. Then we can replace K by the hypercovering of U given by $\{U_n\}$ in degree n where $U_n = \coprod_{i \in I_n} U_{n,i}$. This is allowed by Hypercoverings, Remark 12.9. \square

In the following lemma we use the Čech complex $\mathcal{F}(K)$ associated to a hypercovering K in a site. See Hypercoverings, Section 5. If K is a hypercovering of U and $K_n = \{U_n \rightarrow U\}$, then the Čech complex looks like this:

$$\mathcal{F}(K) = (\mathcal{F}(U_0) \rightarrow \mathcal{F}(U_1) \rightarrow \mathcal{F}(U_2) \rightarrow \dots)$$

- 09A2 **Lemma 18.2.** *Let X be a scheme. Let $E \in D^+(X_{\text{pro-étale}})$ be represented by a bounded below complex \mathcal{E}^\bullet of abelian sheaves. Let K be a hypercovering of $U \in \text{Ob}(X_{\text{pro-étale}})$ with $K_n = \{U_n \rightarrow U\}$ where U_n is a weakly contractible object of $X_{\text{pro-étale}}$. Then*

$$R\Gamma(U, E) = \text{Tot}(\mathcal{E}^\bullet(K))$$

in $D(\text{Ab})$.

Proof. If $E = \mathcal{E}[n]$ is the object associated to a single abelian sheaf on $X_{\text{pro-étale}}$, then the spectral sequence of Hypercoverings, Lemma 5.3 implies that

$$R\Gamma(X_{\text{pro-étale}}, \mathcal{E}) = \mathcal{E}(K)$$

because the higher cohomology groups of any sheaf over U_n vanish, see Cohomology on Sites, Lemma 41.1.

If \mathcal{E}^\bullet is bounded below, then we can choose an injective resolution $\mathcal{E}^\bullet \rightarrow \mathcal{I}^\bullet$ and consider the map of complexes

$$\text{Tot}(\mathcal{E}^\bullet(K)) \longrightarrow \text{Tot}(\mathcal{I}^\bullet(K))$$

For every n the map $\mathcal{E}^\bullet(U_n) \rightarrow \mathcal{I}^\bullet(U_n)$ is a quasi-isomorphism because taking sections over U_n is exact. Hence the displayed map is a quasi-isomorphism by one of the spectral sequences of Homology, Lemma 22.6. Using the result of the first

paragraph we see that for every p the complex $\mathcal{I}^p(K)$ is acyclic in degrees $n > 0$ and computes $\mathcal{I}^p(U)$ in degree 0. Thus the other spectral sequence of Homology, Lemma 22.6 shows $\text{Tot}(\mathcal{I}^\bullet(K))$ computes $R\Gamma(U, E) = \mathcal{I}^\bullet(U)$. \square

09A3 **Lemma 18.3.** *Let X be a quasi-compact and quasi-separated scheme. The functor $R\Gamma(X, -) : D^+(X_{\text{pro-étale}}) \rightarrow D(\text{Ab})$ commutes with direct sums and homotopy colimits.*

Proof. The statement means the following: Suppose we have a family of objects E_i of $D^+(X_{\text{pro-étale}})$ such that $\bigoplus E_i$ is an object of $D^+(X_{\text{pro-étale}})$. Then $R\Gamma(X, \bigoplus E_i) = \bigoplus R\Gamma(X, E_i)$. To see this choose a hypercovering K of X with $K_n = \{U_n \rightarrow X\}$ where U_n is an affine and weakly contractible scheme, see Lemma 18.1. Let N be an integer such that $H^p(E_i) = 0$ for $p < N$. Choose a complex of abelian sheaves \mathcal{E}_i^\bullet representing E_i with $\mathcal{E}_i^p = 0$ for $p < N$. The termwise direct sum $\bigoplus \mathcal{E}_i^\bullet$ represents $\bigoplus E_i$ in $D(X_{\text{pro-étale}})$, see Injectives, Lemma 13.4. By Lemma 18.2 we have

$$R\Gamma(X, \bigoplus E_i) = \text{Tot}((\bigoplus \mathcal{E}_i^\bullet)(K))$$

and

$$R\Gamma(X, E_i) = \text{Tot}(\mathcal{E}_i^\bullet(K))$$

Since each U_n is quasi-compact we see that

$$\text{Tot}((\bigoplus \mathcal{E}_i^\bullet)(K)) = \bigoplus \text{Tot}(\mathcal{E}_i^\bullet(K))$$

by Modules on Sites, Lemma 29.2. The statement on homotopy colimits is a formal consequence of the fact that $R\Gamma$ is an exact functor of triangulated categories and the fact (just proved) that it commutes with direct sums. \square

09A4 **Remark 18.4.** Let X be a scheme. Because $X_{\text{pro-étale}}$ has enough weakly contractible objects for all K in $D(X_{\text{pro-étale}})$ we have $K = R\lim \tau_{\geq -n} K$ by Cohomology on Sites, Proposition 41.2. Since $R\Gamma$ commutes with $R\lim$ by Injectives, Lemma 13.6 we see that

$$R\Gamma(X, K) = R\lim R\Gamma(X, \tau_{\geq -n} K)$$

in $D(\text{Ab})$. This will allow us to extend some results from bounded below complexes to all complexes.

19. Functoriality of the pro-étale site

09A5 Let $f : X \rightarrow Y$ be a morphism of schemes. The functor $Y_{\text{pro-étale}} \rightarrow X_{\text{pro-étale}}$, $V \mapsto X \times_Y V$ induces a morphism of sites $f_{\text{pro-étale}} : X_{\text{pro-étale}} \rightarrow Y_{\text{pro-étale}}$, see Sites, Proposition 14.6. In fact, we obtain a commutative diagram of morphisms of sites

$$\begin{array}{ccc} X_{\text{pro-étale}} & \xrightarrow{\epsilon} & X_{\text{étale}} \\ f_{\text{pro-étale}} \downarrow & & \downarrow f_{\text{étale}} \\ Y_{\text{pro-étale}} & \xrightarrow{\epsilon} & Y_{\text{étale}} \end{array}$$

where ϵ is as in Section 16. In particular we have $\epsilon^{-1} f_{\text{étale}}^{-1} = f_{\text{pro-étale}}^{-1} \epsilon^{-1}$. Here is the corresponding result for pushforward.

09A6 **Lemma 19.1.** *Let $f : X \rightarrow Y$ be a morphism of schemes.*

- (1) Let \mathcal{F} be a sheaf of sets on $X_{\acute{e}tale}$. Then we have $f_{pro\text{-}\acute{e}tale,*}\epsilon^{-1}\mathcal{F} = \epsilon^{-1}f_{\acute{e}tale,*}\mathcal{F}$.
- (2) Let \mathcal{F} be an abelian sheaf on $X_{\acute{e}tale}$. Then we have $Rf_{pro\text{-}\acute{e}tale,*}\epsilon^{-1}\mathcal{F} = \epsilon^{-1}Rf_{\acute{e}tale,*}\mathcal{F}$.

Proof. Proof of (1). Let \mathcal{F} be a sheaf of sets on $X_{\acute{e}tale}$. There is a canonical map $\epsilon^{-1}f_{\acute{e}tale,*}\mathcal{F} \rightarrow f_{pro\text{-}\acute{e}tale,*}\epsilon^{-1}\mathcal{F}$, see Sites, Section 44. To show it is an isomorphism we may work (Zariski) locally on Y , hence we may assume Y is affine. In this case every object of $Y_{pro\text{-}\acute{e}tale}$ has a covering by objects $V = \lim V_i$ which are limits of affine schemes V_i étale over Y (by Proposition 9.1 for example). Evaluating the map $\epsilon^{-1}f_{\acute{e}tale,*}\mathcal{F} \rightarrow f_{pro\text{-}\acute{e}tale,*}\epsilon^{-1}\mathcal{F}$ on V we obtain a map

$$\operatorname{colim} \Gamma(X \times_Y V_i, \mathcal{F}) \longrightarrow \Gamma(X \times_Y V, \epsilon^*\mathcal{F}).$$

see Lemma 16.1 for the left hand side. By Lemma 16.1 we have

$$\Gamma(X \times_Y V, \epsilon^*\mathcal{F}) = \Gamma(X \times_Y V, \mathcal{F})$$

Hence the result holds by Étale Cohomology, Lemma 51.3.

Proof of (2). Arguing in exactly the same manner as above we see that it suffices to show that

$$\operatorname{colim} H_{\acute{e}tale}^i(X \times_Y V_i, \mathcal{F}) \longrightarrow H_{\acute{e}tale}^i(X \times_Y V, \mathcal{F})$$

which follows once more from Étale Cohomology, Lemma 51.3. \square

20. Finite morphisms and pro-étale sites

- 09A7 It is not clear that a finite morphism of schemes determines an exact pushforward on abelian pro-étale sheaves.
- 09A8 **Lemma 20.1.** *Let $f : Z \rightarrow X$ be a finite morphism of schemes which is locally of finite presentation. Then $f_{pro\text{-}\acute{e}tale,*} : Ab(Z_{pro\text{-}\acute{e}tale}) \rightarrow Ab(X_{pro\text{-}\acute{e}tale})$ is exact.*

Proof. To prove this we may work (Zariski) locally on X and assume that X is affine, say $X = \operatorname{Spec}(A)$. Then $Z = \operatorname{Spec}(B)$ for some finite A -algebra B of finite presentation. The construction in the proof of Proposition 10.3 produces a faithfully flat, ind-étale ring map $A \rightarrow D$ with D w-contractible. We may check exactness of a sequence of sheaves by evaluating on $U = \operatorname{Spec}(D)$ be such an object. Then $f_{pro\text{-}\acute{e}tale,*}\mathcal{F}$ evaluated at U is equal to \mathcal{F} evaluated at $V = \operatorname{Spec}(D \otimes_A B)$. Since $D \otimes_A B$ is w-contractible by Lemma 10.6 evaluation at V is exact. \square

21. Closed immersions and pro-étale sites

- 09A9 It is not clear (and likely false) that a closed immersion of schemes determines an exact pushforward on abelian pro-étale sheaves.
- 09BK **Lemma 21.1.** *Let $i : Z \rightarrow X$ be a closed immersion morphism of affine schemes. Denote X_{app} and Z_{app} the sites introduced in Lemma 11.24. The base change functor*

$$u : X_{app} \rightarrow Z_{app}, \quad U \mapsto u(U) = U \times_X Z$$

is continuous and has a fully faithful left adjoint v . For V in Z_{app} the morphism $V \rightarrow v(V)$ is a closed immersion identifying V with $u(v(V)) = v(V) \times_X Z$ and every point of $v(V)$ specializes to a point of V . The functor v is cocontinuous and sends coverings to coverings.

Proof. The existence of the adjoint follows immediately from Lemma 7.7 and the definitions. It is clear that u is continuous from the definition of coverings in X_{app} .

Write $X = \text{Spec}(A)$ and $Z = \text{Spec}(A/I)$. Let $V = \text{Spec}(\overline{C})$ be an object of Z_{app} and let $v(V) = \text{Spec}(C)$. We have seen in the statement of Lemma 7.7 that V equals $v(V) \times_X Z = \text{Spec}(C/IC)$. Any $g \in C$ which maps to an invertible element of $C/IC = \overline{C}$ is invertible in C . Namely, we have the A -algebra maps $C \rightarrow C_g \rightarrow C/IC$ and by adjointness we obtain an C -algebra map $C_g \rightarrow C$. Thus every point of $v(V)$ specializes to a point of V .

Suppose that $\{V_i \rightarrow V\}$ is a covering in Z_{app} . Then $\{v(V_i) \rightarrow v(V)\}$ is a finite family of morphisms of Z_{app} such that every point of $V \subset v(V)$ is in the image of one of the maps $v(V_i) \rightarrow v(V)$. As the morphisms $v(V_i) \rightarrow v(V)$ are flat (since they are weakly étale) we conclude that $\{v(V_i) \rightarrow v(V)\}$ is jointly surjective. This proves that v sends coverings to coverings.

Let V be an object of Z_{app} and let $\{U_i \rightarrow v(V)\}$ be a covering in X_{app} . Then we see that $\{u(U_i) \rightarrow u(v(V)) = V\}$ is a covering of Z_{app} . By adjointness we obtain morphisms $v(u(U_i)) \rightarrow U_i$. Thus the family $\{v(u(U_i)) \rightarrow v(V)\}$ refines the given covering and we conclude that v is cocontinuous. \square

09BL **Lemma 21.2.** *Let $Z \rightarrow X$ be a closed immersion morphism of affine schemes. The corresponding morphism of topoi $i = i_{pro\text{-}\acute{e}tale}$ is equal to the morphism of topoi associated to the fully faithful cocontinuous functor $v : Z_{app} \rightarrow X_{app}$ of Lemma 21.1. It follows that*

- (1) $i^{-1}\mathcal{F}$ is the sheaf associated to the presheaf $V \mapsto \mathcal{F}(v(V))$,
- (2) for a weakly contractible object V of Z_{app} we have $i^{-1}\mathcal{F}(V) = \mathcal{F}(v(V))$,
- (3) $i^{-1} : \text{Sh}(X_{pro\text{-}\acute{e}tale}) \rightarrow \text{Sh}(Z_{pro\text{-}\acute{e}tale})$ has a left adjoint $i_!^{Sh}$,
- (4) $i^{-1} : \text{Ab}(X_{pro\text{-}\acute{e}tale}) \rightarrow \text{Ab}(Z_{pro\text{-}\acute{e}tale})$ has a left adjoint $i_!$,
- (5) $id \rightarrow i^{-1}i_!^{Sh}$, $id \rightarrow i^{-1}i_!$, and $i^{-1}i_* \rightarrow id$ are isomorphisms, and
- (6) i_* , $i_!^{Sh}$ and $i_!$ are fully faithful.

Proof. By Lemma 11.24 we may describe $i_{pro\text{-}\acute{e}tale}$ in terms of the morphism of sites $u : X_{app} \rightarrow Z_{app}$, $V \mapsto V \times_X Z$. The first statement of the lemma follows from Sites, Lemma 21.2 (but with the roles of u and v reversed).

Proof of (1). By the description of i as the morphism of topoi associated to v this holds by the construction, see Sites, Lemma 20.1.

Proof of (2). Since the functor v sends coverings to coverings by Lemma 21.1 we see that the presheaf $\mathcal{G} : V \mapsto \mathcal{F}(v(V))$ is a separated presheaf (Sites, Definition 10.9). Hence the sheafification of \mathcal{G} is \mathcal{G}^+ , see Sites, Theorem 10.10. Next, let V be a weakly contractible object of Z_{app} . Let $\mathcal{V} = \{V_i \rightarrow V\}_{i=1, \dots, n}$ be any covering in Z_{app} . Set $\mathcal{V}' = \{\coprod V_i \rightarrow V\}$. Since v commutes with finite disjoint unions (as a left adjoint or by the construction) and since \mathcal{F} sends finite disjoint unions into products, we see that

$$H^0(\mathcal{V}, \mathcal{G}) = H^0(\mathcal{V}', \mathcal{G})$$

(notation as in Sites, Section 10; compare with Étale Cohomology, Lemma 22.1). Thus we may assume the covering is given by a single morphism, like so $\{V' \rightarrow V\}$. Since V is weakly contractible, this covering can be refined by the trivial covering $\{V \rightarrow V\}$. It therefore follows that the value of $\mathcal{G}^+ = i^{-1}\mathcal{F}$ on V is simply $\mathcal{F}(v(V))$ and (2) is proved.

Proof of (3). Every object of Z_{app} has a covering by weakly contractible objects (Lemma 11.27). By the above we see that we would have $i_1^{Sh} h_V = h_{v(V)}$ for V weakly contractible if i_1^{Sh} existed. The existence of i_1^{Sh} then follows from Sites, Lemma 23.1.

Proof of (4). Existence of $i_!$ follows in the same way by setting $i_! \mathbf{Z}_V = \mathbf{Z}_{v(V)}$ for V weakly contractible in Z_{app} , using similar for direct sums, and applying Homology, Lemma 26.6. Details omitted.

Proof of (5). Let V be a contractible object of Z_{app} . Then $i^{-1} i_1^{Sh} h_V = i^{-1} h_{v(V)} = h_{u(v(V))} = h_V$. (It is a general fact that $i^{-1} h_U = h_{u(U)}$.) Since the sheaves h_V for V contractible generate $Sh(Z_{app})$ (Sites, Lemma 12.5) we conclude $\text{id} \rightarrow i^{-1} i_1^{Sh}$ is an isomorphism. Similarly for the map $\text{id} \rightarrow i^{-1} i_!$. Then $(i^{-1} i_* \mathcal{H})(V) = i_* \mathcal{H}(v(V)) = \mathcal{H}(u(v(V))) = \mathcal{H}(V)$ and we find that $i^{-1} i_* \rightarrow \text{id}$ is an isomorphism.

The fully faithfulness statements of (6) now follow from Categories, Lemma 24.3. \square

09AA **Lemma 21.3.** *Let $i : Z \rightarrow X$ be a closed immersion of schemes. Then*

- (1) $i_{pro\text{-}\acute{e}tale}^{-1}$ commutes with limits,
- (2) $i_{pro\text{-}\acute{e}tale,*}$ is fully faithful, and
- (3) $i_{pro\text{-}\acute{e}tale}^{-1} i_{pro\text{-}\acute{e}tale,*} \cong \text{id}_{Sh(Z_{pro\text{-}\acute{e}tale})}$.

Proof. Assertions (2) and (3) are equivalent by Sites, Lemma 40.1. Parts (1) and (3) are (Zariski) local on X , hence we may assume that X is affine. In this case the result follows from Lemma 21.2. \square

09AB **Lemma 21.4.** *Let $i : Z \rightarrow X$ be an integral universally injective and surjective morphism of schemes. Then $i_{pro\text{-}\acute{e}tale,*}$ and $i_{pro\text{-}\acute{e}tale}^{-1}$ are quasi-inverse equivalences of categories of pro-étale topoi.*

Proof. There is an immediate reduction to the case that X is affine. Then Z is affine too. Set $A = \mathcal{O}(X)$ and $B = \mathcal{O}(Z)$. Then the categories of étale algebras over A and B are equivalent, see Étale Cohomology, Theorem 45.2 and Remark 45.3. Thus the categories of ind-étale algebras over A and B are equivalent. In other words the categories X_{app} and Z_{app} of Lemma 11.24 are equivalent. We omit the verification that this equivalence sends coverings to coverings and vice versa. Thus the result as Lemma 11.24 tells us the pro-étale topos is the topos of sheaves on X_{app} . \square

09AC **Lemma 21.5.** *Let $i : Z \rightarrow X$ be a closed immersion of schemes. Let $U \rightarrow X$ be an object of $X_{pro\text{-}\acute{e}tale}$ such that*

- (1) U is affine and weakly contractible, and
- (2) every point of U specializes to a point of $U \times_X Z$.

Then $i_{pro\text{-}\acute{e}tale}^{-1} \mathcal{F}(U \times_X Z) = \mathcal{F}(U)$ for all abelian sheaves on $X_{pro\text{-}\acute{e}tale}$.

Proof. Since pullback commutes with restriction, we may replace X by U . Thus we may assume that X is affine and weakly contractible and that every point of X specializes to a point of Z . By Lemma 21.2 part (1) it suffices to show that $v(Z) = X$ in this case. Thus we have to show: If A is a w-contractible ring, $I \subset A$ an ideal contained in the radical of A and $A \rightarrow B \rightarrow A/I$ is a factorization with $A \rightarrow B$ ind-étale, then there is a unique section $B \rightarrow A$ compatible with maps

to A/I . Observe that $B/IB = A/I \times R$ as A/I -algebras. After replacing B by a localization we may assume $B/IB = A/I$. Note that $\text{Spec}(B) \rightarrow \text{Spec}(A)$ is surjective as the image contains $V(I)$ and hence all closed points and is closed under specialization. Since A is w-contractible there is a section $B \rightarrow A$. Since $B/IB = A/I$ this section is compatible with the map to A/I . We omit the proof of uniqueness (hint: use that A and B have isomorphic local rings at maximal ideals of A). \square

09BM **Lemma 21.6.** *Let $i : Z \rightarrow X$ be a closed immersion of schemes. If $X \setminus i(Z)$ is a retrocompact open of X , then $i_{\text{pro-étale},*}$ is exact.*

Proof. The question is local on X hence we may assume X is affine. Say $X = \text{Spec}(A)$ and $Z = \text{Spec}(A/I)$. There exist $f_1, \dots, f_r \in I$ such that $Z = V(f_1, \dots, f_r)$ set theoretically, see Algebra, Lemma 28.1. By Lemma 21.4 we may assume that $Z = \text{Spec}(A/(f_1, \dots, f_r))$. In this case the functor $i_{\text{pro-étale},*}$ is exact by Lemma 20.1. \square

22. Extension by zero

09AD The general material in Modules on Sites, Section 19 allows us to make the following definition.

09AE **Definition 22.1.** Let $j : U \rightarrow X$ be a weakly étale morphism of schemes.

- (1) The restriction functor $j^{-1} : \text{Sh}(X_{\text{pro-étale}}) \rightarrow \text{Sh}(U_{\text{pro-étale}})$ has a left adjoint $j_!^{\text{Sh}} : \text{Sh}(X_{\text{pro-étale}}) \rightarrow \text{Sh}(U_{\text{pro-étale}})$.
- (2) The restriction functor $j^{-1} : \text{Ab}(X_{\text{pro-étale}}) \rightarrow \text{Ab}(U_{\text{pro-étale}})$ has a left adjoint which is denoted $j_! : \text{Ab}(U_{\text{pro-étale}}) \rightarrow \text{Ab}(X_{\text{pro-étale}})$ and called *extension by zero*.
- (3) Let Λ be a ring. The functor $j^{-1} : \text{Mod}(X_{\text{pro-étale}}, \Lambda) \rightarrow \text{Mod}(U_{\text{pro-étale}}, \Lambda)$ has a left adjoint $j_! : \text{Mod}(U_{\text{pro-étale}}, \Lambda) \rightarrow \text{Mod}(X_{\text{pro-étale}}, \Lambda)$ and called *extension by zero*.

As usual we compare this to what happens in the étale case.

09AF **Lemma 22.2.** *Let $j : U \rightarrow X$ be an étale morphism of schemes. Let \mathcal{G} be an abelian sheaf on $U_{\text{étale}}$. Then $\epsilon^{-1}j_!\mathcal{G} = j_!\epsilon^{-1}\mathcal{G}$ as sheaves on $X_{\text{pro-étale}}$.*

Proof. This is true because both are left adjoints to $j_{\text{pro-étale},*}\epsilon^{-1} = \epsilon^{-1}j_{\text{étale},*}$, see Lemma 19.1. \square

09AG **Lemma 22.3.** *Let $j : U \rightarrow X$ be a weakly étale morphism of schemes. Let $i : Z \rightarrow X$ be a closed immersion such that $U \times_X Z = \emptyset$. Let $V \rightarrow X$ be an affine object of $X_{\text{pro-étale}}$ such that every point of V specializes to a point of $V_Z = Z \times_X V$. Then $j_!\mathcal{F}(V) = 0$ for all abelian sheaves on $U_{\text{pro-étale}}$.*

Proof. Let $\{V_i \rightarrow V\}$ be a pro-étale covering. The lemma follows if we can refine this covering to a covering where the members have no morphisms into U over X (see construction of $j_!$ in Modules on Sites, Section 19). First refine the covering to get a finite covering with V_i affine. For each i let $V_i = \text{Spec}(A_i)$ and let $Z_i \subset V_i$ be the inverse image of Z . Set $W_i = \text{Spec}(A_{i,Z_i}^\sim)$ with notation as in Lemma 5.1. Then $\coprod W_i \rightarrow V$ is weakly étale and the image contains all points of V_Z . Hence the image contains all points of V by our assumption on specializations. Thus $\{W_i \rightarrow V\}$ is a pro-étale covering refining the given one. But each point in W_i specializes to a point lying over Z , hence there are no morphisms $W_i \rightarrow U$ over X . \square

09BN **Lemma 22.4.** *Let $j : U \rightarrow X$ be an open immersion of schemes. Then $id \cong j^{-1}j_!$ and $j^{-1}j_* \cong id$ and the functors $j_!$ and j_* are fully faithful.*

Proof. See Sites, Lemma 26.4 and Categories, Lemma 24.3. \square

Here is the relationship between extension by zero and restriction to the complementary closed subscheme.

09AH **Lemma 22.5.** *Let X be a scheme. Let $Z \subset X$ be a closed subscheme and let $U \subset X$ be the complement. Denote $i : Z \rightarrow X$ and $j : U \rightarrow X$ the inclusion morphisms. Assume that j is a quasi-compact morphism. For every abelian sheaf on $X_{pro\text{-}\acute{e}tale}$ there is a canonical short exact sequence*

$$0 \rightarrow j_!j^{-1}\mathcal{F} \rightarrow \mathcal{F} \rightarrow i_*i^{-1}\mathcal{F} \rightarrow 0$$

on $X_{pro\text{-}\acute{e}tale}$ where all the functors are for the pro-étale topology.

Proof. We obtain the maps by the adjointness properties of the functors involved. It suffices to show that $X_{pro\text{-}\acute{e}tale}$ has enough objects (Sites, Definition 39.2) on which the sequence evaluates to a short exact sequence. Let $V = \text{Spec}(A)$ be an affine object of $X_{pro\text{-}\acute{e}tale}$ such that A is w-contractible (there are enough objects of this type). Then $V \times_X Z$ is cut out by an ideal $I \subset A$. The assumption that j is quasi-compact implies there exist $f_1, \dots, f_r \in I$ such that $V(I) = V(f_1, \dots, f_r)$. We obtain a faithfully flat, ind-Zariski ring map

$$A \longrightarrow A_{f_1} \times \dots \times A_{f_r} \times A_{\tilde{V}(I)}$$

with $A_{\tilde{V}(I)}$ as in Lemma 5.1. Since $V_i = \text{Spec}(A_{f_i}) \rightarrow X$ factors through U we have

$$j_!j^{-1}\mathcal{F}(V_i) = \mathcal{F}(V_i) \quad \text{and} \quad i_*i^{-1}\mathcal{F}(V_i) = 0$$

On the other hand, for the scheme $V^\sim = \text{Spec}(A_{\tilde{V}(I)})$ we have

$$j_!j^{-1}\mathcal{F}(V^\sim) = 0 \quad \text{and} \quad \mathcal{F}(V^\sim) = i_*i^{-1}\mathcal{F}(V^\sim)$$

the first equality by Lemma 22.3 and the second by Lemmas 21.5 and 10.7. Thus the sequence evaluates to an exact sequence on $\text{Spec}(A_{f_1} \times \dots \times A_{f_r} \times A_{\tilde{V}(I)})$ and the lemma is proved. \square

09BP **Lemma 22.6.** *Let $j : U \rightarrow X$ be a quasi-compact open immersion morphism of schemes. The functor $j_! : Ab(U_{pro\text{-}\acute{e}tale}) \rightarrow Ab(X_{pro\text{-}\acute{e}tale})$ commutes with limits.*

Proof. Since $j_!$ is exact it suffices to show that $j_!$ commutes with products. The question is local on X , hence we may assume X affine. Let \mathcal{G} be an abelian sheaf on $U_{pro\text{-}\acute{e}tale}$. Note that there always is a canonical map

$$j_!\mathcal{G} \rightarrow j_*\mathcal{G}$$

see Modules on Sites, Remark 19.8. In our particular case this map can be obtained from the fact that $j^{-1}j_*\mathcal{G} = \mathcal{G}$. Hence applying the exact sequence of Lemma 22.5 we get

$$0 \rightarrow j_!\mathcal{G} \rightarrow j_*\mathcal{G} \rightarrow i_*i^{-1}j_*\mathcal{G} \rightarrow 0$$

where $i : Z \rightarrow X$ is the inclusion of the reduced induced scheme structure on the complement $Z = X \setminus U$. The functors j_* and i_* commute with products as right adjoints. The functor i^{-1} commutes with products by Lemma 21.3. Hence $j_!$ does because on the pro-étale site products are exact (Cohomology on Sites, Proposition 41.2). \square

23. Constructible sheaves on the pro-étale site

09AI We stick to constructible sheaves of Λ -modules for a Noetherian ring. In the future we intend to discuss constructible sheaves of sets, groups, etc.

09AJ **Definition 23.1.** Let X be a scheme. Let Λ be a Noetherian ring. A sheaf of Λ -modules on $X_{\text{pro-étale}}$ is *constructible* if for every affine open $U \subset X$ there exists a finite decomposition of U into constructible locally closed subschemes $U = \coprod_i U_i$ such that $\mathcal{F}|_{U_i}$ is of finite type and locally constant for all i .

Again this does not give anything “new”.

09AK **Lemma 23.2.** *Let X be a scheme. Let Λ be a Noetherian ring. The functor ϵ^{-1} defines an equivalence of categories*

$$\left\{ \begin{array}{l} \text{constructible sheaves of} \\ \Lambda\text{-modules on } X_{\text{étale}} \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{l} \text{constructible sheaves of} \\ \Lambda\text{-modules on } X_{\text{pro-étale}} \end{array} \right\}$$

between constructible sheaves of Λ -modules on $X_{\text{étale}}$ and constructible sheaves of Λ -modules on $X_{\text{pro-étale}}$.

Proof. By Lemma 16.2 the functor ϵ^{-1} is fully faithful and commutes with pullback (restriction) to the strata. Hence ϵ^{-1} of a constructible étale sheaf is a constructible pro-étale sheaf. To finish the proof let \mathcal{F} be a constructible sheaf of Λ -modules on $X_{\text{pro-étale}}$ as in Definition 23.1. There is a canonical map

$$\epsilon^{-1}\epsilon_*\mathcal{F} \longrightarrow \mathcal{F}$$

We will show this map is an isomorphism. This will prove that \mathcal{F} is in the essential image of ϵ^{-1} and finish the proof (details omitted).

To prove this we may assume that X is affine. In this case we have a finite partition $X = \coprod_i X_i$ by constructible locally closed strata such that $\mathcal{F}|_{X_i}$ is locally constant of finite type. Let $U \subset X$ be one of the open strata in the partition and let $Z \subset X$ be the reduced induced structure on the complement. By Lemma 22.5 we have a short exact sequence

$$0 \rightarrow j_!j^{-1}\mathcal{F} \rightarrow \mathcal{F} \rightarrow i_*i^{-1}\mathcal{F} \rightarrow 0$$

on $X_{\text{pro-étale}}$. Functoriality gives a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \epsilon^{-1}\epsilon_*j_!j^{-1}\mathcal{F} & \longrightarrow & \epsilon^{-1}\epsilon_*\mathcal{F} & \longrightarrow & \epsilon^{-1}\epsilon_*i_*i^{-1}\mathcal{F} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & j_!j^{-1}\mathcal{F} & \longrightarrow & \mathcal{F} & \longrightarrow & i_*i^{-1}\mathcal{F} \longrightarrow 0 \end{array}$$

By induction on the length of the partition we know that on the one hand $\epsilon^{-1}\epsilon_*i^{-1}\mathcal{F} \rightarrow i^{-1}\mathcal{F}$ and $\epsilon^{-1}\epsilon_*j^{-1}\mathcal{F} \rightarrow j^{-1}\mathcal{F}$ are isomorphisms and on the other that $i^{-1}\mathcal{F} = \epsilon^{-1}\mathcal{A}$ and $j^{-1}\mathcal{F} = \epsilon^{-1}\mathcal{B}$ for some constructible sheaves of Λ -modules \mathcal{A} on $Z_{\text{étale}}$ and \mathcal{B} on $U_{\text{étale}}$. Then

$$\epsilon^{-1}\epsilon_*j_!j^{-1}\mathcal{F} = \epsilon^{-1}\epsilon_*j_!\epsilon^{-1}\mathcal{B} = \epsilon^{-1}\epsilon_*\epsilon^{-1}j_!\mathcal{B} = \epsilon^{-1}j_!\mathcal{B} = j_!\epsilon^{-1}\mathcal{B} = j_!j^{-1}\mathcal{F}$$

the second equality by Lemma 22.2, the third equality by Lemma 16.2, and the fourth equality by Lemma 22.2 again. Similarly, we have

$$\epsilon^{-1}\epsilon_*i_*i^{-1}\mathcal{F} = \epsilon^{-1}\epsilon_*i_*\epsilon^{-1}\mathcal{A} = \epsilon^{-1}\epsilon_*\epsilon^{-1}i_*\mathcal{A} = \epsilon^{-1}i_*\mathcal{A} = i_*\epsilon^{-1}\mathcal{A} = i_*i^{-1}\mathcal{F}$$

this time using Lemma 19.1. By the five lemma we conclude the vertical map in the middle of the big diagram is an isomorphism. \square

09B5 **Lemma 23.3.** *Let X be a scheme. Let Λ be a Noetherian ring. The category of constructible sheaves of Λ -modules on $X_{\text{pro-étale}}$ is a weak Serre subcategory of $\text{Mod}(X_{\text{pro-étale}}, \Lambda)$.*

Proof. This is a formal consequence of Lemmas 23.2 and 16.7 and the result for the étale site (Étale Cohomology, Lemma 70.6). \square

09AL **Lemma 23.4.** *Let X be a scheme. Let Λ be a Noetherian ring. Let $D_c(X_{\text{étale}}, \Lambda)$, resp. $D_c(X_{\text{pro-étale}}, \Lambda)$ be the full subcategory of $D(X_{\text{étale}}, \Lambda)$, resp. $D(X_{\text{pro-étale}}, \Lambda)$ consisting of those complexes whose cohomology sheaves are constructible sheaves of Λ -modules. Then*

$$\epsilon^{-1} : D_c^+(X_{\text{étale}}, \Lambda) \longrightarrow D_c^+(X_{\text{pro-étale}}, \Lambda)$$

is an equivalence of categories.

Proof. The categories $D_c(X_{\text{étale}}, \Lambda)$ and $D_c(X_{\text{pro-étale}}, \Lambda)$ are strictly full, saturated, triangulated subcategories of $D(X_{\text{étale}}, \Lambda)$ and $D(X_{\text{pro-étale}}, \Lambda)$ by Étale Cohomology, Lemma 70.6 and Lemma 23.3 and Derived Categories, Section 13. The statement of the lemma follows by combining Lemmas 16.7 and 23.2. \square

09BQ **Lemma 23.5.** *Let X be a scheme. Let Λ be a Noetherian ring. Let $K, L \in D_c^-(X_{\text{pro-étale}}, \Lambda)$. Then $K \otimes_{\Lambda}^{\mathbf{L}} L$ is in $D_c^-(X_{\text{pro-étale}}, \Lambda)$.*

Proof. Note that $H^i(K \otimes_{\Lambda}^{\mathbf{L}} L)$ is the same as $H^i(\tau_{\geq i-1} K \otimes_{\Lambda}^{\mathbf{L}} \tau_{\geq i-1} L)$. Thus we may assume K and L are bounded. In this case we can apply Lemma 23.4 to reduce to the case of the étale site, see Étale Cohomology, Lemma 96.6. \square

09BR **Lemma 23.6.** *Let X be a scheme. Let Λ be a Noetherian ring. Let K be an object of $D(X_{\text{pro-étale}}, \Lambda)$. Set $K_n = K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n$. If K_1 is in $D_c^-(X_{\text{pro-étale}}, \Lambda/I)$, then K_n is in $D_c^-(X_{\text{pro-étale}}, \Lambda/I^n)$ for all n .*

Proof. Consider the distinguished triangles

$$K \otimes_{\Lambda}^{\mathbf{L}} \underline{I^n/I^{n+1}} \rightarrow K_{n+1} \rightarrow K_n \rightarrow K \otimes_{\Lambda}^{\mathbf{L}} \underline{I^n/I^{n+1}}[1]$$

and the isomorphisms

$$K \otimes_{\Lambda}^{\mathbf{L}} \underline{I^n/I^{n+1}} = K_1 \otimes_{\Lambda/I}^{\mathbf{L}} \underline{I^n/I^{n+1}}$$

By Lemma 23.5 we see that this tensor product has constructible cohomology sheaves (and vanishing when K_1 has vanishing cohomology). Hence by induction on n using Lemma 23.3 we see that each K_n has constructible cohomology sheaves. \square

24. Constructible adic sheaves

09BS In this section we define the notion of a constructible Λ -sheaf as well as some variants.

09BT **Definition 24.1.** Let Λ be a Noetherian ring and let $I \subset \Lambda$ be an ideal. Let X be a scheme. Let \mathcal{F} be a sheaf of Λ -modules on $X_{\text{pro-étale}}$.

- (1) We say \mathcal{F} is a *constructible Λ -sheaf* if $\mathcal{F} = \lim \mathcal{F}/I^n \mathcal{F}$ and each $\mathcal{F}/I^n \mathcal{F}$ is a constructible sheaf of Λ/I^n -modules.

- (2) If \mathcal{F} is a constructible Λ -sheaf, then we say \mathcal{F} is *lisse* if each $\mathcal{F}/I^n\mathcal{F}$ is locally constant.
- (3) We say \mathcal{F} is *adic lisse*³ if there exists a I -adically complete Λ -module M with M/IM finite such that \mathcal{F} is locally isomorphic to

$$\underline{M}^\wedge = \varinjlim M/I^n M.$$

- (4) We say \mathcal{F} is *adic constructible*⁴ if for every affine open $U \subset X$ there exists a decomposition $U = \coprod U_i$ into constructible locally closed subschemes such that $\mathcal{F}|_{U_i}$ is adic lisse.

The definition of a constructible Λ -sheaf is equivalent to the one in [Gro77, Exposé VI, Definition 1.1.1] when $\Lambda = \mathbf{Z}_\ell$ and $I = (\ell)$. It is clear that we have the implications

$$\begin{array}{ccc} \text{lisse adic} & \xlongequal{\quad} & \text{adic constructible} \\ \Downarrow & & \Downarrow \\ \text{lisse constructible } \Lambda\text{-sheaf} & \xlongequal{\quad} & \text{constructible } \Lambda\text{-sheaf} \end{array}$$

The vertical arrows can be inverted in some cases (see Lemmas 24.2 and 24.5). In general neither the category of adic constructible sheaves nor the category of adic constructible sheaves is closed under kernels and cokernels.

Namely, let X be an affine scheme whose underlying topological space $|X|$ is homeomorphic to $\Lambda = \mathbf{Z}_\ell$, see Example 6.3. Denote $f : |X| \rightarrow \mathbf{Z}_\ell = \Lambda$ a homeomorphism. We can think of f as a section of $\underline{\Lambda}^\wedge$ over X and multiplication by f then defines a two term complex

$$\underline{\Lambda}^\wedge \xrightarrow{f} \underline{\Lambda}^\wedge$$

on $X_{\text{pro-étale}}$. The sheaf $\underline{\Lambda}^\wedge$ is adic lisse. However, the cokernel of the map above, is not adic constructible, as the isomorphism type of the stalks of this cokernel attains infinitely many values: $\mathbf{Z}/\ell^n\mathbf{Z}$ and \mathbf{Z}_ℓ . The cokernel is a constructible \mathbf{Z}_ℓ -sheaf. However, the kernel is not even a constructible \mathbf{Z}_ℓ -sheaf as it is zero a non-quasi-compact open but not zero.

09BU **Lemma 24.2.** *Let X be a Noetherian scheme. Let Λ be a Noetherian ring and let $I \subset \Lambda$ be an ideal. Let \mathcal{F} be a constructible Λ -sheaf on $X_{\text{pro-étale}}$. Then there exists a finite partition $X = \coprod X_i$ by locally closed subschemes such that the restriction $\mathcal{F}|_{X_i}$ is lisse.*

Proof. Let $R = \bigoplus I^n/I^{n+1}$. Observe that R is a Noetherian ring. Since each of the sheaves $\mathcal{F}/I^n\mathcal{F}$ is a constructible sheaf of $\Lambda/I^n\Lambda$ -modules also $I^n\mathcal{F}/I^{n+1}\mathcal{F}$ is a constructible sheaf of Λ/I -modules and hence the pullback of a constructible sheaf \mathcal{G}_n on $X_{\text{étale}}$ by Lemma 23.2. Set $\mathcal{G} = \bigoplus \mathcal{G}_n$. This is a sheaf of R -modules on $X_{\text{étale}}$ and the map

$$\mathcal{G}_0 \otimes_{\Lambda/I} \underline{R} \longrightarrow \mathcal{G}$$

is surjective because the maps

$$\mathcal{F}/I\mathcal{F} \otimes \underline{I^n/I^{n+1}} \rightarrow I^n\mathcal{F}/I^{n+1}\mathcal{F}$$

³This may be nonstandard notation.

⁴This may be nonstandard notation.

are surjective. Hence \mathcal{G} is a constructible sheaf of R -modules by Étale Cohomology, Proposition 73.1. Choose a partition $X = \coprod X_i$ such that $\mathcal{G}|_{X_i}$ is a locally constant sheaf of R -modules of finite type (Étale Cohomology, Lemma 70.2). We claim this is a partition as in the lemma. Namely, replacing X by X_i we may assume \mathcal{G} is locally constant. It follows that each of the sheaves $I^n\mathcal{F}/I^{n+1}\mathcal{F}$ is locally constant. Using the short exact sequences

$$0 \rightarrow I^n\mathcal{F}/I^{n+1}\mathcal{F} \rightarrow \mathcal{F}/I^{n+1}\mathcal{F} \rightarrow \mathcal{F}/I^n\mathcal{F} \rightarrow 0$$

induction and Modules on Sites, Lemma 42.5 the lemma follows. \square

09BV **Lemma 24.3.** *Let X be a weakly contractible affine scheme. Let Λ be a Noetherian ring and $I \subset \Lambda$ be an ideal. Let \mathcal{F} be a sheaf of Λ -modules on $X_{\text{pro-étale}}$ such that*

- (1) $\mathcal{F} = \varinjlim \mathcal{F}/I^n\mathcal{F}$,
- (2) $\mathcal{F}/I^n\mathcal{F}$ is a constant sheaf of Λ/I^n -modules,
- (3) $\mathcal{F}/I\mathcal{F}$ is of finite type.

Then $\mathcal{F} \cong \underline{M}^\wedge$ where M is a finite Λ^\wedge -module.

Proof. Pick a Λ/I^n -module M_n such that $\mathcal{F}/I^n\mathcal{F} \cong \underline{M}_n$. Since we have the surjections $\mathcal{F}/I^{n+1}\mathcal{F} \rightarrow \mathcal{F}/I^n\mathcal{F}$ we conclude that there exist surjections $M_{n+1} \rightarrow M_n$ inducing isomorphisms $M_{n+1}/I^n M_{n+1} \rightarrow M_n$. Fix a choice of such surjections and set $M = \varinjlim M_n$. Then M is an I -adically complete Λ -module with $M/I^n M = M_n$, see Algebra, Lemma 97.1. Since M_1 is a finite type Λ -module (Modules on Sites, Lemma 41.5) we see that M is a finite Λ^\wedge -module. Consider the sheaves

$$\mathcal{I}_n = \text{Isom}(\underline{M}_n, \mathcal{F}/I^n\mathcal{F})$$

on $X_{\text{pro-étale}}$. Modding out by I^n defines a transition map

$$\mathcal{I}_{n+1} \longrightarrow \mathcal{I}_n$$

By our choice of M_n the sheaf \mathcal{I}_n is a torsor under

$$\text{Isom}(\underline{M}_n, \underline{M}_n) = \text{Isom}_\Lambda(M_n, M_n)$$

(Modules on Sites, Lemma 42.4) since $\mathcal{F}/I^n\mathcal{F}$ is (étale) locally isomorphic to \underline{M}_n . It follows from More on Algebra, Lemma 86.1 that the system of sheaves (\mathcal{I}_n) is Mittag-Leffler. For each n let $\mathcal{I}'_n \subset \mathcal{I}_n$ be the image of $\mathcal{I}_N \rightarrow \mathcal{I}_n$ for all $N \gg n$. Then

$$\dots \rightarrow \mathcal{I}'_3 \rightarrow \mathcal{I}'_2 \rightarrow \mathcal{I}'_1 \rightarrow *$$

is a sequence of sheaves of sets on $X_{\text{pro-étale}}$ with surjective transition maps. Since $*(X)$ is a singleton (not empty) and since evaluating at X transforms surjective maps of sheaves of sets into surjections of sets, we can pick $s \in \varinjlim \mathcal{I}'_n(X)$. The sections define isomorphisms $\underline{M}^\wedge \rightarrow \varinjlim \mathcal{F}/I^n\mathcal{F} = \mathcal{F}$ and the proof is done. \square

09BW **Lemma 24.4.** *Let X be a connected scheme. Let Λ be a Noetherian ring and let $I \subset \Lambda$ be an ideal. If \mathcal{F} is a lisse constructible Λ -sheaf on $X_{\text{pro-étale}}$, then \mathcal{F} is adic lisse.*

Proof. By Lemma 16.8 we have $\mathcal{F}/I^n\mathcal{F} = \epsilon^{-1}\mathcal{G}_n$ for some locally constant sheaf \mathcal{G}_n of Λ/I^n -modules. By Étale Cohomology, Lemma 63.8 there exists a finite Λ/I^n -module M_n such that \mathcal{G}_n is locally isomorphic to \underline{M}_n . Choose a covering $\{W_t \rightarrow X\}_{t \in T}$ with each W_t affine and weakly contractible. Then $\mathcal{F}|_{W_t}$ satisfies the assumptions of Lemma 24.3 and hence $\mathcal{F}|_{W_t} \cong \underline{N}_t^\wedge$ for some finite Λ^\wedge -module

N_t . Note that $N_t/I^n N_t \cong M_n$ for all t and n . Hence $N_t \cong N_{t'}$ for all $t, t' \in T$, see More on Algebra, Lemma 86.2. This proves that \mathcal{F} is adic lisse. \square

09BX **Lemma 24.5.** *Let X be a Noetherian scheme. Let Λ be a Noetherian ring and let $I \subset \Lambda$ be an ideal. Let \mathcal{F} be a constructible Λ -sheaf on $X_{\text{pro-étale}}$. Then \mathcal{F} is adic constructible.*

Proof. This is a consequence of Lemmas 24.2 and 24.4, the fact that a Noetherian scheme is locally connected (Topology, Lemma 9.6), and the definitions. \square

It will be useful to identify the constructible Λ -sheaves inside the category of derived complete sheaves of Λ -modules. It turns out that the naive analogue of More on Algebra, Lemma 82.5 is wrong in this setting. However, here is the analogue of More on Algebra, Lemma 82.4.

09BY **Lemma 24.6.** *Let X be a scheme. Let Λ be a ring and let $I \subset \Lambda$ be a finitely generated ideal. Let \mathcal{F} be a sheaf of Λ -modules on $X_{\text{pro-étale}}$. If \mathcal{F} is derived complete and $\mathcal{F}/I\mathcal{F} = 0$, then $\mathcal{F} = 0$.*

Proof. Assume that $\mathcal{F}/I\mathcal{F}$ is zero. Let $I = (f_1, \dots, f_r)$. Let $i < r$ be the largest integer such that $\mathcal{G} = \mathcal{F}/(f_1, \dots, f_i)\mathcal{F}$ is nonzero. If i does not exist, then $\mathcal{F} = 0$ which is what we want to show. Then \mathcal{G} is derived complete as a cokernel of a map between derived complete modules, see Proposition 15.1. By our choice of i we have that $f_{i+1} : \mathcal{G} \rightarrow \mathcal{G}$ is surjective. Hence

$$\lim(\dots \rightarrow \mathcal{G} \xrightarrow{f_{i+1}} \mathcal{G} \xrightarrow{f_{i+1}} \mathcal{G})$$

is nonzero, contradicting the derived completeness of \mathcal{G} . \square

09BZ **Lemma 24.7.** *Let X be a weakly contractible affine scheme. Let Λ be a Noetherian ring and let $I \subset \Lambda$ be an ideal. Let \mathcal{F} be a derived complete sheaf of Λ -modules on $X_{\text{pro-étale}}$ with $\mathcal{F}/I\mathcal{F}$ a locally constant sheaf of Λ/I -modules of finite type. Then there exists an integer t and a surjective map*

$$(\underline{\Lambda}^\wedge)^{\oplus t} \rightarrow \mathcal{F}$$

Proof. Since X is weakly contractible, there exists a finite disjoint open covering $X = \coprod U_i$ such that $\mathcal{F}/I\mathcal{F}|_{U_i}$ is isomorphic to the constant sheaf associated to a finite Λ/I -module M_i . Choose finitely many generators m_{ij} of M_i . We can find sections $s_{ij} \in \mathcal{F}(X)$ restricting to m_{ij} viewed as a section of $\mathcal{F}/I\mathcal{F}$ over U_i . Let t be the total number of s_{ij} . Then we obtain a map

$$\alpha : \underline{\Lambda}^{\oplus t} \rightarrow \mathcal{F}$$

which is surjective modulo I by construction. By Lemma 14.1 the derived completion of $\underline{\Lambda}^{\oplus t}$ is the sheaf $(\underline{\Lambda}^\wedge)^{\oplus t}$. Since \mathcal{F} is derived complete we see that α factors through a map

$$\alpha^\wedge : (\underline{\Lambda}^\wedge)^{\oplus t} \rightarrow \mathcal{F}$$

Then $\mathcal{Q} = \text{Coker}(\alpha^\wedge)$ is a derived complete sheaf of Λ -modules by Proposition 15.1. By construction $\mathcal{Q}/I\mathcal{Q} = 0$. It follows from Lemma 24.6 that $\mathcal{Q} = 0$ which is what we wanted to show. \square

25. A suitable derived category

09C0 Let X be a scheme. It will turn out that for many schemes X a suitable derived category of ℓ -adic sheaves can be gotten by considering the derived complete objects K of $D(X_{\text{pro-étale}}, \Lambda)$ with the property that $K \otimes_{\Lambda}^{\mathbf{L}} \mathbf{F}_{\ell}$ is bounded with constructible cohomology sheaves. Here is the general definition.

09C1 **Definition 25.1.** Let Λ be a Noetherian ring and let $I \subset \Lambda$ be an ideal. Let X be a scheme. An object K of $D(X_{\text{pro-étale}}, \Lambda)$ is called *constructible* if

- (1) K is derived complete with respect to I ,
- (2) $K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I$ has constructible cohomology sheaves and locally has finite tor dimension.

We denote $D_{\text{cons}}(X, \Lambda)$ the full subcategory of constructible K in $D(X_{\text{pro-étale}}, \Lambda)$.

Recall that with our conventions a complex of finite tor dimension is bounded (Cohomology on Sites, Definition 38.1). In fact, let's collect everything proved so far in a lemma.

09C2 **Lemma 25.2.** *In the situation above suppose K is in $D_{\text{cons}}(X, \Lambda)$ and X is quasi-compact. Set $K_n = K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n$. There exist a, b such that*

- (1) $K = R\lim K_n$ and $H^i(K) = 0$ for $i \notin [a, b]$,
- (2) each K_n has tor amplitude in $[a, b]$,
- (3) each K_n has constructible cohomology sheaves,
- (4) each $K_n = \epsilon^{-1}L_n$ for some $L_n \in D_{\text{ctf}}(X_{\text{étale}}, \Lambda/I^n)$ (*Étale Cohomology, Definition 96.7*).

Proof. By definition of local having finite tor dimension, we can find a, b such that K_1 has tor amplitude in $[a, b]$. Part (2) follows from Cohomology on Sites, Lemma 38.8. Then (1) follows as K is derived complete by the description of limits in Cohomology on Sites, Proposition 41.2 and the fact that $H^b(K_{n+1}) \rightarrow H^b(K_n)$ is surjective as $K_n = K_{n+1} \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n$. Part (3) follows from Lemma 23.6, Part (4) follows from Lemma 23.4 and the fact that L_n has finite tor dimension because K_n does (small argument omitted). \square

09C3 **Lemma 25.3.** *Let X be a weakly contractible affine scheme. Let Λ be a Noetherian ring and let $I \subset \Lambda$ be an ideal. Let K be an object of $D_{\text{cons}}(X, \Lambda)$ such that the cohomology sheaves of $K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I$ are locally constant. Then there exists a finite disjoint open covering $X = \coprod U_i$ and for each i a finite collection of finite projective Λ^{\wedge} -modules M_a, \dots, M_b such that $K|_{U_i}$ is represented by a complex*

$$(\underline{M}^a)^{\wedge} \rightarrow \dots \rightarrow (\underline{M}^b)^{\wedge}$$

in $D(U_{i,\text{pro-étale}}, \Lambda)$ for some maps of sheaves of Λ -modules $(\underline{M}^i)^{\wedge} \rightarrow (\underline{M}^{i+1})^{\wedge}$.

Proof. We freely use the results of Lemma 25.2. Choose a, b as in that lemma. We will prove the lemma by induction on $b - a$. Let $\mathcal{F} = H^b(K)$. Note that \mathcal{F} is a derived complete sheaf of Λ -modules by Proposition 15.1. Moreover $\mathcal{F}/I\mathcal{F}$ is a locally constant sheaf of Λ/I -modules of finite type. Apply Lemma 24.7 to get a surjection $\rho : (\underline{\Lambda}^{\wedge})^{\oplus t} \rightarrow \mathcal{F}$.

If $a = b$, then $K = \mathcal{F}[-b]$. In this case we see that

$$\mathcal{F} \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I = \mathcal{F}/I\mathcal{F}$$

As X is weakly contractible and $\mathcal{F}/I\mathcal{F}$ locally constant, we can find a finite disjoint union decomposition $X = \coprod U_i$ by affine opens U_i and Λ/I -modules \overline{M}_i such that $\mathcal{F}/I\mathcal{F}$ restricts to \overline{M}_i on U_i . After refining the covering we may assume the map

$$\rho|_{U_i \bmod I} : \underline{\Lambda/I}^{\oplus t} \longrightarrow \overline{M}_i$$

is equal to α_i for some surjective module map $\alpha_i : \Lambda/I^{\oplus t} \rightarrow \overline{M}_i$, see Modules on Sites, Lemma 42.3. Note that each \overline{M}_i is a finite Λ/I -module. Since $\mathcal{F}/I\mathcal{F}$ has tor amplitude in $[0, 0]$ we conclude that \overline{M}_i is a flat Λ/I -module. Hence \overline{M}_i is finite projective (Algebra, Lemma 77.2). Hence we can find a projector $\overline{p}_i : (\Lambda/I)^{\oplus t} \rightarrow (\Lambda/I)^{\oplus t}$ whose image maps isomorphically to \overline{M}_i under the map α_i . We can lift \overline{p}_i to a projector $p_i : (\Lambda^\wedge)^{\oplus t} \rightarrow (\Lambda^\wedge)^{\oplus t}$ ⁵. Then $M_i = \text{Im}(p_i)$ is a finite I -adically complete Λ^\wedge -module with $M_i/IM_i = \overline{M}_i$. Over U_i consider the maps

$$\underline{M}_i^\wedge \rightarrow (\underline{\Lambda}^\wedge)^{\oplus t} \rightarrow \mathcal{F}|_{U_i}$$

By construction the composition induces an isomorphism modulo I . The source and target are derived complete, hence so are the cokernel \mathcal{Q} and the kernel \mathcal{K} . We have $\mathcal{Q}/I\mathcal{Q} = 0$ by construction hence \mathcal{Q} is zero by Lemma 24.6. Then

$$0 \rightarrow \mathcal{K}/I\mathcal{K} \rightarrow \overline{M}_i \rightarrow \mathcal{F}/I\mathcal{F} \rightarrow 0$$

is exact by the vanishing of Tor_1 see at the start of this paragraph; also use that $\underline{\Lambda}^\wedge/I\overline{\Lambda}^\wedge$ by Modules on Sites, Lemma 41.4 to see that $\underline{M}_i^\wedge/IM_i^\wedge = \overline{M}_i$. Hence $\mathcal{K}/I\mathcal{K} = 0$ by construction and we conclude that $\mathcal{K} = 0$ as before. This proves the result in case $a = b$.

If $b > a$, then we lift the map ρ to a map

$$\tilde{\rho} : (\underline{\Lambda}^\wedge)^{\oplus t}[-b] \longrightarrow K$$

in $D(X_{\text{pro-étale}}, \Lambda)$. This is possible as we can think of K as a complex of $\underline{\Lambda}^\wedge$ -modules by discussion in the introduction to Section 14 and because $X_{\text{pro-étale}}$ is weakly contractible hence there is no obstruction to lifting the elements $\rho(e_s) \in H^0(X, \mathcal{F})$ to elements of $H^b(X, K)$. Fitting $\tilde{\rho}$ into a distinguished triangle

$$(\underline{\Lambda}^\wedge)^{\oplus t}[-b] \rightarrow K \rightarrow L \rightarrow (\underline{\Lambda}^\wedge)^{\oplus t}[-b+1]$$

we see that L is an object of $D_{\text{cons}}(X, \Lambda)$ such that $L \otimes_{\Lambda}^L \underline{\Lambda}/I$ has tor amplitude contained in $[a, b-1]$ (details omitted). By induction we can describe L locally as stated in the lemma, say L is isomorphic to

$$(\underline{M}^a)^\wedge \rightarrow \dots \rightarrow (\underline{M}^{b-1})^\wedge$$

The map $L \rightarrow (\underline{\Lambda}^\wedge)^{\oplus t}[-b+1]$ corresponds to a map $(\underline{M}^{b-1})^\wedge \rightarrow (\underline{\Lambda}^\wedge)^{\oplus t}$ which allows us to extend the complex by one. The corresponding complex is isomorphic to K in the derived category by the properties of triangulated categories. This finishes the proof. \square

Motivated by what happens for constructible Λ -sheaves we introduce the following notion.

09C4 **Definition 25.4.** Let X be a scheme. Let Λ be a Noetherian ring and let $I \subset \Lambda$ be an ideal. Let $K \in D(X_{\text{pro-étale}}, \Lambda)$.

⁵Proof: by Algebra, Lemma 31.6 we can lift \overline{p}_i to a compatible system of projectors $p_{i,n} : (\Lambda/I^n)^{\oplus t} \rightarrow (\Lambda/I^n)^{\oplus t}$ and then we set $p_i = \lim p_{i,n}$ which works because $\Lambda^\wedge = \lim \Lambda/I^n$.

- (1) We say K is *adic lisse*⁶ if there exists a finite complex of finite projective Λ^\wedge -modules M^\bullet such that K is locally isomorphic to

$$\underline{M}^{a^\wedge} \rightarrow \dots \rightarrow \underline{M}^{b^\wedge}$$

- (2) We say K is *adic constructible*⁷ if for every affine open $U \subset X$ there exists a decomposition $U = \coprod U_i$ into constructible locally closed subschemes such that $K|_{U_i}$ is adic lisse.

The difference between the local structure obtained in Lemma 25.3 and the structure of an adic lisse complex is that the maps $\underline{M}^{i^\wedge} \rightarrow \underline{M}^{i+1^\wedge}$ in Lemma 25.3 need not be constant, whereas in the definition above they are required to be constant.

09C5 **Lemma 25.5.** *Let X be a weakly contractible affine scheme. Let Λ be a Noetherian ring and let $I \subset \Lambda$ be an ideal. Let K be an object of $D_{\text{cons}}(X, \Lambda)$ such that $K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n$ is isomorphic in $D(X_{\text{pro-étale}}, \Lambda/I^n)$ to a complex of constant sheaves of Λ/I^n -modules. Then*

$$H^0(X, K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n)$$

has the Mittag-Leffler condition.

Proof. Say $K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n$ is isomorphic to \underline{E}_n for some object E_n of $D(\Lambda/I^n)$. Since $K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I$ has finite tor dimension and has finite type cohomology sheaves we see that E_1 is perfect (see More on Algebra, Lemma 67.2). The transition maps

$$K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^{n+1} \rightarrow K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n$$

locally come from (possibly many distinct) maps of complexes $E_{n+1} \rightarrow E_n$ in $D(\Lambda/I^{n+1})$ see Cohomology on Sites, Lemma 43.3. For each n choose one such map and observe that it induces an isomorphism $E_{n+1} \otimes_{\Lambda/I^{n+1}}^{\mathbf{L}} \Lambda/I^n \rightarrow E_n$ in $D(\Lambda/I^n)$. By More on Algebra, Lemma 83.4 we can find a finite complex M^\bullet of finite projective Λ^\wedge -modules and isomorphisms $M^\bullet/I^n M^\bullet \rightarrow E_n$ in $D(\Lambda/I^n)$ compatible with the transition maps.

Now observe that for each finite collection of indices $n > m > k$ the triple of maps

$$H^0(X, K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n) \rightarrow H^0(X, K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^m) \rightarrow H^0(X, K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^k)$$

is isomorphic to

$$H^0(X, \underline{M^\bullet/I^n M^\bullet}) \rightarrow H^0(X, \underline{M^\bullet/I^m M^\bullet}) \rightarrow H^0(X, \underline{M^\bullet/I^k M^\bullet})$$

Namely, choose any isomorphism

$$\underline{M^\bullet/I^n M^\bullet} \rightarrow K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n$$

induces similar isomorphisms module I^m and I^k and we see that the assertion is true. Thus to prove the lemma it suffices to show that the system $H^0(X, \underline{M^\bullet/I^n M^\bullet})$ has Mittag-Leffler. Since taking sections over X is exact, it suffices to prove that the system of Λ -modules

$$H^0(M^\bullet/I^n M^\bullet)$$

has Mittag-Leffler. Set $A = \Lambda^\wedge$ and consider the spectral sequence

$$\text{Tor}_{-p}^A(H^q(M^\bullet), A/I^n A) \Rightarrow H^{p+q}(M^\bullet/I^n M^\bullet)$$

⁶This may be nonstandard notation

⁷This may be nonstandard notation.

By More on Algebra, Lemma 25.3 the pro-systems $\{\mathrm{Tor}_{-p}^A(H^q(M^\bullet), A/I^n A)\}$ are zero for $p > 0$. Thus the pro-system $\{H^0(M^\bullet/I^n M^\bullet)\}$ is equal to the pro-system $\{H^0(M^\bullet)/I^n H^0(M^\bullet)\}$ and the lemma is proved. \square

09C6 **Lemma 25.6.** *Let X be a connected scheme. Let Λ be a Noetherian ring and let $I \subset \Lambda$ be an ideal. If K is in $D_{\mathrm{cons}}(X, \Lambda)$ such that $K \otimes_{\Lambda} \Lambda/I$ has locally constant cohomology sheaves, then K is adic lisse (Definition 25.4).*

Proof. Write $K_n = K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n$. We will use the results of Lemma 25.2 without further mention. By Cohomology on Sites, Lemma 43.5 we see that K_n has locally constant cohomology sheaves for all n . We have $K_n = \epsilon^{-1} L_n$ some L_n in $D_{\mathrm{ctf}}(X_{\mathrm{\acute{e}tate}}, \Lambda/I^n)$ with locally constant cohomology sheaves. By Étale Cohomology, Lemma 96.14 there exist perfect $M_n \in D(\Lambda/I^n)$ such that L_n is étale locally isomorphic to \underline{M}_n . The maps $L_{n+1} \rightarrow L_n$ corresponding to $K_{n+1} \rightarrow K_n$ induces isomorphisms $\underline{L}_{n+1} \otimes_{\Lambda/I^{n+1}}^{\mathbf{L}} \Lambda/I^n \rightarrow L_n$. Looking locally on X we conclude that there exist maps $M_{n+1} \rightarrow M_n$ in $D(\Lambda/I^{n+1})$ inducing isomorphisms $M_{n+1} \otimes_{\Lambda/I^{n+1}} \Lambda/I^n \rightarrow M_n$, see Cohomology on Sites, Lemma 43.3. Fix a choice of such maps. By More on Algebra, Lemma 83.4 we can find a finite complex M^\bullet of finite projective Λ^\wedge -modules and isomorphisms $M^\bullet/I^n M^\bullet \rightarrow M_n$ in $D(\Lambda/I^n)$ compatible with the transition maps. To finish the proof we will show that K is locally isomorphic to

$$\underline{M}^{\wedge} = \lim \underline{M^\bullet/I^n M^\bullet} = R \lim \underline{M^\bullet/I^n M^\bullet}$$

Let E^\bullet be the dual complex to M^\bullet , see More on Algebra, Lemma 67.14 and its proof. Consider the objects

$$H_n = R \mathcal{H}om_{\Lambda/I^n}(M^\bullet/I^n M^\bullet, K_n) = \underline{E^\bullet/I^n E^\bullet} \otimes_{\Lambda/I^n}^{\mathbf{L}} K_n$$

of $D(X_{\mathrm{pro-étale}}, \Lambda/I^n)$. Modding out by I^n defines a transition map $H_{n+1} \rightarrow H_n$. Set $H = R \lim H_n$. Then H is an object of $D_{\mathrm{cons}}(X, \Lambda)$ (details omitted) with $H \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n = H_n$. Choose a covering $\{W_t \rightarrow X\}_{t \in T}$ with each W_t affine and weakly contractible. By our choice of M^\bullet we see that

$$\begin{aligned} H_n|_{W_t} &\cong R \mathcal{H}om_{\Lambda/I^n}(M^\bullet/I^n M^\bullet, M^\bullet/I^n M^\bullet) \\ &= \underline{\mathrm{Tot}(E^\bullet/I^n E^\bullet \otimes_{\Lambda/I^n} M^\bullet/I^n M^\bullet)} \end{aligned}$$

Thus we may apply Lemma 25.5 to $H = R \lim H_n$. We conclude the system $H^0(W_t, H_n)$ satisfies Mittag-Leffler. Since for all $n \gg 1$ there is an element of $H^0(W_t, H_n)$ which maps to an isomorphism in

$$H^0(W_t, H_1) = \mathrm{Hom}(\underline{M^\bullet/IM^\bullet}, K_1)$$

we find an element $(\varphi_{t,n})$ in the inverse limit which produces an isomorphism mod I . Then

$$R \lim \varphi_{t,n} : \underline{M}^{\wedge}|_{W_t} = R \lim \underline{M^\bullet/I^n M^\bullet}|_{W_t} \longrightarrow R \lim K_n|_{W_t} = K|_{W_t}$$

is an isomorphism. This finishes the proof. \square

09C7 **Proposition 25.7.** *Let X be a Noetherian scheme. Let Λ be a Noetherian ring and let $I \subset \Lambda$ be an ideal. Let K be an object of $D_{\mathrm{cons}}(X, \Lambda)$. Then K is adic constructible (Definition 25.4).*

Proof. This is a consequence of Lemma 25.6 and the fact that a Noetherian scheme is locally connected (Topology, Lemma 9.6), and the definitions. \square

26. Proper base change

09C8 In this section we explain how to prove the proper base change theorem for derived complete objects on the pro-étale site using the proper base change theorem for étale cohomology following the general theme that we use the pro-étale topology only to deal with “limit issues” and we use results proved for the étale topology to handle everything else.

09C9 **Theorem 26.1.** *Let $f : X \rightarrow Y$ be a proper morphism of schemes. Let $g : Y' \rightarrow Y$ be a morphism of schemes giving rise to the base change diagram*

$$\begin{array}{ccc} X' & \xrightarrow{g'} & X \\ f' \downarrow & & \downarrow f \\ Y' & \xrightarrow{g} & Y \end{array}$$

Let Λ be a Noetherian ring and let $I \subset \Lambda$ be an ideal such that Λ/I is torsion. Let K be an object of $D(X_{\text{pro-étale}})$ such that

- (1) K is derived complete, and
- (2) $K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n$ is bounded below with cohomology sheaves coming from $X_{\text{étale}}$,
- (3) Λ/I^n is a perfect Λ -module⁸.

Then the base change map

$$Lg_{\text{comp}}^* Rf_* K \longrightarrow Rf'_* L(g')^*_{\text{comp}} K$$

is an isomorphism.

Proof. We omit the construction of the base change map (this uses only formal properties of derived pushforward and completed derived pullback, compare with Cohomology on Sites, Remark 20.3). Write $K_n = K \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n$. By Lemma 14.1 we have $K = R\lim K_n$ because K is derived complete. By Lemmas 14.2 and 14.1 we can unwind the left hand side

$$Lg_{\text{comp}}^* Rf_* K = R\lim Lg^*(Rf_* K) \otimes_{\Lambda}^{\mathbf{L}} \Lambda/I^n = R\lim Lg^* Rf_* K_n$$

the last equality because Λ/I^n is a perfect module and the projection formula (Cohomology on Sites, Lemma 40.1). Using Lemma 14.2 we can unwind the right hand side

$$Rf'_* L(g')^*_{\text{comp}} K = Rf'_* R\lim L(g')^* K_n = R\lim Rf'_* L(g')^* K_n$$

the last equality because Rf'_* commutes with $R\lim$ (Cohomology on Sites, Lemma 22.3). Thus it suffices to show the maps

$$Lg^* Rf_* K_n \longrightarrow Rf'_* L(g')^* K_n$$

are isomorphisms. By Lemma 16.7 and our second condition we can write $K_n = \epsilon^{-1} L_n$ for some $L_n \in D^+(X_{\text{étale}}, \Lambda/I^n)$. By Lemma 19.1 and the fact that ϵ^{-1} commutes with pullbacks we obtain

$$Lg^* Rf_* K_n = Lg^* Rf_* \epsilon^* L_n = Lg^* \epsilon^{-1} Rf_* L_n = \epsilon^{-1} Lg^* Rf_* L_n$$

and

$$Rf'_* L(g')^* K_n = Rf'_* L(g')^* \epsilon^{-1} L_n = Rf'_* \epsilon^{-1} L(g')^* L_n = \epsilon^{-1} Rf'_* L(g')^* L_n$$

⁸This assumption can be removed if K is a constructible complex, see [BS13].

(this also uses that L_n is bounded below). Finally, by the proper base change theorem for étale cohomology (Étale Cohomology, Theorem 79.11) we have

$$Lg^*Rf_*L_n = Rf'_*L(g')^*L_n$$

(again using that L_n is bounded below) and the theorem is proved. □

27. Other chapters

<p>Preliminaries</p> <ul style="list-style-type: none"> (1) Introduction (2) Conventions (3) Set Theory (4) Categories (5) Topology (6) Sheaves on Spaces (7) Sites and Sheaves (8) Stacks (9) Fields (10) Commutative Algebra (11) Brauer Groups (12) Homological Algebra (13) Derived Categories (14) Simplicial Methods (15) More on Algebra (16) Smoothing Ring Maps (17) Sheaves of Modules (18) Modules on Sites (19) Injectives (20) Cohomology of Sheaves (21) Cohomology on Sites (22) Differential Graded Algebra (23) Divided Power Algebra (24) Hypercoverings <p>Schemes</p> <ul style="list-style-type: none"> (25) Schemes (26) Constructions of Schemes (27) Properties of Schemes (28) Morphisms of Schemes (29) Cohomology of Schemes (30) Divisors (31) Limits of Schemes (32) Varieties (33) Topologies on Schemes (34) Descent (35) Derived Categories of Schemes (36) More on Morphisms (37) More on Flatness (38) Groupoid Schemes 	<ul style="list-style-type: none"> (39) More on Groupoid Schemes (40) Étale Morphisms of Schemes <p>Topics in Scheme Theory</p> <ul style="list-style-type: none"> (41) Chow Homology (42) Intersection Theory (43) Picard Schemes of Curves (44) Adequate Modules (45) Dualizing Complexes (46) Duality for Schemes (47) Discriminants and Differents (48) Local Cohomology (49) Algebraic Curves (50) Resolution of Surfaces (51) Semistable Reduction (52) Fundamental Groups of Schemes (53) Étale Cohomology (54) Crystalline Cohomology (55) Pro-étale Cohomology <p>Algebraic Spaces</p> <ul style="list-style-type: none"> (56) Algebraic Spaces (57) Properties of Algebraic Spaces (58) Morphisms of Algebraic Spaces (59) Decent Algebraic Spaces (60) Cohomology of Algebraic Spaces (61) Limits of Algebraic Spaces (62) Divisors on Algebraic Spaces (63) Algebraic Spaces over Fields (64) Topologies on Algebraic Spaces (65) Descent and Algebraic Spaces (66) Derived Categories of Spaces (67) More on Morphisms of Spaces (68) Flatness on Algebraic Spaces (69) Groupoids in Algebraic Spaces (70) More on Groupoids in Spaces (71) Bootstrap (72) Pushouts of Algebraic Spaces <p>Topics in Geometry</p> <ul style="list-style-type: none"> (73) Quotients of Groupoids (74) More on Cohomology of Spaces (75) Simplicial Spaces
---	---

- | | |
|---------------------------------------|--------------------------------------|
| (76) Duality for Spaces | (93) Cohomology of Algebraic Stacks |
| (77) Formal Algebraic Spaces | (94) Derived Categories of Stacks |
| (78) Restricted Power Series | (95) Introducing Algebraic Stacks |
| (79) Resolution of Surfaces Revisited | (96) More on Morphisms of Stacks |
| Deformation Theory | (97) The Geometry of Stacks |
| (80) Formal Deformation Theory | Topics in Moduli Theory |
| (81) Deformation Theory | (98) Moduli Stacks |
| (82) The Cotangent Complex | (99) Moduli of Curves |
| (83) Deformation Problems | |
| Algebraic Stacks | Miscellany |
| (84) Algebraic Stacks | (100) Examples |
| (85) Examples of Stacks | (101) Exercises |
| (86) Sheaves on Algebraic Stacks | (102) Guide to Literature |
| (87) Criteria for Representability | (103) Desirables |
| (88) Artin's Axioms | (104) Coding Style |
| (89) Quot and Hilbert Spaces | (105) Obsolete |
| (90) Properties of Algebraic Stacks | (106) GNU Free Documentation License |
| (91) Morphisms of Algebraic Stacks | (107) Auto Generated Index |
| (92) Limits of Algebraic Stacks | |

References

- [BBD82] Alexander A. Beilinson, Joseph Bernstein, and Pierre Deligne, *Faisceaux pervers*, Analysis and topology on singular spaces, I (Luminy, 1981), Astérisque, vol. 100, Soc. Math. France, Paris, 1982, pp. 5–171.
- [BS13] Bhargav Bhatt and Peter Scholze, *The pro-étale topology for schemes*, preprint, 2013.
- [Del74] Pierre Deligne, *La conjecture de Weil. I*, Inst. Hautes Études Sci. Publ. Math. (1974), no. 43, 273–307.
- [Eke90] Torsten Ekedahl, *On the adic formalism*, 197–218.
- [Gro77] Alexandre Grothendieck, *Séminaire de géométrie algébrique du bois-marie 1965-66, cohomologie l-adique et fonctions l*, sga5, Springer Lecture Notes, vol. 589, Springer-Verlag, 1977.
- [Jan88] Uwe Jannsen, *Continuous étale cohomology*, Math. Ann. **280** (1988), 207–245.
- [Tat76] John Tate, *Relations between k_2 and galois cohomology*, Inventiones mathematicae **36** (1976), 257–274.