

DESCENT AND ALGEBRAIC SPACES

03YE

Contents

1. Introduction	1
2. Conventions	1
3. Descent data for quasi-coherent sheaves	2
4. Fpqc descent of quasi-coherent sheaves	3
5. Descent of finiteness properties of modules	4
6. Fpqc coverings	5
7. Descent of finiteness and smoothness properties of morphisms	6
8. Descending properties of spaces	8
9. Descending properties of morphisms	9
10. Descending properties of morphisms in the fpqc topology	11
11. Descending properties of morphisms in the fppf topology	19
12. Application of descent of properties of morphisms	20
13. Properties of morphisms local on the source	21
14. Properties of morphisms local in the fpqc topology on the source	22
15. Properties of morphisms local in the fppf topology on the source	23
16. Properties of morphisms local in the syntomic topology on the source	23
17. Properties of morphisms local in the smooth topology on the source	23
18. Properties of morphisms local in the étale topology on the source	24
19. Properties of morphisms smooth local on source-and-target	24
20. Properties of morphisms étale-smooth local on source-and-target	27
21. Descent data for spaces over spaces	30
22. Descent data in terms of sheaves	34
23. Other chapters	35
References	36

1. Introduction

03YF In the chapter on topologies on algebraic spaces (see Topologies on Spaces, Section 1) we introduced étale, fppf, smooth, syntomic and fpqc coverings of algebraic spaces. In this chapter we discuss what kind of structures over algebraic spaces can be descended through such coverings. See for example [Gro95a], [Gro95b], [Gro95e], [Gro95f], [Gro95c], and [Gro95d].

2. Conventions

041I

The standing assumption is that all schemes are contained in a big fppf site Sch_{fppf} . And all rings A considered have the property that $\text{Spec}(A)$ is (isomorphic) to an object of this big site.

Let S be a scheme and let X be an algebraic space over S . In this chapter and the following we will write $X \times_S X$ for the product of X with itself (in the category of algebraic spaces over S), instead of $X \times X$.

3. Descent data for quasi-coherent sheaves

04W2 This section is the analogue of Descent, Section 2 for algebraic spaces. It makes sense to read that section first.

04W3 **Definition 3.1.** Let S be a scheme. Let $\{f_i : X_i \rightarrow X\}_{i \in I}$ be a family of morphisms of algebraic spaces over S with fixed target X .

- (1) A *descent datum* $(\mathcal{F}_i, \varphi_{ij})$ for quasi-coherent sheaves with respect to the given family is given by a quasi-coherent sheaf \mathcal{F}_i on X_i for each $i \in I$, an isomorphism of quasi-coherent $\mathcal{O}_{X_i \times_X X_j}$ -modules $\varphi_{ij} : \text{pr}_0^* \mathcal{F}_i \rightarrow \text{pr}_1^* \mathcal{F}_j$ for each pair $(i, j) \in I^2$ such that for every triple of indices $(i, j, k) \in I^3$ the diagram

$$\begin{array}{ccc} \text{pr}_0^* \mathcal{F}_i & \xrightarrow{\quad \text{pr}_{02}^* \varphi_{ik} \quad} & \text{pr}_2^* \mathcal{F}_k \\ & \searrow \text{pr}_{01}^* \varphi_{ij} & \nearrow \text{pr}_{12}^* \varphi_{jk} \\ & \text{pr}_1^* \mathcal{F}_j & \end{array}$$

of $\mathcal{O}_{X_i \times_X X_j \times_X X_k}$ -modules commutes. This is called the *cocycle condition*.

- (2) A *morphism* $\psi : (\mathcal{F}_i, \varphi_{ij}) \rightarrow (\mathcal{F}'_i, \varphi'_{ij})$ of descent data is given by a family $\psi = (\psi_i)_{i \in I}$ of morphisms of \mathcal{O}_{X_i} -modules $\psi_i : \mathcal{F}_i \rightarrow \mathcal{F}'_i$ such that all the diagrams

$$\begin{array}{ccc} \text{pr}_0^* \mathcal{F}_i & \xrightarrow{\quad \varphi_{ij} \quad} & \text{pr}_1^* \mathcal{F}_j \\ \text{pr}_0^* \psi_i \downarrow & & \downarrow \text{pr}_1^* \psi_j \\ \text{pr}_0^* \mathcal{F}'_i & \xrightarrow{\quad \varphi'_{ij} \quad} & \text{pr}_1^* \mathcal{F}'_j \end{array}$$

commute.

04W4 **Lemma 3.2.** Let S be a scheme. Let $\mathcal{U} = \{U_i \rightarrow U\}_{i \in I}$ and $\mathcal{V} = \{V_j \rightarrow V\}_{j \in J}$ be families of morphisms of algebraic spaces over S with fixed targets. Let $(g, \alpha : I \rightarrow J, (g_i)) : \mathcal{U} \rightarrow \mathcal{V}$ be a morphism of families of maps with fixed target, see Sites, Definition 8.1. Let $(\mathcal{F}_j, \varphi_{jj'})$ be a descent datum for quasi-coherent sheaves with respect to the family $\{V_j \rightarrow V\}_{j \in J}$. Then

- (1) The system

$$(g_i^* \mathcal{F}_{\alpha(i)}, (g_i \times g_{i'})^* \varphi_{\alpha(i)\alpha(i')})$$

is a descent datum with respect to the family $\{U_i \rightarrow U\}_{i \in I}$.

- (2) This construction is functorial in the descent datum $(\mathcal{F}_j, \varphi_{jj'})$.
(3) Given a second morphism $(g', \alpha' : I \rightarrow J, (g'_i))$ of families of maps with fixed target with $g = g'$ there exists a functorial isomorphism of descent data

$$(g_i^* \mathcal{F}_{\alpha(i)}, (g_i \times g_{i'})^* \varphi_{\alpha(i)\alpha(i')}) \cong ((g'_i)^* \mathcal{F}_{\alpha'(i)}, (g'_i \times g'_{i'})^* \varphi_{\alpha'(i)\alpha'(i')}).$$

Proof. Omitted. Hint: The maps $g_i^* \mathcal{F}_{\alpha(i)} \rightarrow (g'_i)^* \mathcal{F}_{\alpha'(i)}$ which give the isomorphism of descent data in part (3) are the pullbacks of the maps $\varphi_{\alpha(i)\alpha'(i)}$ by the morphisms $(g_i, g'_i) : U_i \rightarrow V_{\alpha(i)} \times_V V_{\alpha'(i)}$. \square

Let $g : U \rightarrow V$ be a morphism of algebraic spaces. The lemma above tells us that there is a well defined pullback functor between the categories of descent data relative to families of maps with target V and U provided there is a morphism between those families of maps which “lives over g ”.

04W5 **Definition 3.3.** Let S be a scheme. Let $\{U_i \rightarrow U\}_{i \in I}$ be a family of morphisms of algebraic spaces over S with fixed target.

- (1) Let \mathcal{F} be a quasi-coherent \mathcal{O}_U -module. We call the unique descent on \mathcal{F} datum with respect to the covering $\{U \rightarrow U\}$ the *trivial descent datum*.
- (2) The pullback of the trivial descent datum to $\{U_i \rightarrow U\}$ is called the *canonical descent datum*. Notation: $(\mathcal{F}|_{U_i}, \text{can})$.
- (3) A descent datum $(\mathcal{F}_i, \varphi_{ij})$ for quasi-coherent sheaves with respect to the given family is said to be *effective* if there exists a quasi-coherent sheaf \mathcal{F} on U such that $(\mathcal{F}_i, \varphi_{ij})$ is isomorphic to $(\mathcal{F}|_{U_i}, \text{can})$.

04W6 **Lemma 3.4.** Let S be a scheme. Let U be an algebraic space over S . Let $\{U_i \rightarrow U\}$ be a Zariski covering of U , see *Topologies on Spaces, Definition 3.1*. Any descent datum on quasi-coherent sheaves for the family $\mathcal{U} = \{U_i \rightarrow U\}$ is effective. Moreover, the functor from the category of quasi-coherent \mathcal{O}_U -modules to the category of descent data with respect to $\{U_i \rightarrow U\}$ is fully faithful.

Proof. Omitted. \square

4. Fpqc descent of quasi-coherent sheaves

04W7 The main application of flat descent for modules is the corresponding descent statement for quasi-coherent sheaves with respect to fpqc-coverings.

04W8 **Proposition 4.1.** Let S be a scheme. Let $\{X_i \rightarrow X\}$ be an fpqc covering of algebraic spaces over S , see *Topologies on Spaces, Definition 9.1*. Any descent datum on quasi-coherent sheaves for $\{X_i \rightarrow X\}$ is effective. Moreover, the functor from the category of quasi-coherent \mathcal{O}_X -modules to the category of descent data with respect to $\{X_i \rightarrow X\}$ is fully faithful.

Proof. This is more or less a formal consequence of the corresponding result for schemes, see Descent, Proposition 5.2. Here is a strategy for a proof:

- (1) The fact that $\{X_i \rightarrow X\}$ is a refinement of the trivial covering $\{X \rightarrow X\}$ gives, via Lemma 3.2, a functor $QCoh(\mathcal{O}_X) \rightarrow DD(\{X_i \rightarrow X\})$ from the category of quasi-coherent \mathcal{O}_X -modules to the category of descent data for the given family.
- (2) In order to prove the proposition we will construct a quasi-inverse functor $\text{back} : DD(\{X_i \rightarrow X\}) \rightarrow QCoh(\mathcal{O}_X)$.
- (3) Applying again Lemma 3.2 we see that there is a functor $DD(\{X_i \rightarrow X\}) \rightarrow DD(\{T_j \rightarrow X\})$ if $\{T_j \rightarrow X\}$ is a refinement of the given family. Hence in order to construct the functor back we may assume that each X_i is a scheme, see *Topologies on Spaces, Lemma 9.5*. This reduces us to the case where all the X_i are schemes.

- (4) A quasi-coherent sheaf on X is by definition a quasi-coherent \mathcal{O}_X -module on $X_{\acute{e}tale}$. Now for any $U \in \text{Ob}(X_{\acute{e}tale})$ we get an fppf covering $\{U_i \times_X X_i \rightarrow U\}$ by schemes and a morphism $g : \{U_i \times_X X_i \rightarrow U\} \rightarrow \{X_i \rightarrow X\}$ of coverings lying over $U \rightarrow X$. Given a descent datum $\xi = (\mathcal{F}_i, \varphi_{ij})$ we obtain a quasi-coherent \mathcal{O}_U -module $\mathcal{F}_{\xi, U}$ corresponding to the pullback $g^*\xi$ of Lemma 3.2 to the covering of U and using effectivity for fppf covering of schemes, see Descent, Proposition 5.2.
- (5) Check that $\xi \mapsto \mathcal{F}_{\xi, U}$ is functorial in ξ . Omitted.
- (6) Check that $\xi \mapsto \mathcal{F}_{\xi, U}$ is compatible with morphisms $U \rightarrow U'$ of the site $X_{\acute{e}tale}$, so that the system of sheaves $\mathcal{F}_{\xi, U}$ corresponds to a quasi-coherent \mathcal{F}_{ξ} on $X_{\acute{e}tale}$, see Properties of Spaces, Lemma 28.3. Details omitted.
- (7) Check that $back : \xi \mapsto \mathcal{F}_{\xi}$ is quasi-inverse to the functor constructed in (1). Omitted.

This finishes the proof. \square

5. Descent of finiteness properties of modules

060T This section is the analogue for the case of algebraic spaces of Descent, Section 7. The goal is to show that one can check a quasi-coherent module has a certain finiteness conditions by checking on the members of a covering. We will repeatedly use the following proof scheme. Suppose that X is an algebraic space, and that $\{X_i \rightarrow X\}$ is a fppf (resp. fpqc) covering. Let $U \rightarrow X$ be a surjective étale morphism such that U is a scheme. Then there exists an fppf (resp. fpqc) covering $\{Y_j \rightarrow X\}$ such that

- (1) $\{Y_j \rightarrow X\}$ is a refinement of $\{X_i \rightarrow X\}$,
- (2) each Y_j is a scheme, and
- (3) each morphism $Y_j \rightarrow X$ factors through U , and
- (4) $\{Y_j \rightarrow U\}$ is an fppf (resp. fpqc) covering of U .

Namely, first refine $\{X_i \rightarrow X\}$ by an fppf (resp. fpqc) covering such that each X_i is a scheme, see Topologies on Spaces, Lemma 7.4, resp. Lemma 9.5. Then set $Y_i = U \times_X X_i$. A quasi-coherent \mathcal{O}_X -module \mathcal{F} is of finite type, of finite presentation, etc if and only if the quasi-coherent \mathcal{O}_U -module $\mathcal{F}|_U$ is of finite type, of finite presentation, etc. Hence we can use the existence of the refinement $\{Y_j \rightarrow X\}$ to reduce the proof of the following lemmas to the case of schemes. We will indicate this by saying that “*the result follows from the case of schemes by étale localization*”.

060U **Lemma 5.1.** *Let X be an algebraic space over a scheme S . Let \mathcal{F} be a quasi-coherent \mathcal{O}_X -module. Let $\{f_i : X_i \rightarrow X\}_{i \in I}$ be an fpqc covering such that each $f_i^*\mathcal{F}$ is a finite type \mathcal{O}_{X_i} -module. Then \mathcal{F} is a finite type \mathcal{O}_X -module.*

Proof. This follows from the case of schemes, see Descent, Lemma 7.1, by étale localization. \square

060V **Lemma 5.2.** *Let X be an algebraic space over a scheme S . Let \mathcal{F} be a quasi-coherent \mathcal{O}_X -module. Let $\{f_i : X_i \rightarrow X\}_{i \in I}$ be an fpqc covering such that each $f_i^*\mathcal{F}$ is an \mathcal{O}_{X_i} -module of finite presentation. Then \mathcal{F} is an \mathcal{O}_X -module of finite presentation.*

Proof. This follows from the case of schemes, see Descent, Lemma 7.3, by étale localization. \square

060W **Lemma 5.3.** *Let X be an algebraic space over a scheme S . Let \mathcal{F} be a quasi-coherent \mathcal{O}_X -module. Let $\{f_i : X_i \rightarrow X\}_{i \in I}$ be an fpqc covering such that each $f_i^* \mathcal{F}$ is a flat \mathcal{O}_{X_i} -module. Then \mathcal{F} is a flat \mathcal{O}_X -module.*

Proof. This follows from the case of schemes, see Descent, Lemma 7.5, by étale localization. \square

060X **Lemma 5.4.** *Let X be an algebraic space over a scheme S . Let \mathcal{F} be a quasi-coherent \mathcal{O}_X -module. Let $\{f_i : X_i \rightarrow X\}_{i \in I}$ be an fpqc covering such that each $f_i^* \mathcal{F}$ is a finite locally free \mathcal{O}_{X_i} -module. Then \mathcal{F} is a finite locally free \mathcal{O}_X -module.*

Proof. This follows from the case of schemes, see Descent, Lemma 7.6, by étale localization. \square

The definition of a locally projective quasi-coherent sheaf can be found in Properties of Spaces, Section 30. It is also proved there that this notion is preserved under pullback.

060Y **Lemma 5.5.** *Let X be an algebraic space over a scheme S . Let \mathcal{F} be a quasi-coherent \mathcal{O}_X -module. Let $\{f_i : X_i \rightarrow X\}_{i \in I}$ be an fpqc covering such that each $f_i^* \mathcal{F}$ is a locally projective \mathcal{O}_{X_i} -module. Then \mathcal{F} is a locally projective \mathcal{O}_X -module.*

Proof. This follows from the case of schemes, see Descent, Lemma 7.7, by étale localization. \square

We also add here two results which are related to the results above, but are of a slightly different nature.

060Z **Lemma 5.6.** *Let S be a scheme. Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . Let \mathcal{F} be a quasi-coherent \mathcal{O}_X -module. Assume f is a finite morphism. Then \mathcal{F} is an \mathcal{O}_X -module of finite type if and only if $f_* \mathcal{F}$ is an \mathcal{O}_Y -module of finite type.*

Proof. As f is finite it is representable. Choose a scheme V and a surjective étale morphism $V \rightarrow Y$. Then $U = V \times_Y X$ is a scheme with a surjective étale morphism towards X and a finite morphism $\psi : U \rightarrow V$ (the base change of f). Since $\psi_*(\mathcal{F}|_U) = f_* \mathcal{F}|_V$ the result of the lemma follows immediately from the schemes version which is Descent, Lemma 7.9. \square

0610 **Lemma 5.7.** *Let S be a scheme. Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . Let \mathcal{F} be a quasi-coherent \mathcal{O}_X -module. Assume f is finite and of finite presentation. Then \mathcal{F} is an \mathcal{O}_X -module of finite presentation if and only if $f_* \mathcal{F}$ is an \mathcal{O}_Y -module of finite presentation.*

Proof. As f is finite it is representable. Choose a scheme V and a surjective étale morphism $V \rightarrow Y$. Then $U = V \times_Y X$ is a scheme with a surjective étale morphism towards X and a finite morphism $\psi : U \rightarrow V$ (the base change of f). Since $\psi_*(\mathcal{F}|_U) = f_* \mathcal{F}|_V$ the result of the lemma follows immediately from the schemes version which is Descent, Lemma 7.10. \square

6. Fpqc coverings

04P0 This section is the analogue of Descent, Section 10. At the moment we do not know if all of the material for fpqc coverings of schemes holds also for algebraic spaces.

04P1 **Lemma 6.1.** *Let S be a scheme. Let $\{f_i : T_i \rightarrow T\}_{i \in I}$ be an fpqc covering of algebraic spaces over S . Suppose that for each i we have an open subspace $W_i \subset T_i$ such that for all $i, j \in I$ we have $pr_0^{-1}(W_i) = pr_1^{-1}(W_j)$ as open subspaces of $T_i \times_T T_j$. Then there exists a unique open subspace $W \subset T$ such that $W_i = f_i^{-1}(W)$ for each i .*

Proof. By Topologies on Spaces, Lemma 9.5 we may assume each T_i is a scheme. Choose a scheme U and a surjective étale morphism $U \rightarrow T$. Then $\{T_i \times_T U \rightarrow U\}$ is an fpqc covering of U and $T_i \times_T U$ is a scheme for each i . Hence we see that the collection of opens $W_i \times_T U$ comes from a unique open subscheme $W' \subset U$ by Descent, Lemma 10.2. As $U \rightarrow T$ is open we can define $W \subset T$ the Zariski open which is the image of W' , see Properties of Spaces, Section 4. We omit the verification that this works, i.e., that W_i is the inverse image of W for each i . \square

04P2 **Lemma 6.2.** *Let S be a scheme. Let $\{T_i \rightarrow T\}$ be an fpqc covering of algebraic spaces over S , see Topologies on Spaces, Definition 9.1. Then given an algebraic space B over S the sequence*

$$\text{Mor}_S(T, B) \longrightarrow \prod_i \text{Mor}_S(T_i, B) \rightrightarrows \prod_{i,j} \text{Mor}_S(T_i \times_T T_j, B)$$

is an equalizer diagram. In other words, every representable functor on the category of algebraic spaces over S satisfies the sheaf condition for fpqc coverings.

Proof. We know this is true if $\{T_i \rightarrow T\}$ is an fpqc covering of schemes, see Properties of Spaces, Proposition 16.1. This is the key fact and we encourage the reader to skip the rest of the proof which is formal. Choose a scheme U and a surjective étale morphism $U \rightarrow T$. Let U_i be a scheme and let $U_i \rightarrow T_i \times_T U$ be a surjective étale morphism. Then $\{U_i \rightarrow U\}$ is an fpqc covering. This follows from Topologies on Spaces, Lemmas 9.3 and 9.4. By the above we have the result for $\{U_i \rightarrow U\}$.

What this means is the following: Suppose that $b_i : T_i \rightarrow B$ is a family of morphisms with $b_i \circ pr_0 = b_j \circ pr_1$ as morphisms $T_i \times_T T_j \rightarrow B$. Then we let $a_i : U_i \rightarrow B$ be the composition of $U_i \rightarrow T_i \times_T U$ with b_i . By what was said above we find a unique morphism $a : U \rightarrow B$ such that a_i is the composition of a with $U_i \rightarrow U$. The uniqueness guarantees that $a \circ pr_0 = a \circ pr_1$ as morphisms $U \times_T U \rightarrow B$. Then since $T = U/(U \times_T U)$ as a sheaf, we find that a comes from a unique morphism $b : T \rightarrow B$. Chasing diagrams we find that b is the morphism we are looking for. \square

7. Descent of finiteness and smoothness properties of morphisms

06NQ The following type of lemma is occasionally useful.

06NR **Lemma 7.1.** *Let S be a scheme. Let $X \rightarrow Y \rightarrow Z$ be morphism of algebraic spaces. Let P be one of the following properties of morphisms of algebraic spaces over S : flat, locally finite type, locally finite presentation. Assume that $X \rightarrow Z$ has P and that $X \rightarrow Y$ is a surjection of sheaves on $(\text{Sch}/S)_{\text{fppf}}$. Then $Y \rightarrow Z$ is P .*

Proof. Choose a scheme W and a surjective étale morphism $W \rightarrow Z$. Choose a scheme V and a surjective étale morphism $V \rightarrow W \times_Z Y$. Choose a scheme U and a surjective étale morphism $U \rightarrow V \times_Y X$. By assumption we can find an fppf covering $\{V_i \rightarrow V\}$ and lifts $V_i \rightarrow X$ of the morphism $V_i \rightarrow Y$. Since $U \rightarrow X$ is surjective étale we see that over the members of the fppf covering $\{V_i \times_X U \rightarrow V\}$

we have lifts into U . Hence $U \rightarrow V$ induces a surjection of sheaves on $(Sch/S)_{fppf}$. By our definition of what it means to have property P for a morphism of algebraic spaces (see Morphisms of Spaces, Definition 29.1, Definition 23.1, and Definition 28.1) we see that $U \rightarrow W$ has P and we have to show $V \rightarrow W$ has P . Thus we reduce the question to the case of morphisms of schemes which is treated in Descent, Lemma 11.8. \square

A more standard case of the above lemma is the following. (The version with “flat” follows from Morphisms of Spaces, Lemma 30.5.)

0AHC **Lemma 7.2.** *Let S be a scheme. Let*

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow p & \swarrow q \\ & & B \end{array}$$

be a commutative diagram of morphisms of algebraic spaces over S . Assume that f is surjective, flat, and locally of finite presentation and assume that p is locally of finite presentation (resp. locally of finite type). Then q is locally of finite presentation (resp. locally of finite type).

Proof. Since $\{X \rightarrow Y\}$ is an fppf covering, it induces a surjection of fppf sheaves (Topologies on Spaces, Lemma 7.5) and the lemma is a special case of Lemma 7.1. On the other hand, an easier argument is to deduce it from the analogue for schemes. Namely, the problem is étale local on B and Y (Morphisms of Spaces, Lemmas 23.4 and 28.4). Hence we may assume that B and Y are affine schemes. Since $|X| \rightarrow |Y|$ is open (Morphisms of Spaces, Lemma 29.6), we can choose an affine scheme U and an étale morphism $U \rightarrow X$ such that the composition $U \rightarrow Y$ is surjective. In this case the result follows from Descent, Lemma 11.3. \square

0AHD **Lemma 7.3.** *Let S be a scheme. Let*

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow p & \swarrow q \\ & & B \end{array}$$

be a commutative diagram of morphisms of algebraic spaces over S . Assume that

- (1) *f is surjective, and syntomic (resp. smooth, resp. étale),*
- (2) *p is syntomic (resp. smooth, resp. étale).*

Then q is syntomic (resp. smooth, resp. étale).

Proof. We deduce this from the analogue for schemes. Namely, the problem is étale local on B and Y (Morphisms of Spaces, Lemmas 35.4, 36.4, and 38.2). Hence we may assume that B and Y are affine schemes. Since $|X| \rightarrow |Y|$ is open (Morphisms of Spaces, Lemma 29.6), we can choose an affine scheme U and an étale morphism $U \rightarrow X$ such that the composition $U \rightarrow Y$ is surjective. In this case the result follows from Descent, Lemma 11.4. \square

Actually we can strengthen this result as follows.

0AHE **Lemma 7.4.** *Let S be a scheme. Let*

$$\begin{array}{ccc} X & \xrightarrow{\quad} & Y \\ & \searrow p & \swarrow q \\ & & B \end{array}$$

be a commutative diagram of morphisms of algebraic spaces over S . Assume that

- (1) *f is surjective, flat, and locally of finite presentation,*
- (2) *p is smooth (resp. étale).*

Then q is smooth (resp. étale).

Proof. We deduce this from the analogue for schemes. Namely, the problem is étale local on B and Y (Morphisms of Spaces, Lemmas 36.4 and 38.2). Hence we may assume that B and Y are affine schemes. Since $|X| \rightarrow |Y|$ is open (Morphisms of Spaces, Lemma 29.6), we can choose an affine scheme U and an étale morphism $U \rightarrow X$ such that the composition $U \rightarrow Y$ is surjective. In this case the result follows from Descent, Lemma 11.5. \square

0AHF **Lemma 7.5.** *Let S be a scheme. Let*

$$\begin{array}{ccc} X & \xrightarrow{\quad} & Y \\ & \searrow p & \swarrow q \\ & & B \end{array}$$

be a commutative diagram of morphisms of algebraic spaces over S . Assume that

- (1) *f is surjective, flat, and locally of finite presentation,*
- (2) *p is syntomic.*

Then both q and f are syntomic.

Proof. We deduce this from the analogue for schemes. Namely, the problem is étale local on B and Y (Morphisms of Spaces, Lemma 35.4). Hence we may assume that B and Y are affine schemes. Since $|X| \rightarrow |Y|$ is open (Morphisms of Spaces, Lemma 29.6), we can choose an affine scheme U and an étale morphism $U \rightarrow X$ such that the composition $U \rightarrow Y$ is surjective. In this case the result follows from Descent, Lemma 11.7. \square

8. Descending properties of spaces

06DP In this section we put some results of the following kind.

06DQ **Lemma 8.1.** *Let S be a scheme. Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . Let $x \in |X|$. If f is flat at x and X is geometrically unibranch at x , then Y is geometrically unibranch at $f(x)$.*

Proof. Consider the map of étale local rings $\mathcal{O}_{Y,f(\bar{x})} \rightarrow \mathcal{O}_{X,\bar{x}}$. By Morphisms of Spaces, Lemma 29.8 this is flat. Hence if $\mathcal{O}_{X,\bar{x}}$ has a unique minimal prime, so does $\mathcal{O}_{Y,f(\bar{x})}$ (by going down, see Algebra, Lemma 38.18). \square

06MI **Lemma 8.2.** *Let S be a scheme. Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . If f is flat and surjective and X is reduced, then Y is reduced.*

Proof. Choose a scheme V and a surjective étale morphism $V \rightarrow Y$. Choose a scheme U and a surjective étale morphism $U \rightarrow X \times_Y V$. As f is surjective and flat, the morphism of schemes $U \rightarrow V$ is surjective and flat. In this way we reduce the problem to the case of schemes (as reducedness of X and Y is defined in terms of reducedness of U and V , see Properties of Spaces, Section 7). The case of schemes is Descent, Lemma 16.1. \square

06MJ **Lemma 8.3.** *Let $f : X \rightarrow Y$ be a morphism of algebraic spaces. If f is locally of finite presentation, flat, and surjective and X is locally Noetherian, then Y is locally Noetherian.*

Proof. Choose a scheme V and a surjective étale morphism $V \rightarrow Y$. Choose a scheme U and a surjective étale morphism $U \rightarrow X \times_Y V$. As f is surjective, flat, and locally of finite presentation the morphism of schemes $U \rightarrow V$ is surjective, flat, and locally of finite presentation. In this way we reduce the problem to the case of schemes (as being locally Noetherian for X and Y is defined in terms of being locally Noetherian of U and V , see Properties of Spaces, Section 7). In the case of schemes the result follows from Descent, Lemma 13.1. \square

06MK **Lemma 8.4.** *Let $f : X \rightarrow Y$ be a morphism of algebraic spaces. If f is locally of finite presentation, flat, and surjective and X is regular, then Y is regular.*

Proof. By Lemma 8.3 we know that Y is locally Noetherian. Choose a scheme V and a surjective étale morphism $V \rightarrow Y$. It suffices to prove that the local rings of V are all regular local rings, see Properties, Lemma 9.2. Choose a scheme U and a surjective étale morphism $U \rightarrow X \times_Y V$. As f is surjective and flat the morphism of schemes $U \rightarrow V$ is surjective and flat. By assumption U is a regular scheme in particular all of its local rings are regular (by the lemma above). Hence the lemma follows from Algebra, Lemma 109.9. \square

9. Descending properties of morphisms

03YG In this section we introduce the notion of when a property of morphisms of algebraic spaces is local on the target in a topology. Please compare with Descent, Section 19.

03YH **Definition 9.1.** Let S be a scheme. Let \mathcal{P} be a property of morphisms of algebraic spaces over S . Let $\tau \in \{fpqc, fppf, syntomic, smooth, \acute{e}tale\}$. We say \mathcal{P} is τ local on the base, or τ local on the target, or local on the base for the τ -topology if for any τ -covering $\{Y_i \rightarrow Y\}_{i \in I}$ of algebraic spaces and any morphism of algebraic spaces $f : X \rightarrow Y$ we have

$$f \text{ has } \mathcal{P} \Leftrightarrow \text{each } Y_i \times_Y X \rightarrow Y_i \text{ has } \mathcal{P}.$$

To be sure, since isomorphisms are always coverings we see (or require) that property \mathcal{P} holds for $X \rightarrow Y$ if and only if it holds for any arrow $X' \rightarrow Y'$ isomorphic to $X \rightarrow Y$. If a property is τ -local on the target then it is preserved by base changes by morphisms which occur in τ -coverings. Here is a formal statement.

06EM **Lemma 9.2.** *Let S be a scheme. Let $\tau \in \{fpqc, fppf, syntomic, smooth, \acute{e}tale\}$. Let \mathcal{P} be a property of morphisms of algebraic spaces over S which is τ local on the target. Let $f : X \rightarrow Y$ have property \mathcal{P} . For any morphism $Y' \rightarrow Y$ which is flat, resp. flat and locally of finite presentation, resp. syntomic, resp. étale, the base change $f' : Y' \times_Y X \rightarrow Y'$ of f has property \mathcal{P} .*

Proof. This is true because we can fit $Y' \rightarrow Y$ into a family of morphisms which forms a τ -covering. \square

A simple often used consequence of the above is that if $f : X \rightarrow Y$ has property \mathcal{P} which is τ -local on the target and $f(X) \subset V$ for some open subspace $V \subset Y$, then also the induced morphism $X \rightarrow V$ has \mathcal{P} . Proof: The base change f by $V \rightarrow Y$ gives $X \rightarrow V$.

06R2 **Lemma 9.3.** *Let S be a scheme. Let $\tau \in \{fppf, \text{syntomic}, \text{smooth}, \text{étale}\}$. Let \mathcal{P} be a property of morphisms of algebraic spaces over S which is τ local on the target. For any morphism of algebraic spaces $f : X \rightarrow Y$ over S there exists a largest open subspace $W(f) \subset Y$ such that the restriction $X_{W(f)} \rightarrow W(f)$ has \mathcal{P} . Moreover,*

- (1) *if $g : Y' \rightarrow Y$ is a morphism of algebraic spaces which is flat and locally of finite presentation, syntomic, smooth, or étale and the base change $f' : X_{Y'} \rightarrow Y'$ has \mathcal{P} , then g factors through $W(f)$,*
- (2) *if $g : Y' \rightarrow Y$ is flat and locally of finite presentation, syntomic, smooth, or étale, then $W(f') = g^{-1}(W(f))$, and*
- (3) *if $\{g_i : Y_i \rightarrow Y\}$ is a τ -covering, then $g_i^{-1}(W(f)) = W(f_i)$, where f_i is the base change of f by $Y_i \rightarrow Y$.*

Proof. Consider the union $W_{set} \subset |Y|$ of the images $g(|Y'|) \subset |Y|$ of morphisms $g : Y' \rightarrow Y$ with the properties:

- (1) g is flat and locally of finite presentation, syntomic, smooth, or étale, and
- (2) the base change $Y' \times_{g,Y} X \rightarrow Y'$ has property \mathcal{P} .

Since such a morphism g is open (see Morphisms of Spaces, Lemma 29.6) we see that W_{set} is an open subset of $|Y|$. Denote $W \subset Y$ the open subspace whose underlying set of points is W_{set} , see Properties of Spaces, Lemma 4.8. Since \mathcal{P} is local in the τ topology the restriction $X_W \rightarrow W$ has property \mathcal{P} because we are given a covering $\{Y' \rightarrow W\}$ of W such that the pullbacks have \mathcal{P} . This proves the existence and proves that $W(f)$ has property (1). To see property (2) note that $W(f') \supset g^{-1}(W(f))$ because \mathcal{P} is stable under base change by flat and locally of finite presentation, syntomic, smooth, or étale morphisms, see Lemma 9.2. On the other hand, if $Y'' \subset Y'$ is an open such that $X_{Y''} \rightarrow Y''$ has property \mathcal{P} , then $Y'' \rightarrow Y$ factors through W by construction, i.e., $Y'' \subset g^{-1}(W(f))$. This proves (2). Assertion (3) follows from (2) because each morphism $Y_i \rightarrow Y$ is flat and locally of finite presentation, syntomic, smooth, or étale by our definition of a τ -covering. \square

041J **Lemma 9.4.** *Let S be a scheme. Let \mathcal{P} be a property of morphisms of algebraic spaces over S . Assume*

- (1) *if $X_i \rightarrow Y_i$, $i = 1, 2$ have property \mathcal{P} so does $X_1 \amalg X_2 \rightarrow Y_1 \amalg Y_2$,*
- (2) *a morphism of algebraic spaces $f : X \rightarrow Y$ has property \mathcal{P} if and only if for every affine scheme Z and morphism $Z \rightarrow Y$ the base change $Z \times_Y X \rightarrow Z$ of f has property \mathcal{P} , and*
- (3) *for any surjective flat morphism of affine schemes $Z' \rightarrow Z$ over S and a morphism $f : X \rightarrow Z$ from an algebraic space to Z we have*

$$f' : Z' \times_Z X \rightarrow Z' \text{ has } \mathcal{P} \Rightarrow f \text{ has } \mathcal{P}.$$

Then \mathcal{P} is fpqc local on the base.

Proof. If \mathcal{P} has property (2), then it is automatically stable under any base change. Hence the direct implication in Definition 9.1.

Let $\{Y_i \rightarrow Y\}_{i \in I}$ be an fpqc covering of algebraic spaces over S . Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . Assume each base change $f_i : Y_i \times_Y X \rightarrow Y_i$ has property \mathcal{P} . Our goal is to show that f has \mathcal{P} . Let Z be an affine scheme, and let $Z \rightarrow Y$ be a morphism. By (2) it suffices to show that the morphism of algebraic spaces $Z \times_Y X \rightarrow Z$ has \mathcal{P} . Since $\{Y_i \rightarrow Y\}_{i \in I}$ is an fpqc covering we know there exists a standard fpqc covering $\{Z_j \rightarrow Z\}_{j=1, \dots, n}$ and morphisms $Z_j \rightarrow Y_{i_j}$ over Y for suitable indices $i_j \in I$. Since f_{i_j} has \mathcal{P} we see that

$$Z_j \times_Y X = Z_j \times_{Y_{i_j}} (Y_{i_j} \times_Y X) \longrightarrow Z_j$$

has \mathcal{P} as a base change of f_{i_j} (see first remark of the proof). Set $Z' = \coprod_{j=1, \dots, n} Z_j$, so that $Z' \rightarrow Z$ is a flat and surjective morphism of affine schemes over S . By (1) we conclude that $Z' \times_Y X \rightarrow Z'$ has property \mathcal{P} . Since this is the base change of the morphism $Z \times_Y X \rightarrow Z$ by the morphism $Z' \rightarrow Z$ we conclude that $Z \times_Y X \rightarrow Z$ has property \mathcal{P} as desired. \square

10. Descending properties of morphisms in the fpqc topology

041K In this section we find a large number of properties of morphisms of algebraic spaces which are local on the base in the fpqc topology. Please compare with Descent, Section 20 for the case of morphisms of schemes.

041L **Lemma 10.1.** *Let S be a scheme. The property $\mathcal{P}(f) = \text{“}f \text{ is quasi-compact”}$ is fpqc local on the base on algebraic spaces over S .*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 8.7. Let $Z' \rightarrow Z$ be a surjective flat morphism of affine schemes over S . Let $f : X \rightarrow Z$ be a morphism of algebraic spaces, and assume that the base change $f' : Z' \times_Z X \rightarrow Z'$ is quasi-compact. We have to show that f is quasi-compact. To see this, using Morphisms of Spaces, Lemma 8.7 again, it is enough to show that for every affine scheme Y and morphism $Y \rightarrow Z$ the fibre product $Y \times_Z X$ is quasi-compact. Here is a picture:

041M (10.1.1)

$$\begin{array}{ccccc}
 Y \times_Z Z' \times_Z X & \xrightarrow{\quad\quad\quad} & Z' \times_Z X & & \\
 \downarrow & \searrow & \downarrow & \searrow & \\
 & & Y \times_Z X & \xrightarrow{\quad\quad\quad} & X \\
 & & \downarrow & \downarrow f' & \downarrow f \\
 Y \times_Z Z' & \xrightarrow{\quad\quad\quad} & Z' & & \\
 \downarrow & \searrow & \downarrow & \searrow & \\
 & & Y & \xrightarrow{\quad\quad\quad} & Z
 \end{array}$$

Note that all squares are cartesian and the bottom square consists of affine schemes. The assumption that f' is quasi-compact combined with the fact that $Y \times_Z Z'$ is affine implies that $Y \times_Z Z' \times_Z X$ is quasi-compact. Since

$$Y \times_Z Z' \times_Z X \longrightarrow Y \times_Z X$$

is surjective as a base change of $Z' \rightarrow Z$ we conclude that $Y \times_Z X$ is quasi-compact, see Morphisms of Spaces, Lemma 8.5. This finishes the proof. \square

041N **Lemma 10.2.** *Let S be a scheme. The property $\mathcal{P}(f) = “f$ is quasi-separated” is fpqc local on the base on algebraic spaces over S .*

Proof. A base change of a quasi-separated morphism is quasi-separated, see Morphisms of Spaces, Lemma 4.4. Hence the direct implication in Definition 9.1.

Let $\{Y_i \rightarrow Y\}_{i \in I}$ be an fpqc covering of algebraic spaces over S . Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . Assume each base change $X_i := Y_i \times_Y X \rightarrow Y_i$ is quasi-separated. This means that each of the morphisms

$$\Delta_i : X_i \longrightarrow X_i \times_{Y_i} X_i = Y_i \times_Y (X \times_Y X)$$

is quasi-compact. The base change of a fpqc covering is an fpqc covering, see Topologies on Spaces, Lemma 9.3 hence $\{Y_i \times_Y (X \times_Y X) \rightarrow X \times_Y X\}$ is an fpqc covering of algebraic spaces. Moreover, each Δ_i is the base change of the morphism $\Delta : X \rightarrow X \times_Y X$. Hence it follows from Lemma 10.1 that Δ is quasi-compact, i.e., f is quasi-separated. \square

041O **Lemma 10.3.** *Let S be a scheme. The property $\mathcal{P}(f) = “f$ is universally closed” is fpqc local on the base on algebraic spaces over S .*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 9.5. Let $Z' \rightarrow Z$ be a surjective flat morphism of affine schemes over S . Let $f : X \rightarrow Z$ be a morphism of algebraic spaces, and assume that the base change $f' : Z' \times_Z X \rightarrow Z'$ is universally closed. We have to show that f is universally closed. To see this, using Morphisms of Spaces, Lemma 9.5 again, it is enough to show that for every affine scheme Y and morphism $Y \rightarrow Z$ the map $|Y \times_Z X| \rightarrow |Y|$ is closed. Consider the cube (10.1.1). The assumption that f' is universally closed implies that $|Y \times_Z Z' \times_Z X| \rightarrow |Y \times_Z Z'|$ is closed. As $Y \times_Z Z' \rightarrow Y$ is quasi-compact, surjective, and flat as a base change of $Z' \rightarrow Z$ we see the map $|Y \times_Z Z'| \rightarrow |Y|$ is submersive, see Morphisms, Lemma 24.11. Moreover the map

$$|Y \times_Z Z' \times_Z X| \longrightarrow |Y \times_Z Z'| \times_{|Y|} |Y \times_Z X|$$

is surjective, see Properties of Spaces, Lemma 4.3. It follows by elementary topology that $|Y \times_Z X| \rightarrow |Y|$ is closed. \square

041P **Lemma 10.4.** *Let S be a scheme. The property $\mathcal{P}(f) = “f$ is universally open” is fpqc local on the base on algebraic spaces over S .*

Proof. The proof is the same as the proof of Lemma 10.3. \square

0CFW **Lemma 10.5.** *The property $\mathcal{P}(f) = “f$ is universally submersive” is fpqc local on the base.*

Proof. The proof is the same as the proof of Lemma 10.3. \square

041Q **Lemma 10.6.** *The property $\mathcal{P}(f) = “f$ is surjective” is fpqc local on the base.*

Proof. Omitted. (Hint: Use Properties of Spaces, Lemma 4.3.) \square

041R **Lemma 10.7.** *The property $\mathcal{P}(f) = “f$ is universally injective” is fpqc local on the base.*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 9.5. Let $Z' \rightarrow Z$ be a flat surjective morphism of affine schemes over S and let $f : X \rightarrow Z$ be a morphism from an algebraic space to Z . Assume that the base change $f' : X' \rightarrow Z'$ is universally injective. Let K be a field, and let $a, b : \text{Spec}(K) \rightarrow X$ be two morphisms such that $f \circ a = f \circ b$. As $Z' \rightarrow Z$ is surjective there exists a field extension $K \subset K'$ and a morphism $\text{Spec}(K') \rightarrow Z'$ such that the following solid diagram commutes

$$\begin{array}{ccccc}
 \text{Spec}(K') & & & & \\
 \downarrow & \swarrow & & \searrow & \\
 & a', b' & & & \\
 & & X' & \longrightarrow & Z' \\
 & & \downarrow & & \downarrow \\
 \text{Spec}(K) & \xrightarrow{a, b} & X & \longrightarrow & Z
 \end{array}$$

As the square is cartesian we get the two dotted arrows a', b' making the diagram commute. Since $X' \rightarrow Z'$ is universally injective we get $a' = b'$. This forces $a = b$ as $\{\text{Spec}(K') \rightarrow \text{Spec}(K)\}$ is an fpqc covering, see Properties of Spaces, Proposition 16.1. Hence f is universally injective as desired. \square

0CFX **Lemma 10.8.** *The property $\mathcal{P}(f) = "f \text{ is a universal homeomorphism}"$ is fpqc local on the base.*

Proof. This can be proved in exactly the same manner as Lemma 10.3. Alternatively, one can use that a map of topological spaces is a homeomorphism if and only if it is injective, surjective, and open. Thus a universal homeomorphism is the same thing as a surjective, universally injective, and universally open morphism. See Morphisms of Spaces, Lemma 5.5 and Morphisms of Spaces, Definitions 19.3, 5.2, 6.2, 51.2. Thus the lemma follows from Lemmas 10.6, 10.7, and 10.4. \square

041S **Lemma 10.9.** *The property $\mathcal{P}(f) = "f \text{ is locally of finite type}"$ is fpqc local on the base.*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 23.4. Let $Z' \rightarrow Z$ be a surjective flat morphism of affine schemes over S . Let $f : X \rightarrow Z$ be a morphism of algebraic spaces, and assume that the base change $f' : Z' \times_Z X \rightarrow Z'$ is locally of finite type. We have to show that f is locally of finite type. Let U be a scheme and let $U \rightarrow X$ be surjective and étale. By Morphisms of Spaces, Lemma 23.4 again, it is enough to show that $U \rightarrow Z$ is locally of finite type. Since f' is locally of finite type, and since $Z' \times_Z U$ is a scheme étale over $Z' \times_Z X$ we conclude (by the same lemma again) that $Z' \times_Z U \rightarrow Z'$ is locally of finite type. As $\{Z' \rightarrow Z\}$ is an fpqc covering we conclude that $U \rightarrow Z$ is locally of finite type by Descent, Lemma 20.10 as desired. \square

041T **Lemma 10.10.** *The property $\mathcal{P}(f) = "f \text{ is locally of finite presentation}"$ is fpqc local on the base.*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 28.4. Let $Z' \rightarrow Z$ be a surjective flat morphism of affine schemes over S . Let $f : X \rightarrow Z$ be a morphism of algebraic

spaces, and assume that the base change $f' : Z' \times_Z X \rightarrow Z'$ is locally of finite presentation. We have to show that f is locally of finite presentation. Let U be a scheme and let $U \rightarrow X$ be surjective and étale. By Morphisms of Spaces, Lemma 28.4 again, it is enough to show that $U \rightarrow Z$ is locally of finite presentation. Since f' is locally of finite presentation, and since $Z' \times_Z U$ is a scheme étale over $Z' \times_Z X$ we conclude (by the same lemma again) that $Z' \times_Z U \rightarrow Z'$ is locally of finite presentation. As $\{Z' \rightarrow Z\}$ is an fpqc covering we conclude that $U \rightarrow Z$ is locally of finite presentation by Descent, Lemma 20.11 as desired. \square

041U **Lemma 10.11.** *The property $\mathcal{P}(f) = “f \text{ is of finite type}”$ is fpqc local on the base.*

Proof. Combine Lemmas 10.1 and 10.9. \square

041V **Lemma 10.12.** *The property $\mathcal{P}(f) = “f \text{ is of finite presentation}”$ is fpqc local on the base.*

Proof. Combine Lemmas 10.1, 10.2 and 10.10. \square

041W **Lemma 10.13.** *The property $\mathcal{P}(f) = “f \text{ is flat}”$ is fpqc local on the base.*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 29.5. Let $Z' \rightarrow Z$ be a surjective flat morphism of affine schemes over S . Let $f : X \rightarrow Z$ be a morphism of algebraic spaces, and assume that the base change $f' : Z' \times_Z X \rightarrow Z'$ is flat. We have to show that f is flat. Let U be a scheme and let $U \rightarrow X$ be surjective and étale. By Morphisms of Spaces, Lemma 29.5 again, it is enough to show that $U \rightarrow Z$ is flat. Since f' is flat, and since $Z' \times_Z U$ is a scheme étale over $Z' \times_Z X$ we conclude (by the same lemma again) that $Z' \times_Z U \rightarrow Z'$ is flat. As $\{Z' \rightarrow Z\}$ is an fpqc covering we conclude that $U \rightarrow Z$ is flat by Descent, Lemma 20.15 as desired. \square

041X **Lemma 10.14.** *The property $\mathcal{P}(f) = “f \text{ is an open immersion}”$ is fpqc local on the base.*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 12.1. Consider a cartesian diagram

$$\begin{array}{ccc} X' & \longrightarrow & X \\ \downarrow & & \downarrow \\ Z' & \longrightarrow & Z \end{array}$$

of algebraic spaces over S where $Z' \rightarrow Z$ is a surjective flat morphism of affine schemes, and $X' \rightarrow Z'$ is an open immersion. We have to show that $X \rightarrow Z$ is an open immersion. Note that $|X'| \subset |Z'|$ corresponds to an open subscheme $U' \subset Z'$ (isomorphic to X') with the property that $\text{pr}_0^{-1}(U') = \text{pr}_1^{-1}(U')$ as open subschemes of $Z' \times_Z Z'$. Hence there exists an open subscheme $U \subset Z$ such that $X' = (Z' \rightarrow Z)^{-1}(U)$, see Descent, Lemma 10.2. By Properties of Spaces, Proposition 16.1 we see that X satisfies the sheaf condition for the fpqc topology. Now we have the fpqc covering $\mathcal{U} = \{U' \rightarrow U\}$ and the element $U' \rightarrow X' \rightarrow X \in \check{H}^0(\mathcal{U}, X)$. By the

sheaf condition we obtain a morphism $U \rightarrow X$ such that

$$\begin{array}{ccc} U' & \longrightarrow & U \\ \downarrow \cong & & \downarrow \\ X' & \longrightarrow & X \\ \downarrow & & \downarrow \\ Z' & \longrightarrow & Z \end{array}$$

is commutative. On the other hand, we know that for any scheme T over S and T -valued point $T \rightarrow X$ the composition $T \rightarrow X \rightarrow Z$ is a morphism such that $Z' \times_Z T \rightarrow Z'$ factors through U' . Clearly this means that $T \rightarrow Z$ factors through U . In other words the map of sheaves $U \rightarrow X$ is bijective and we win. \square

041Y **Lemma 10.15.** *The property $\mathcal{P}(f) = "f \text{ is an isomorphism}"$ is fpqc local on the base.*

Proof. Combine Lemmas 10.6 and 10.14. \square

041Z **Lemma 10.16.** *The property $\mathcal{P}(f) = "f \text{ is affine}"$ is fpqc local on the base.*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 20.3. Let $Z' \rightarrow Z$ be a surjective flat morphism of affine schemes over S . Let $f : X \rightarrow Z$ be a morphism of algebraic spaces, and assume that the base change $f' : Z' \times_Z X \rightarrow Z'$ is affine. Let X' be a scheme representing $Z' \times_Z X$. We obtain a canonical isomorphism

$$\varphi : X' \times_{Z'} Z' \longrightarrow Z' \times_Z X'$$

since both schemes represent the algebraic space $Z' \times_Z Z' \times_Z X$. This is a descent datum for $X'/Z'/Z$, see Descent, Definition 31.1 (verification omitted, compare with Descent, Lemma 36.1). Since $X' \rightarrow Z'$ is affine this descent datum is effective, see Descent, Lemma 34.1. Thus there exists a scheme $Y \rightarrow Z$ over Z and an isomorphism $\psi : Z' \times_Z Y \rightarrow X'$ compatible with descent data. Of course $Y \rightarrow Z$ is affine (by construction or by Descent, Lemma 20.18). Note that $\mathcal{Y} = \{Z' \times_Z Y \rightarrow Y\}$ is a fpqc covering, and interpreting ψ as an element of $X(Z' \times_Z Y)$ we see that $\psi \in \check{H}^0(\mathcal{Y}, X)$. By the sheaf condition for X with respect to this covering (see Properties of Spaces, Proposition 16.1) we obtain a morphism $Y \rightarrow X$. By construction the base change of this to Z' is an isomorphism, hence an isomorphism by Lemma 10.15. This proves that X is representable by an affine scheme and we win. \square

0420 **Lemma 10.17.** *The property $\mathcal{P}(f) = "f \text{ is a closed immersion}"$ is fpqc local on the base.*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 12.1. Consider a cartesian diagram

$$\begin{array}{ccc} X' & \longrightarrow & X \\ \downarrow & & \downarrow \\ Z' & \longrightarrow & Z \end{array}$$

of algebraic spaces over S where $Z' \rightarrow Z$ is a surjective flat morphism of affine schemes, and $X' \rightarrow Z'$ is a closed immersion. We have to show that $X \rightarrow Z$ is a closed immersion. The morphism $X' \rightarrow Z'$ is affine. Hence by Lemma 10.16 we see that X is a scheme and $X \rightarrow Z$ is affine. It follows from Descent, Lemma 20.19 that $X \rightarrow Z$ is a closed immersion as desired. \square

0421 **Lemma 10.18.** *The property $\mathcal{P}(f) = “f \text{ is separated}”$ is fpqc local on the base.*

Proof. A base change of a separated morphism is separated, see Morphisms of Spaces, Lemma 4.4. Hence the direct implication in Definition 9.1.

Let $\{Y_i \rightarrow Y\}_{i \in I}$ be an fpqc covering of algebraic spaces over S . Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . Assume each base change $X_i := Y_i \times_Y X \rightarrow Y_i$ is separated. This means that each of the morphisms

$$\Delta_i : X_i \longrightarrow X_i \times_{Y_i} X_i = Y_i \times_Y (X \times_Y X)$$

is a closed immersion. The base change of a fpqc covering is an fpqc covering, see Topologies on Spaces, Lemma 9.3 hence $\{Y_i \times_Y (X \times_Y X) \rightarrow X \times_Y X\}$ is an fpqc covering of algebraic spaces. Moreover, each Δ_i is the base change of the morphism $\Delta : X \rightarrow X \times_Y X$. Hence it follows from Lemma 10.17 that Δ is a closed immersion, i.e., f is separated. \square

0422 **Lemma 10.19.** *The property $\mathcal{P}(f) = “f \text{ is proper}”$ is fpqc local on the base.*

Proof. The lemma follows by combining Lemmas 10.3, 10.18 and 10.11. \square

0423 **Lemma 10.20.** *The property $\mathcal{P}(f) = “f \text{ is quasi-affine}”$ is fpqc local on the base.*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 21.3. Let $Z' \rightarrow Z$ be a surjective flat morphism of affine schemes over S . Let $f : X \rightarrow Z$ be a morphism of algebraic spaces, and assume that the base change $f' : Z' \times_Z X \rightarrow Z'$ is quasi-affine. Let X' be a scheme representing $Z' \times_Z X$. We obtain a canonical isomorphism

$$\varphi : X' \times_Z Z' \longrightarrow Z' \times_Z X'$$

since both schemes represent the algebraic space $Z' \times_Z Z' \times_Z X$. This is a descent datum for $X'/Z'/Z$, see Descent, Definition 31.1 (verification omitted, compare with Descent, Lemma 36.1). Since $X' \rightarrow Z'$ is quasi-affine this descent datum is effective, see Descent, Lemma 35.1. Thus there exists a scheme $Y \rightarrow Z$ over Z and an isomorphism $\psi : Z' \times_Z Y \rightarrow X'$ compatible with descent data. Of course $Y \rightarrow Z$ is quasi-affine (by construction or by Descent, Lemma 20.20). Note that $\mathcal{Y} = \{Z' \times_Z Y \rightarrow Y\}$ is a fpqc covering, and interpreting ψ as an element of $X(Z' \times_Z Y)$ we see that $\psi \in \check{H}^0(\mathcal{Y}, X)$. By the sheaf condition for X (see Properties of Spaces, Proposition 16.1) we obtain a morphism $Y \rightarrow X$. By construction the base change of this to Z' is an isomorphism, hence an isomorphism by Lemma 10.15. This proves that X is representable by a quasi-affine scheme and we win. \square

0424 **Lemma 10.21.** *The property $\mathcal{P}(f) = “f \text{ is a quasi-compact immersion}”$ is fpqc local on the base.*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemmas 12.1 and 8.7. Consider a cartesian

diagram

$$\begin{array}{ccc} X' & \longrightarrow & X \\ \downarrow & & \downarrow \\ Z' & \longrightarrow & Z \end{array}$$

of algebraic spaces over S where $Z' \rightarrow Z$ is a surjective flat morphism of affine schemes, and $X' \rightarrow Z'$ is a quasi-compact immersion. We have to show that $X \rightarrow Z$ is a closed immersion. The morphism $X' \rightarrow Z'$ is quasi-affine. Hence by Lemma 10.20 we see that X is a scheme and $X \rightarrow Z$ is quasi-affine. It follows from Descent, Lemma 20.21 that $X \rightarrow Z$ is a quasi-compact immersion as desired. \square

0425 **Lemma 10.22.** *The property $\mathcal{P}(f) = “f \text{ is integral}”$ is fpqc local on the base.*

Proof. An integral morphism is the same thing as an affine, universally closed morphism. See Morphisms of Spaces, Lemma 44.7. Hence the lemma follows on combining Lemmas 10.3 and 10.16. \square

0426 **Lemma 10.23.** *The property $\mathcal{P}(f) = “f \text{ is finite}”$ is fpqc local on the base.*

Proof. A finite morphism is the same thing as an integral, morphism which is locally of finite type. See Morphisms of Spaces, Lemma 44.6. Hence the lemma follows on combining Lemmas 10.9 and 10.22. \square

0427 **Lemma 10.24.** *The properties $\mathcal{P}(f) = “f \text{ is locally quasi-finite}”$ and $\mathcal{P}(f) = “f \text{ is quasi-finite}”$ are fpqc local on the base.*

Proof. We have already seen that “quasi-compact” is fpqc local on the base, see Lemma 10.1. Hence it is enough to prove the lemma for “locally quasi-finite”. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 27.6. Let $Z' \rightarrow Z$ be a surjective flat morphism of affine schemes over S . Let $f : X \rightarrow Z$ be a morphism of algebraic spaces, and assume that the base change $f' : Z' \times_Z X \rightarrow Z'$ is locally quasi-finite. We have to show that f is locally quasi-finite. Let U be a scheme and let $U \rightarrow X$ be surjective and étale. By Morphisms of Spaces, Lemma 27.6 again, it is enough to show that $U \rightarrow Z$ is locally quasi-finite. Since f' is locally quasi-finite, and since $Z' \times_Z U$ is a scheme étale over $Z' \times_Z X$ we conclude (by the same lemma again) that $Z' \times_Z U \rightarrow Z'$ is locally quasi-finite. As $\{Z' \rightarrow Z\}$ is an fpqc covering we conclude that $U \rightarrow Z$ is locally quasi-finite by Descent, Lemma 20.24 as desired. \square

0428 **Lemma 10.25.** *The property $\mathcal{P}(f) = “f \text{ is syntomic}”$ is fpqc local on the base.*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 35.4. Let $Z' \rightarrow Z$ be a surjective flat morphism of affine schemes over S . Let $f : X \rightarrow Z$ be a morphism of algebraic spaces, and assume that the base change $f' : Z' \times_Z X \rightarrow Z'$ is syntomic. We have to show that f is syntomic. Let U be a scheme and let $U \rightarrow X$ be surjective and étale. By Morphisms of Spaces, Lemma 35.4 again, it is enough to show that $U \rightarrow Z$ is syntomic. Since f' is syntomic, and since $Z' \times_Z U$ is a scheme étale over $Z' \times_Z X$ we conclude (by the same lemma again) that $Z' \times_Z U \rightarrow Z'$ is syntomic. As $\{Z' \rightarrow Z\}$ is an fpqc covering we conclude that $U \rightarrow Z$ is syntomic by Descent, Lemma 20.26 as desired. \square

0429 **Lemma 10.26.** *The property $\mathcal{P}(f) = “f \text{ is smooth}”$ is fpqc local on the base.*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 36.4. Let $Z' \rightarrow Z$ be a surjective flat morphism of affine schemes over S . Let $f : X \rightarrow Z$ be a morphism of algebraic spaces, and assume that the base change $f' : Z' \times_Z X \rightarrow Z'$ is smooth. We have to show that f is smooth. Let U be a scheme and let $U \rightarrow X$ be surjective and étale. By Morphisms of Spaces, Lemma 36.4 again, it is enough to show that $U \rightarrow Z$ is smooth. Since f' is smooth, and since $Z' \times_Z U$ is a scheme étale over $Z' \times_Z X$ we conclude (by the same lemma again) that $Z' \times_Z U \rightarrow Z'$ is smooth. As $\{Z' \rightarrow Z\}$ is an fpqc covering we conclude that $U \rightarrow Z$ is smooth by Descent, Lemma 20.27 as desired. \square

042A **Lemma 10.27.** *The property $\mathcal{P}(f) = “f \text{ is unramified}”$ is fpqc local on the base.*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 37.5. Let $Z' \rightarrow Z$ be a surjective flat morphism of affine schemes over S . Let $f : X \rightarrow Z$ be a morphism of algebraic spaces, and assume that the base change $f' : Z' \times_Z X \rightarrow Z'$ is unramified. We have to show that f is unramified. Let U be a scheme and let $U \rightarrow X$ be surjective and étale. By Morphisms of Spaces, Lemma 37.5 again, it is enough to show that $U \rightarrow Z$ is unramified. Since f' is unramified, and since $Z' \times_Z U$ is a scheme étale over $Z' \times_Z X$ we conclude (by the same lemma again) that $Z' \times_Z U \rightarrow Z'$ is unramified. As $\{Z' \rightarrow Z\}$ is an fpqc covering we conclude that $U \rightarrow Z$ is unramified by Descent, Lemma 20.28 as desired. \square

042B **Lemma 10.28.** *The property $\mathcal{P}(f) = “f \text{ is étale}”$ is fpqc local on the base.*

Proof. We will use Lemma 9.4 to prove this. Assumptions (1) and (2) of that lemma follow from Morphisms of Spaces, Lemma 38.2. Let $Z' \rightarrow Z$ be a surjective flat morphism of affine schemes over S . Let $f : X \rightarrow Z$ be a morphism of algebraic spaces, and assume that the base change $f' : Z' \times_Z X \rightarrow Z'$ is étale. We have to show that f is étale. Let U be a scheme and let $U \rightarrow X$ be surjective and étale. By Morphisms of Spaces, Lemma 38.2 again, it is enough to show that $U \rightarrow Z$ is étale. Since f' is étale, and since $Z' \times_Z U$ is a scheme étale over $Z' \times_Z X$ we conclude (by the same lemma again) that $Z' \times_Z U \rightarrow Z'$ is étale. As $\{Z' \rightarrow Z\}$ is an fpqc covering we conclude that $U \rightarrow Z$ is étale by Descent, Lemma 20.29 as desired. \square

042C **Lemma 10.29.** *The property $\mathcal{P}(f) = “f \text{ is finite locally free}”$ is fpqc local on the base.*

Proof. Being finite locally free is equivalent to being finite, flat and locally of finite presentation (Morphisms of Spaces, Lemma 45.6). Hence this follows from Lemmas 10.23, 10.13, and 10.10. \square

042D **Lemma 10.30.** *The property $\mathcal{P}(f) = “f \text{ is a monomorphism}”$ is fpqc local on the base.*

Proof. Let $f : X \rightarrow Y$ be a morphism of algebraic spaces. Let $\{Y_i \rightarrow Y\}$ be an fpqc covering, and assume each of the base changes $f_i : X_i \rightarrow Y_i$ of f is a monomorphism. We have to show that f is a monomorphism.

First proof. Note that f is a monomorphism if and only if $\Delta : X \rightarrow X \times_Y X$ is an isomorphism. By applying this to f_i we see that each of the morphisms

$$\Delta_i : X_i \longrightarrow X_i \times_{Y_i} X_i = Y_i \times_Y (X \times_Y X)$$

is an isomorphism. The base change of an fpqc covering is an fpqc covering, see Topologies on Spaces, Lemma 9.3 hence $\{Y_i \times_Y (X \times_Y X) \rightarrow X \times_Y X\}$ is an fpqc covering of algebraic spaces. Moreover, each Δ_i is the base change of the morphism $\Delta : X \rightarrow X \times_Y X$. Hence it follows from Lemma 10.15 that Δ is an isomorphism, i.e., f is a monomorphism.

Second proof. Let V be a scheme, and let $V \rightarrow Y$ be a surjective étale morphism. If we can show that $V \times_Y X \rightarrow V$ is a monomorphism, then it follows that $X \rightarrow Y$ is a monomorphism. Namely, given any cartesian diagram of sheaves

$$\begin{array}{ccc} \mathcal{F} & \xrightarrow{a} & \mathcal{G} \\ b \downarrow & & \downarrow c \\ \mathcal{H} & \xrightarrow{d} & \mathcal{I} \end{array} \quad \mathcal{F} = \mathcal{H} \times_{\mathcal{I}} \mathcal{G}$$

if c is a surjection of sheaves, and a is injective, then also d is injective. This reduces the problem to the case where Y is a scheme. Moreover, in this case we may assume that the algebraic spaces Y_i are schemes also, since we can always refine the covering to place ourselves in this situation, see Topologies on Spaces, Lemma 9.5.

Assume $\{Y_i \rightarrow Y\}$ is an fpqc covering of schemes. Let $a, b : T \rightarrow X$ be two morphisms such that $f \circ a = f \circ b$. We have to show that $a = b$. Since f_i is a monomorphism we see that $a_i = b_i$, where $a_i, b_i : Y_i \times_Y T \rightarrow X_i$ are the base changes. In particular the compositions $Y_i \times_Y T \rightarrow T \rightarrow X$ are equal. Since $\{Y_i \times_Y T \rightarrow T\}$ is an fpqc covering we deduce that $a = b$ from Properties of Spaces, Proposition 16.1. \square

11. Descending properties of morphisms in the fppf topology

042E In this section we find some properties of morphisms of algebraic spaces for which we could not (yet) show they are local on the base in the fpqc topology which, however, are local on the base in the fppf topology.

042U **Lemma 11.1.** *The property $\mathcal{P}(f) = "f \text{ is an immersion}"$ is fppf local on the base.*

Proof. Let $f : X \rightarrow Y$ be a morphism of algebraic spaces. Let $\{Y_i \rightarrow Y\}_{i \in I}$ be an fppf covering of Y . Let $f_i : X_i \rightarrow Y_i$ be the base change of f .

If f is an immersion, then each f_i is an immersion by Spaces, Lemma 12.3. This proves the direct implication in Definition 9.1.

Conversely, assume each f_i is an immersion. By Morphisms of Spaces, Lemma 10.7 this implies each f_i is separated. By Morphisms of Spaces, Lemma 27.7 this implies each f_i is locally quasi-finite. Hence we see that f is locally quasi-finite and separated, by applying Lemmas 10.18 and 10.24. By Morphisms of Spaces, Lemma 49.1 this implies that f is representable!

By Morphisms of Spaces, Lemma 12.1 it suffices to show that for every scheme Z and morphism $Z \rightarrow Y$ the base change $Z \times_Y X \rightarrow Z$ is an immersion. By Topologies on Spaces, Lemma 7.4 we can find an fppf covering $\{Z_i \rightarrow Z\}$ by schemes which refines the pullback of the covering $\{Y_i \rightarrow Y\}$ to Z . Hence we see that $Z \times_Y X \rightarrow Z$ (which is a morphism of schemes according to the result of the preceding paragraph) becomes an immersion after pulling back to the members of an fppf (by schemes) of Z . Hence $Z \times_Y X \rightarrow Z$ is an immersion by the result for schemes, see Descent, Lemma 21.1. \square

042F **Lemma 11.2.** *The property $\mathcal{P}(f) = “f \text{ is locally separated}”$ is fppf local on the base.*

Proof. A base change of a locally separated morphism is locally separated, see Morphisms of Spaces, Lemma 4.4. Hence the direct implication in Definition 9.1.

Let $\{Y_i \rightarrow Y\}_{i \in I}$ be an fppf covering of algebraic spaces over S . Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . Assume each base change $X_i := Y_i \times_Y X \rightarrow Y_i$ is locally separated. This means that each of the morphisms

$$\Delta_i : X_i \longrightarrow X_i \times_{Y_i} X_i = Y_i \times_Y (X \times_Y X)$$

is an immersion. The base change of a fppf covering is an fppf covering, see Topologies on Spaces, Lemma 7.3 hence $\{Y_i \times_Y (X \times_Y X) \rightarrow X \times_Y X\}$ is an fppf covering of algebraic spaces. Moreover, each Δ_i is the base change of the morphism $\Delta : X \rightarrow X \times_Y X$. Hence it follows from Lemma 11.1 that Δ is a immersion, i.e., f is locally separated. \square

12. Application of descent of properties of morphisms

0D3B This section is the analogue of Descent, Section 22.

0D3C **Lemma 12.1.** *Let S be a scheme. Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . Let \mathcal{L} be an invertible \mathcal{O}_X -module. Let $\{g_i : Y_i \rightarrow Y\}_{i \in I}$ be an fpqc covering. Let $f_i : X_i \rightarrow Y_i$ be the base change of f and let \mathcal{L}_i be the pullback of \mathcal{L} to X_i . The following are equivalent*

- (1) \mathcal{L} is ample on X/Y , and
- (2) \mathcal{L}_i is ample on X_i/Y_i for every $i \in I$.

Proof. The implication (1) \Rightarrow (2) follows from Divisors on Spaces, Lemma 12.3. Assume (2). To check \mathcal{L} is ample on X/Y we may work étale locally on Y , see Divisors on Spaces, Lemma 12.6. Thus we may assume that Y is a scheme and then we may in turn assume each Y_i is a scheme too, see Topologies on Spaces, Lemma 9.5. In other words, we may assume that $\{Y_i \rightarrow Y\}$ is an fpqc covering of schemes.

By Divisors on Spaces, Lemma 12.4 we see that $X_i \rightarrow Y_i$ is representable (i.e., X_i is a scheme), quasi-compact, and separated. Hence f is quasi-compact and separated by Lemmas 10.1 and 10.18. This means that $\mathcal{A} = \bigoplus_{d \geq 0} f_* \mathcal{L}^{\otimes d}$ is a quasi-coherent graded \mathcal{O}_Y -algebra (Morphisms of Spaces, Lemma 11.2). Moreover, the formation of \mathcal{A} commutes with flat base change by Cohomology of Spaces, Lemma 11.2. In particular, if we set $\mathcal{A}_i = \bigoplus_{d \geq 0} f_{i,*} \mathcal{L}_i^{\otimes d}$ then we have $\mathcal{A}_i = g_i^* \mathcal{A}$. It follows that the natural maps $\psi_d : f^* \mathcal{A}_d \rightarrow \mathcal{L}^{\otimes d}$ of \mathcal{O}_X pullback to give the natural maps $\psi_{i,d} : f_i^* (\mathcal{A}_i)_d \rightarrow \mathcal{L}_i^{\otimes d}$ of \mathcal{O}_{X_i} -modules. Since \mathcal{L}_i is ample on X_i/Y_i we see that for any point $x_i \in X_i$, there exists a $d \geq 1$ such that $f_i^* (\mathcal{A}_i)_d \rightarrow \mathcal{L}_i^{\otimes d}$ is surjective on stalks at x_i . This follows either directly from the definition of a relatively ample module or from Morphisms, Lemma 35.4. If $x \in |X|$, then we can choose an i and an $x_i \in X_i$ mapping to x . Since $\mathcal{O}_{X, \bar{x}} \rightarrow \mathcal{O}_{X_i, \bar{x}_i}$ is flat hence faithfully flat, we conclude that for every $x \in |X|$ there exists a $d \geq 1$ such that $f^* \mathcal{A}_d \rightarrow \mathcal{L}^{\otimes d}$ is surjective on stalks at x . This implies that the open subset $U(\psi) \subset X$ of Divisors on Spaces, Lemma 11.1 corresponding to the map $\psi : f^* \mathcal{A} \rightarrow \bigoplus_{d \geq 0} \mathcal{L}^{\otimes d}$ of graded \mathcal{O}_X -algebras is equal to X . Consider the corresponding morphism

$$r_{\mathcal{L}, \psi} : X \longrightarrow \underline{\text{Proj}}_Y(\mathcal{A})$$

It is clear from the above that the base change of $r_{\mathcal{L},\psi}$ to Y_i is the morphism $r_{\mathcal{L}_i,\psi_i}$ which is an open immersion by Morphisms, Lemma 35.4. Hence $r_{\mathcal{L},\psi}$ is an open immersion by Lemma 10.14. Hence X is a scheme and we conclude \mathcal{L} is ample on X/Y by Morphisms, Lemma 35.4. \square

0D3D **Lemma 12.2.** *Let S be a scheme. Let $f : X \rightarrow Y$ be a proper morphism of algebraic spaces over S . Let \mathcal{L} be an invertible \mathcal{O}_X -module. There exists an open subspace $V \subset Y$ characterized by the following property: A morphism $Y' \rightarrow Y$ of algebraic spaces factors through V if and only if the pullback \mathcal{L}' of \mathcal{L} to $X' = Y' \times_Y X$ is ample on X'/Y' (as in Divisors on Spaces, Definition 12.1).*

Proof. Suppose that the lemma holds whenever Y is a scheme. Let U be a scheme and let $U \rightarrow Y$ be a surjective étale morphism. Let $R = U \times_Y U$ with projections $t, s : R \rightarrow U$. Denote $X_U = U \times_Y X$ and \mathcal{L}_U the pullback. Then we get an open subscheme $V' \subset U$ as in the lemma for $(X_U \rightarrow U, \mathcal{L}_U)$. By the functorial characterization we see that $s^{-1}(V') = t^{-1}(V')$. Thus there is an open subspace $V \subset Y$ such that V' is the inverse image of V in U . In particular $V' \rightarrow V$ is surjective étale and we conclude that \mathcal{L}_V is ample on X_V/V (Divisors on Spaces, Lemma 12.6). Now, if $Y' \rightarrow Y$ is a morphism such that \mathcal{L}' is ample on X'/Y' , then $U \times_Y Y' \rightarrow Y'$ must factor through V' and we conclude that $Y' \rightarrow Y$ factors through V . Hence $V \subset Y$ is as in the statement of the lemma. In this way we reduce to the case dealt with in the next paragraph.

Assume Y is a scheme. Since the question is local on Y we may assume Y is an affine scheme. We will show the following:

- (A) If $\text{Spec}(k) \rightarrow Y$ is a morphism such that \mathcal{L}_k is ample on X_k/k , then there is an open neighbourhood $V \subset Y$ of the image of $\text{Spec}(k) \rightarrow Y$ such that \mathcal{L}_V is ample on X_V/V .

It is clear that (A) implies the truth of the lemma.

Let $X \rightarrow Y$, \mathcal{L} , $\text{Spec}(k) \rightarrow Y$ be as in (A). By Lemma 12.1 we may assume that $k = \kappa(y)$ is the residue field of a point y of Y .

As Y is affine we can find a directed set I and an inverse system of morphisms $X_i \rightarrow Y_i$ of algebraic spaces with Y_i of finite presentation over \mathbf{Z} , with affine transition morphisms $X_i \rightarrow X_{i'}$ and $Y_i \rightarrow Y_{i'}$, with $X_i \rightarrow Y_i$ proper and of finite presentation, and such that $X \rightarrow Y = \lim(X_i \rightarrow Y_i)$. See Limits of Spaces, Lemma 12.2. After shrinking I we may assume Y_i is an (affine) scheme for all i , see Limits of Spaces, Lemma 5.10. After shrinking I we can assume we have a compatible system of invertible \mathcal{O}_{X_i} -modules \mathcal{L}_i pulling back to \mathcal{L} , see Limits of Spaces, Lemma 7.3. Let $y_i \in Y_i$ be the image of y . Then $\kappa(y) = \text{colim } \kappa(y_i)$. Hence $X_y = \lim X_{i,y_i}$ and after shrinking I we may assume X_{i,y_i} is a scheme for all i , see Limits of Spaces, Lemma 5.11. Hence for some i we have \mathcal{L}_{i,y_i} is ample on X_{i,y_i} by Limits, Lemma 4.15. By Divisors on Spaces, Lemma 13.3 we find an open neighbourhood $V_i \subset Y_i$ of y_i such that \mathcal{L}_i restricted to $f_i^{-1}(V_i)$ is ample relative to V_i . Letting $V \subset Y$ be the inverse image of V_i finishes the proof (hints: use Morphisms, Lemma 35.9 and the fact that $X \rightarrow Y \times_{Y_i} X_i$ is affine and the fact that the pullback of an ample invertible sheaf by an affine morphism is ample by Morphisms, Lemma 35.7). \square

13. Properties of morphisms local on the source

In this section we define what it means for a property of morphisms of algebraic spaces to be local on the source. Please compare with Descent, Section 23.

06EP **Definition 13.1.** Let S be a scheme. Let \mathcal{P} be a property of morphisms of algebraic spaces over S . Let $\tau \in \{fpqc, fppf, syntomic, smooth, \acute{e}tale\}$. We say \mathcal{P} is τ local on the source, or local on the source for the τ -topology if for any morphism $f : X \rightarrow Y$ of algebraic spaces over S , and any τ -covering $\{X_i \rightarrow X\}_{i \in I}$ of algebraic spaces we have

$$f \text{ has } \mathcal{P} \Leftrightarrow \text{each } X_i \rightarrow Y \text{ has } \mathcal{P}.$$

To be sure, since isomorphisms are always coverings we see (or require) that property \mathcal{P} holds for $X \rightarrow Y$ if and only if it holds for any arrow $X' \rightarrow Y'$ isomorphic to $X \rightarrow Y$. If a property is τ -local on the source then it is preserved by precomposing with morphisms which occur in τ -coverings. Here is a formal statement.

06EQ **Lemma 13.2.** *Let S be a scheme. Let $\tau \in \{fpqc, fppf, syntomic, smooth, \acute{e}tale\}$. Let \mathcal{P} be a property of morphisms of algebraic spaces over S which is τ local on the source. Let $f : X \rightarrow Y$ have property \mathcal{P} . For any morphism $a : X' \rightarrow X$ which is flat, resp. flat and locally of finite presentation, resp. syntomic, resp. smooth, resp. $\acute{e}tale$, the composition $f \circ a : X' \rightarrow Y$ has property \mathcal{P} .*

Proof. This is true because we can fit $X' \rightarrow X$ into a family of morphisms which forms a τ -covering. \square

06ER **Lemma 13.3.** *Let S be a scheme. Let $\tau \in \{fpqc, fppf, syntomic, smooth, \acute{e}tale\}$. Suppose that \mathcal{P} is a property of morphisms of schemes over S which is $\acute{e}tale$ local on the source-and-target. Denote \mathcal{P}_{spaces} the corresponding property of morphisms of algebraic spaces over S , see Morphisms of Spaces, Definition 22.2. If \mathcal{P} is local on the source for the τ -topology, then \mathcal{P}_{spaces} is local on the source for the τ -topology.*

Proof. Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . Let $\{X_i \rightarrow X\}_{i \in I}$ be a τ -covering of algebraic spaces. Choose a scheme V and a surjective $\acute{e}tale$ morphism $V \rightarrow Y$. Choose a scheme U and a surjective $\acute{e}tale$ morphism $U \rightarrow X \times_Y V$. For each i choose a scheme U_i and a surjective $\acute{e}tale$ morphism $U_i \rightarrow X_i \times_X U$.

Note that $\{X_i \times_X U \rightarrow U\}_{i \in I}$ is a τ -covering. Note that each $\{U_i \rightarrow X_i \times_X U\}$ is an $\acute{e}tale$ covering, hence a τ -covering. Hence $\{U_i \rightarrow U\}_{i \in I}$ is a τ -covering of algebraic spaces over S . But since U and each U_i is a scheme we see that $\{U_i \rightarrow U\}_{i \in I}$ is a τ -covering of schemes over S .

Now we have

$$\begin{aligned} f \text{ has } \mathcal{P}_{spaces} &\Leftrightarrow U \rightarrow V \text{ has } \mathcal{P} \\ &\Leftrightarrow \text{each } U_i \rightarrow V \text{ has } \mathcal{P} \\ &\Leftrightarrow \text{each } X_i \rightarrow Y \text{ has } \mathcal{P}_{spaces}. \end{aligned}$$

the first and last equivalence by the definition of \mathcal{P}_{spaces} the middle equivalence because we assumed \mathcal{P} is local on the source in the τ -topology. \square

14. Properties of morphisms local in the fpqc topology on the source

06ES Here are some properties of morphisms that are fpqc local on the source.

06ET **Lemma 14.1.** *The property $\mathcal{P}(f) = \text{“}f \text{ is flat”}$ is fpqc local on the source.*

Proof. Follows from Lemma 13.3 using Morphisms of Spaces, Definition 29.1 and Descent, Lemma 24.1. \square

15. Properties of morphisms local in the fppf topology on the source

06EU Here are some properties of morphisms that are fppf local on the source.

06EV **Lemma 15.1.** *The property $\mathcal{P}(f) = “f \text{ is locally of finite presentation}”$ is fppf local on the source.*

Proof. Follows from Lemma 13.3 using Morphisms of Spaces, Definition 28.1 and Descent, Lemma 25.1. \square

06EW **Lemma 15.2.** *The property $\mathcal{P}(f) = “f \text{ is locally of finite type}”$ is fppf local on the source.*

Proof. Follows from Lemma 13.3 using Morphisms of Spaces, Definition 23.1 and Descent, Lemma 25.2. \square

06EX **Lemma 15.3.** *The property $\mathcal{P}(f) = “f \text{ is open}”$ is fppf local on the source.*

Proof. Follows from Lemma 13.3 using Morphisms of Spaces, Definition 6.2 and Descent, Lemma 25.3. \square

06EY **Lemma 15.4.** *The property $\mathcal{P}(f) = “f \text{ is universally open}”$ is fppf local on the source.*

Proof. Follows from Lemma 13.3 using Morphisms of Spaces, Definition 6.2 and Descent, Lemma 25.4. \square

16. Properties of morphisms local in the syntomic topology on the source

06EZ Here are some properties of morphisms that are syntomic local on the source.

06F0 **Lemma 16.1.** *The property $\mathcal{P}(f) = “f \text{ is syntomic}”$ is syntomic local on the source.*

Proof. Follows from Lemma 13.3 using Morphisms of Spaces, Definition 35.1 and Descent, Lemma 26.1. \square

17. Properties of morphisms local in the smooth topology on the source

06F1 Here are some properties of morphisms that are smooth local on the source.

06F2 **Lemma 17.1.** *The property $\mathcal{P}(f) = “f \text{ is smooth}”$ is smooth local on the source.*

Proof. Follows from Lemma 13.3 using Morphisms of Spaces, Definition 36.1 and Descent, Lemma 27.1. \square

18. Properties of morphisms local in the étale topology on the source

06F3 Here are some properties of morphisms that are étale local on the source.

06F4 **Lemma 18.1.** *The property $\mathcal{P}(f) = “f \text{ is étale}”$ is étale local on the source.*

Proof. Follows from Lemma 13.3 using Morphisms of Spaces, Definition 38.1 and Descent, Lemma 28.1. \square

06F5 **Lemma 18.2.** *The property $\mathcal{P}(f) = “f \text{ is locally quasi-finite}”$ is étale local on the source.*

Proof. Follows from Lemma 13.3 using Morphisms of Spaces, Definition 27.1 and Descent, Lemma 28.2. \square

06F6 **Lemma 18.3.** *The property $\mathcal{P}(f) = “f \text{ is unramified}”$ is étale local on the source.*

Proof. Follows from Lemma 13.3 using Morphisms of Spaces, Definition 37.1 and Descent, Lemma 28.3. \square

19. Properties of morphisms smooth local on source-and-target

06F7 Let \mathcal{P} be a property of morphisms of algebraic spaces. There is an intuitive meaning to the phrase “ \mathcal{P} is smooth local on the source and target”. However, it turns out that this notion is not the same as asking \mathcal{P} to be both smooth local on the source and smooth local on the target. We have discussed a similar phenomenon (for the étale topology and the category of schemes) in great detail in Descent, Section 29 (for a quick overview take a look at Descent, Remark 29.8). However, there is an important difference between the case of the smooth and the étale topology. To see this difference we encourage the reader to ponder the difference between Descent, Lemma 29.4 and Lemma 19.2 as well as the difference between Descent, Lemma 29.5 and Lemma 19.3. Namely, in the étale setting the choice of the étale “covering” of the target is immaterial, whereas in the smooth setting it is not.

06F8 **Definition 19.1.** Let S be a scheme. Let \mathcal{P} be a property of morphisms of algebraic spaces over S . We say \mathcal{P} is *smooth local on source-and-target* if

- (1) (stable under precomposing with smooth maps) if $f : X \rightarrow Y$ is smooth and $g : Y \rightarrow Z$ has \mathcal{P} , then $g \circ f$ has \mathcal{P} ,
- (2) (stable under smooth base change) if $f : X \rightarrow Y$ has \mathcal{P} and $Y' \rightarrow Y$ is smooth, then the base change $f' : Y' \times_Y X \rightarrow Y'$ has \mathcal{P} , and
- (3) (locality) given a morphism $f : X \rightarrow Y$ the following are equivalent
 - (a) f has \mathcal{P} ,
 - (b) for every $x \in |X|$ there exists a commutative diagram

$$\begin{array}{ccc} U & \xrightarrow{\quad} & V \\ a \downarrow & \searrow h & \downarrow b \\ X & \xrightarrow{f} & Y \end{array}$$

with smooth vertical arrows and $u \in |U|$ with $a(u) = x$ such that h has \mathcal{P} .

The above serves as our definition. In the lemmas below we will show that this is equivalent to \mathcal{P} being smooth local on the target, smooth local on the source, and stable under post-composing by smooth morphisms.

06F9 **Lemma 19.2.** *Let S be a scheme. Let \mathcal{P} be a property of morphisms of algebraic spaces over S which is smooth local on source-and-target. Then*

- (1) \mathcal{P} is smooth local on the source,
- (2) \mathcal{P} is smooth local on the target,
- (3) \mathcal{P} is stable under postcomposing with smooth morphisms: if $f : X \rightarrow Y$ has \mathcal{P} and $g : Y \rightarrow Z$ is smooth, then $g \circ f$ has \mathcal{P} .

Proof. We write everything out completely.

Proof of (1). Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . Let $\{X_i \rightarrow X\}_{i \in I}$ be a smooth covering of X . If each composition $h_i : X_i \rightarrow Y$ has \mathcal{P} , then for each $|x| \in X$ we can find an $i \in I$ and a point $x_i \in |X_i|$ mapping to x . Then $(X_i, x_i) \rightarrow (X, x)$ is a smooth morphism of pairs, and $\text{id}_Y : Y \rightarrow Y$ is a smooth morphism, and h_i is as in part (3) of Definition 19.1. Thus we see that f has \mathcal{P} . Conversely, if f has \mathcal{P} then each $X_i \rightarrow Y$ has \mathcal{P} by Definition 19.1 part (1).

Proof of (2). Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . Let $\{Y_i \rightarrow Y\}_{i \in I}$ be a smooth covering of Y . Write $X_i = Y_i \times_Y X$ and $h_i : X_i \rightarrow Y_i$ for the base change of f . If each $h_i : X_i \rightarrow Y_i$ has \mathcal{P} , then for each $x \in |X|$ we pick an $i \in I$ and a point $x_i \in |X_i|$ mapping to x . Then $(X_i, x_i) \rightarrow (X, x)$ is a smooth morphism of pairs, $Y_i \rightarrow Y$ is smooth, and h_i is as in part (3) of Definition 19.1. Thus we see that f has \mathcal{P} . Conversely, if f has \mathcal{P} , then each $X_i \rightarrow Y_i$ has \mathcal{P} by Definition 19.1 part (2).

Proof of (3). Assume $f : X \rightarrow Y$ has \mathcal{P} and $g : Y \rightarrow Z$ is smooth. For every $x \in |X|$ we can think of $(X, x) \rightarrow (X, x)$ as a smooth morphism of pairs, $Y \rightarrow Z$ is a smooth morphism, and $h = f$ is as in part (3) of Definition 19.1. Thus we see that $g \circ f$ has \mathcal{P} . \square

The following lemma is the analogue of Morphisms, Lemma 13.4.

06FA **Lemma 19.3.** *Let S be a scheme. Let \mathcal{P} be a property of morphisms of algebraic spaces over S which is smooth local on source-and-target. Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . The following are equivalent:*

- (a) f has property \mathcal{P} ,
- (b) for every $x \in |X|$ there exists a smooth morphism of pairs $a : (U, u) \rightarrow (X, x)$, a smooth morphism $b : V \rightarrow Y$, and a morphism $h : U \rightarrow V$ such that $f \circ a = b \circ h$ and h has \mathcal{P} ,
- (c) for some commutative diagram

$$\begin{array}{ccc} U & \longrightarrow & V \\ a \downarrow & & \downarrow b \\ X & \xrightarrow{f} & Y \end{array}$$

with a, b smooth and a surjective the morphism h has \mathcal{P} ,

- (d) for any commutative diagram

$$\begin{array}{ccc} U & \longrightarrow & V \\ a \downarrow & & \downarrow b \\ X & \xrightarrow{f} & Y \end{array}$$

with b smooth and $U \rightarrow X \times_Y V$ smooth the morphism h has \mathcal{P} ,

- (e) *there exists a smooth covering $\{Y_i \rightarrow Y\}_{i \in I}$ such that each base change $Y_i \times_Y X \rightarrow Y_i$ has \mathcal{P} ,*
- (f) *there exists a smooth covering $\{X_i \rightarrow X\}_{i \in I}$ such that each composition $X_i \rightarrow Y$ has \mathcal{P} ,*
- (g) *there exists a smooth covering $\{Y_i \rightarrow Y\}_{i \in I}$ and for each $i \in I$ a smooth covering $\{X_{ij} \rightarrow Y_i \times_Y X\}_{j \in J_i}$ such that each morphism $X_{ij} \rightarrow Y_i$ has \mathcal{P} .*

Proof. The equivalence of (a) and (b) is part of Definition 19.1. The equivalence of (a) and (e) is Lemma 19.2 part (2). The equivalence of (a) and (f) is Lemma 19.2 part (1). As (a) is now equivalent to (e) and (f) it follows that (a) equivalent to (g).

It is clear that (c) implies (b). If (b) holds, then for any $x \in |X|$ we can choose a smooth morphism of pairs $a_x : (U_x, u_x) \rightarrow (X, x)$, a smooth morphism $b_x : V_x \rightarrow Y$, and a morphism $h_x : U_x \rightarrow V_x$ such that $f \circ a_x = b_x \circ h_x$ and h_x has \mathcal{P} . Then $h = \coprod h_x : \coprod U_x \rightarrow \coprod V_x$ with $a = \coprod a_x$ and $b = \coprod b_x$ is a diagram as in (c). (Note that h has property \mathcal{P} as $\{V_x \rightarrow \coprod V_x\}$ is a smooth covering and \mathcal{P} is smooth local on the target.) Thus (b) is equivalent to (c).

Now we know that (a), (b), (c), (e), (f), and (g) are equivalent. Suppose (a) holds. Let U, V, a, b, h be as in (d). Then $X \times_Y V \rightarrow V$ has \mathcal{P} as \mathcal{P} is stable under smooth base change, whence $U \rightarrow V$ has \mathcal{P} as \mathcal{P} is stable under precomposing with smooth morphisms. Conversely, if (d) holds, then setting $U = X$ and $V = Y$ we see that f has \mathcal{P} . \square

06FB **Lemma 19.4.** *Let S be a scheme. Let \mathcal{P} be a property of morphisms of algebraic spaces over S . Assume*

- (1) *\mathcal{P} is smooth local on the source,*
- (2) *\mathcal{P} is smooth local on the target, and*
- (3) *\mathcal{P} is stable under postcomposing with smooth morphisms: if $f : X \rightarrow Y$ has \mathcal{P} and $Y \rightarrow Z$ is a smooth morphism then $X \rightarrow Z$ has \mathcal{P} .*

Then \mathcal{P} is smooth local on the source-and-target.

Proof. Let \mathcal{P} be a property of morphisms of algebraic spaces which satisfies conditions (1), (2) and (3) of the lemma. By Lemma 13.2 we see that \mathcal{P} is stable under precomposing with smooth morphisms. By Lemma 9.2 we see that \mathcal{P} is stable under smooth base change. Hence it suffices to prove part (3) of Definition 19.1 holds.

More precisely, suppose that $f : X \rightarrow Y$ is a morphism of algebraic spaces over S which satisfies Definition 19.1 part (3)(b). In other words, for every $x \in X$ there exists a smooth morphism $a_x : U_x \rightarrow X$, a point $u_x \in |U_x|$ mapping to x , a smooth morphism $b_x : V_x \rightarrow Y$, and a morphism $h_x : U_x \rightarrow V_x$ such that $f \circ a_x = b_x \circ h_x$ and h_x has \mathcal{P} . The proof of the lemma is complete once we show that f has \mathcal{P} . Set $U = \coprod U_x$, $a = \coprod a_x$, $V = \coprod V_x$, $b = \coprod b_x$, and $h = \coprod h_x$. We obtain a commutative diagram

$$\begin{array}{ccc} U & \xrightarrow{h} & V \\ a \downarrow & & \downarrow b \\ X & \xrightarrow{f} & Y \end{array}$$

with a, b smooth, a surjective. Note that h has \mathcal{P} as each h_x does and \mathcal{P} is smooth local on the target. Because a is surjective and \mathcal{P} is smooth local on the source, it suffices to prove that $b \circ h$ has \mathcal{P} . This follows as we assumed that \mathcal{P} is stable under postcomposing with a smooth morphism and as b is smooth. \square

06FC **Remark 19.5.** Using Lemma 19.4 and the work done in the earlier sections of this chapter it is easy to make a list of types of morphisms which are smooth local on the source-and-target. In each case we list the lemma which implies the property is smooth local on the source and the lemma which implies the property is smooth local on the target. In each case the third assumption of Lemma 19.4 is trivial to check, and we omit it. Here is the list:

- (1) flat, see Lemmas 14.1 and 10.13,
- (2) locally of finite presentation, see Lemmas 15.1 and 10.10,
- (3) locally finite type, see Lemmas 15.2 and 10.9,
- (4) universally open, see Lemmas 15.4 and 10.4,
- (5) syntomic, see Lemmas 16.1 and 10.25,
- (6) smooth, see Lemmas 17.1 and 10.26,
- (7) add more here as needed.

20. Properties of morphisms étale-smooth local on source-and-target

0CFY This section is the analogue of Section 19 for properties of morphisms which are étale local on the source and smooth local on the target. We give this property a ridiculously long name in order to avoid using it too much.

0CFZ **Definition 20.1.** Let S be a scheme. Let \mathcal{P} be a property of morphisms of algebraic spaces over S . We say \mathcal{P} is *étale-smooth local on source-and-target* if

- (1) (stable under precomposing with étale maps) if $f : X \rightarrow Y$ is étale and $g : Y \rightarrow Z$ has \mathcal{P} , then $g \circ f$ has \mathcal{P} ,
- (2) (stable under smooth base change) if $f : X \rightarrow Y$ has \mathcal{P} and $Y' \rightarrow Y$ is smooth, then the base change $f' : Y' \times_Y X \rightarrow Y'$ has \mathcal{P} , and
- (3) (locality) given a morphism $f : X \rightarrow Y$ the following are equivalent
 - (a) f has \mathcal{P} ,
 - (b) for every $x \in |X|$ there exists a commutative diagram

$$\begin{array}{ccc} U & \xrightarrow{\quad} & V \\ a \downarrow & \quad h \quad & \downarrow b \\ X & \xrightarrow{\quad f \quad} & Y \end{array}$$

with b smooth and $U \rightarrow X \times_Y V$ étale and $u \in |U|$ with $a(u) = x$ such that h has \mathcal{P} .

The above serves as our definition. In the lemmas below we will show that this is equivalent to \mathcal{P} being étale local on the target, smooth local on the source, and stable under post-composing by étale morphisms.

0CG0 **Lemma 20.2.** *Let S be a scheme. Let \mathcal{P} be a property of morphisms of algebraic spaces over S which is étale-smooth local on source-and-target. Then*

- (1) \mathcal{P} is étale local on the source,
- (2) \mathcal{P} is smooth local on the target,

- (3) \mathcal{P} is stable under postcomposing with étale morphisms: if $f : X \rightarrow Y$ has \mathcal{P} and $g : Y \rightarrow Z$ is étale, then $g \circ f$ has \mathcal{P} , and
- (4) \mathcal{P} has a permanence property: given $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ étale such that $g \circ f$ has \mathcal{P} , then f has \mathcal{P} .

Proof. We write everything out completely.

Proof of (1). Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . Let $\{X_i \rightarrow X\}_{i \in I}$ be an étale covering of X . If each composition $h_i : X_i \rightarrow Y$ has \mathcal{P} , then for each $|x| \in X$ we can find an $i \in I$ and a point $x_i \in |X_i|$ mapping to x . Then $(X_i, x_i) \rightarrow (X, x)$ is an étale morphism of pairs, and $\text{id}_Y : Y \rightarrow Y$ is a smooth morphism, and h_i is as in part (3) of Definition 20.1. Thus we see that f has \mathcal{P} . Conversely, if f has \mathcal{P} then each $X_i \rightarrow Y$ has \mathcal{P} by Definition 20.1 part (1).

Proof of (2). Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . Let $\{Y_i \rightarrow Y\}_{i \in I}$ be a smooth covering of Y . Write $X_i = Y_i \times_Y X$ and $h_i : X_i \rightarrow Y_i$ for the base change of f . If each $h_i : X_i \rightarrow Y_i$ has \mathcal{P} , then for each $x \in |X|$ we pick an $i \in I$ and a point $x_i \in |X_i|$ mapping to x . Then $X_i \rightarrow X \times_Y Y_i$ is an étale morphism (because it is an isomorphism), $Y_i \rightarrow Y$ is smooth, and h_i is as in part (3) of Definition 19.1. Thus we see that f has \mathcal{P} . Conversely, if f has \mathcal{P} , then each $X_i \rightarrow Y_i$ has \mathcal{P} by Definition 19.1 part (2).

Proof of (3). Assume $f : X \rightarrow Y$ has \mathcal{P} and $g : Y \rightarrow Z$ is étale. The morphism $X \rightarrow Y \times_Z X$ is étale as a morphism between algebraic spaces étale over X (Properties of Spaces, Lemma 15.6). Also $Y \rightarrow Z$ is étale hence a smooth morphism. Thus the diagram

$$\begin{array}{ccc} X & \xrightarrow{\quad} & Y \\ \downarrow & \searrow f & \downarrow \\ X & \xrightarrow{g \circ f} & Z \end{array}$$

works for every $x \in |X|$ in part (3) of Definition 19.1 and we conclude that $g \circ f$ has \mathcal{P} .

Proof of (4). Let $f : X \rightarrow Y$ be a morphism and $g : Y \rightarrow Z$ étale such that $g \circ f$ has \mathcal{P} . Then by Definition 20.1 part (2) we see that $\text{pr}_Y : Y \times_Z X \rightarrow Y$ has \mathcal{P} . But the morphism $(f, 1) : X \rightarrow Y \times_Z X$ is étale as a section to the étale projection $\text{pr}_X : Y \times_Z X \rightarrow X$, see Morphisms of Spaces, Lemma 38.11. Hence $f = \text{pr}_Y \circ (f, 1)$ has \mathcal{P} by Definition 20.1 part (1). \square

0CG1 **Lemma 20.3.** *Let S be a scheme. Let \mathcal{P} be a property of morphisms of algebraic spaces over S which is étale-smooth local on source-and-target. Let $f : X \rightarrow Y$ be a morphism of algebraic spaces over S . The following are equivalent:*

- (a) f has property \mathcal{P} ,
- (b) for every $x \in |X|$ there exists a smooth morphism $b : V \rightarrow Y$, an étale morphism $a : U \rightarrow V \times_Y X$, and a point $u \in |U|$ mapping to x such that $U \rightarrow V$ has \mathcal{P} ,
- (c) for some commutative diagram

$$\begin{array}{ccc} U & \xrightarrow{\quad} & V \\ a \downarrow & \searrow h & \downarrow b \\ X & \xrightarrow{\quad} & Y \end{array}$$

with b smooth, $U \rightarrow V \times_Y X$ étale, and a surjective the morphism h has \mathcal{P} ,

(d) for any commutative diagram

$$\begin{array}{ccc} U & \xrightarrow{\quad} & V \\ a \downarrow & \searrow h & \downarrow b \\ X & \xrightarrow{\quad f \quad} & Y \end{array}$$

with b smooth and $U \rightarrow X \times_Y V$ étale, the morphism h has \mathcal{P} ,

- (e) there exists a smooth covering $\{Y_i \rightarrow Y\}_{i \in I}$ such that each base change $Y_i \times_Y X \rightarrow Y_i$ has \mathcal{P} ,
- (f) there exists an étale covering $\{X_i \rightarrow X\}_{i \in I}$ such that each composition $X_i \rightarrow Y$ has \mathcal{P} ,
- (g) there exists a smooth covering $\{Y_i \rightarrow Y\}_{i \in I}$ and for each $i \in I$ an étale covering $\{X_{ij} \rightarrow Y_i \times_Y X\}_{j \in J_i}$ such that each morphism $X_{ij} \rightarrow Y_i$ has \mathcal{P} .

Proof. The equivalence of (a) and (b) is part of Definition 20.1. The equivalence of (a) and (e) is Lemma 20.2 part (2). The equivalence of (a) and (f) is Lemma 20.2 part (1). As (a) is now equivalent to (e) and (f) it follows that (a) equivalent to (g).

It is clear that (c) implies (b). If (b) holds, then for any $x \in |X|$ we can choose a smooth morphism a smooth morphism $b_x : V_x \rightarrow Y$, an étale morphism $U_x \rightarrow V_x \times_Y X$, and $u_x \in |U_x|$ mapping to x such that $U_x \rightarrow V_x$ has \mathcal{P} . Then $h = \coprod h_x : \coprod U_x \rightarrow \coprod V_x$ with $a = \coprod a_x$ and $b = \coprod b_x$ is a diagram as in (c). (Note that h has property \mathcal{P} as $\{V_x \rightarrow \coprod V_x\}$ is a smooth covering and \mathcal{P} is smooth local on the target.) Thus (b) is equivalent to (c).

Now we know that (a), (b), (c), (e), (f), and (g) are equivalent. Suppose (a) holds. Let U, V, a, b, h be as in (d). Then $X \times_Y V \rightarrow V$ has \mathcal{P} as \mathcal{P} is stable under smooth base change, whence $U \rightarrow V$ has \mathcal{P} as \mathcal{P} is stable under precomposing with étale morphisms. Conversely, if (d) holds, then setting $U = X$ and $V = Y$ we see that f has \mathcal{P} . \square

0CG2 **Lemma 20.4.** *Let S be a scheme. Let \mathcal{P} be a property of morphisms of algebraic spaces over S . Assume*

- (1) \mathcal{P} is étale local on the source,
- (2) \mathcal{P} is smooth local on the target, and
- (3) \mathcal{P} is stable under postcomposing with open immersions: if $f : X \rightarrow Y$ has \mathcal{P} and $Y \subset Z$ is an open embedding then $X \rightarrow Z$ has \mathcal{P} .

Then \mathcal{P} is étale-smooth local on the source-and-target.

Proof. Let \mathcal{P} be a property of morphisms of algebraic spaces which satisfies conditions (1), (2) and (3) of the lemma. By Lemma 13.2 we see that \mathcal{P} is stable under precomposing with étale morphisms. By Lemma 9.2 we see that \mathcal{P} is stable under smooth base change. Hence it suffices to prove part (3) of Definition 19.1 holds.

More precisely, suppose that $f : X \rightarrow Y$ is a morphism of algebraic spaces over S which satisfies Definition 19.1 part (3)(b). In other words, for every $x \in X$ there exists a smooth morphism $b_x : V_x \rightarrow Y$, an étale morphism $U_x \rightarrow V_x \times_Y X$, and a point $u_x \in |U_x|$ mapping to x such that $h_x : U_x \rightarrow V_x$ has \mathcal{P} . The proof of the lemma is complete once we show that f has \mathcal{P} .

Let $a_x : U_x \rightarrow X$ be the composition $U_x \rightarrow V_x \times_Y X \rightarrow X$. Set $U = \coprod U_x$, $a = \coprod a_x$, $V = \coprod V_x$, $b = \coprod b_x$, and $h = \coprod h_x$. We obtain a commutative diagram

$$\begin{array}{ccc} U & \xrightarrow{\quad} & V \\ a \downarrow & \searrow h & \downarrow b \\ X & \xrightarrow{\quad f \quad} & Y \end{array}$$

with b smooth, $U \rightarrow V \times_Y X$ étale, a surjective. Note that h has \mathcal{P} as each h_x does and \mathcal{P} is smooth local on the target. In the next paragraph we prove that we may assume U, V, X, Y are schemes; we encourage the reader to skip it.

Let X, Y, U, V, a, b, f, h be as in the previous paragraph. We have to show f has \mathcal{P} . Let $X' \rightarrow X$ be a surjective étale morphism with X_i a scheme. Set $U' = X' \times_X U$. Then $U' \rightarrow X'$ is surjective and $U' \rightarrow X' \times_Y V$ is étale. Since \mathcal{P} is étale local on the source, we see that $U' \rightarrow V$ has \mathcal{P} and that it suffices to show that $X' \rightarrow Y$ has \mathcal{P} . In other words, we may assume that X is a scheme. Next, choose a surjective étale morphism $Y' \rightarrow Y$ with Y' a scheme. Set $V' = V \times_Y Y'$, $X' = X \times_Y Y'$, and $U' = U \times_Y Y'$. Then $U' \rightarrow X'$ is surjective and $U' \rightarrow X' \times_{Y'} V'$ is étale. Since \mathcal{P} is smooth local on the target, we see that $U' \rightarrow V'$ has \mathcal{P} and that it suffices to prove $X' \rightarrow Y'$ has \mathcal{P} . Thus we may assume both X and Y are schemes. Choose a surjective étale morphism $V' \rightarrow V$ with V' a scheme. Set $U' = U \times_V V'$. Then $U' \rightarrow X$ is surjective and $U' \rightarrow X \times_Y V'$ is étale. Since \mathcal{P} is smooth local on the source, we see that $U' \rightarrow V'$ has \mathcal{P} . Thus we may replace U, V by U', V' and assume X, Y, V are schemes. Finally, we replace U by a scheme surjective étale over U and we see that we may assume U, V, X, Y are all schemes.

If U, V, X, Y are schemes, then f has \mathcal{P} by Descent, Lemma 29.11. □

0CG3 **Remark 20.5.** Using Lemma 20.4 and the work done in the earlier sections of this chapter it is easy to make a list of types of morphisms which are smooth local on the source-and-target. In each case we list the lemma which implies the property is étale local on the source and the lemma which implies the property is smooth local on the target. In each case the third assumption of Lemma 20.4 is trivial to check, and we omit it. Here is the list:

- (1) étale, see Lemmas 18.1 and 10.28,
- (2) locally quasi-finite, see Lemmas 18.2 and 10.24,
- (3) unramified, see Lemmas 18.3 and 10.27, and
- (4) add more here as needed.

Of course any property listed in Remark 19.5 is a fortiori an example that could be listed here.

21. Descent data for spaces over spaces

0ADF This section is the analogue of Descent, Section 31 for algebraic spaces. Most of the arguments in this section are formal relying only on the definition of a descent datum.

0ADG **Definition 21.1.** Let S be a scheme. Let $f : Y \rightarrow X$ be a morphism of algebraic spaces over S .

- (1) Let $V \rightarrow Y$ be a morphism of algebraic spaces. A *descent datum* for $V/Y/X$ is an isomorphism $\varphi : V \times_X Y \rightarrow Y \times_X V$ of algebraic spaces over $Y \times_X Y$ satisfying the *cocycle condition* that the diagram

$$\begin{array}{ccc} V \times_X Y \times_X Y & \xrightarrow{\varphi_{02}} & Y \times_X Y \times_X V \\ & \searrow \varphi_{01} & \nearrow \varphi_{12} \\ & Y \times_X Y \times_X Y & \end{array}$$

commutes (with obvious notation).

- (2) We also say that the pair $(V/Y, \varphi)$ is a *descent datum relative to* $Y \rightarrow X$.
(3) A *morphism* $f : (V/Y, \varphi) \rightarrow (V'/Y, \varphi')$ of descent data relative to $Y \rightarrow X$ is a morphism $f : V \rightarrow V'$ of algebraic spaces over Y such that the diagram

$$\begin{array}{ccc} V \times_X Y & \xrightarrow{\varphi} & Y \times_X V \\ f \times \text{id}_Y \downarrow & & \downarrow \text{id}_Y \times f \\ V' \times_X Y & \xrightarrow{\varphi'} & Y \times_X V' \end{array}$$

commutes.

0ADH **Remark 21.2.** Let S be a scheme. Let $Y \rightarrow X$ be a morphism of algebraic spaces over S . Let $(V/Y, \varphi)$ be a descent datum relative to $Y \rightarrow X$. We may think of the isomorphism φ as an isomorphism

$$(Y \times_X Y) \times_{\text{pr}_0, Y} V \longrightarrow (Y \times_X Y) \times_{\text{pr}_1, Y} V$$

of algebraic spaces over $Y \times_X Y$. So loosely speaking one may think of φ as a map $\varphi : \text{pr}_0^* V \rightarrow \text{pr}_1^* V$. The cocycle condition then says that $\text{pr}_{02}^* \varphi = \text{pr}_{12}^* \varphi \circ \text{pr}_{01}^* \varphi$. In this way it is very similar to the case of a descent datum on quasi-coherent sheaves.

Here is the definition in case you have a family of morphisms with fixed target.

0ADI **Definition 21.3.** Let S be a scheme. Let $\{X_i \rightarrow X\}_{i \in I}$ be a family of morphisms of algebraic spaces over S with fixed target X .

- (1) A *descent datum* (V_i, φ_{ij}) relative to the family $\{X_i \rightarrow X\}$ is given by an algebraic space V_i over X_i for each $i \in I$, an isomorphism $\varphi_{ij} : V_i \times_X X_j \rightarrow X_i \times_X V_j$ of algebraic spaces over $X_i \times_X X_j$ for each pair $(i, j) \in I^2$ such that for every triple of indices $(i, j, k) \in I^3$ the diagram

$$\begin{array}{ccc} V_i \times_X X_j \times_X X_k & \xrightarrow{\text{pr}_{02}^* \varphi_{ik}} & X_i \times_X X_j \times_X V_k \\ & \searrow \text{pr}_{01}^* \varphi_{ij} & \nearrow \text{pr}_{12}^* \varphi_{jk} \\ & X_i \times_X V_j \times_X X_k & \end{array}$$

of algebraic spaces over $X_i \times_X X_j \times_X X_k$ commutes (with obvious notation).

- (2) A *morphism* $\psi : (V_i, \varphi_{ij}) \rightarrow (V'_i, \varphi'_{ij})$ of descent data is given by a family $\psi = (\psi_i)_{i \in I}$ of morphisms $\psi_i : V_i \rightarrow V'_i$ of algebraic spaces over X_i such

¹Unfortunately, we have chosen the “wrong” direction for our arrow here. In Definitions 21.1 and 21.3 we should have the opposite direction to what was done in Definition 3.1 by the general principle that “functions” and “spaces” are dual.

that all the diagrams

$$\begin{array}{ccc} V_i \times_X X_j & \xrightarrow{\varphi_{ij}} & X_i \times_X V_j \\ \psi_i \times \text{id} \downarrow & & \downarrow \text{id} \times \psi_j \\ V'_i \times_X X_j & \xrightarrow{\varphi'_{ij}} & X_i \times_X V'_j \end{array}$$

commute.

0ADJ **Remark 21.4.** Let S be a scheme. Let $\{X_i \rightarrow X\}_{i \in I}$ be a family of morphisms of algebraic spaces over S with fixed target X . Let (V_i, φ_{ij}) be a descent datum relative to $\{X_i \rightarrow X\}$. We may think of the isomorphisms φ_{ij} as isomorphisms

$$(X_i \times_X X_j) \times_{\text{pr}_0, X_i} V_i \longrightarrow (X_i \times_X X_j) \times_{\text{pr}_1, X_j} V_j$$

of algebraic spaces over $X_i \times_X X_j$. So loosely speaking one may think of φ_{ij} as an isomorphism $\text{pr}_0^* V_i \rightarrow \text{pr}_1^* V_j$ over $X_i \times_X X_j$. The cocycle condition then says that $\text{pr}_{02}^* \varphi_{ik} = \text{pr}_{12}^* \varphi_{jk} \circ \text{pr}_{01}^* \varphi_{ij}$. In this way it is very similar to the case of a descent datum on quasi-coherent sheaves.

The reason we will usually work with the version of a family consisting of a single morphism is the following lemma.

0ADK **Lemma 21.5.** *Let S be a scheme. Let $\{X_i \rightarrow X\}_{i \in I}$ be a family of morphisms of algebraic spaces over S with fixed target X . Set $Y = \coprod_{i \in I} X_i$. There is a canonical equivalence of categories*

$$\begin{array}{ccc} \text{category of descent data} & & \text{category of descent data} \\ \text{relative to the family } \{X_i \rightarrow X\}_{i \in I} & \longrightarrow & \text{relative to } Y/X \end{array}$$

which maps (V_i, φ_{ij}) to (V, φ) with $V = \coprod_{i \in I} V_i$ and $\varphi = \coprod \varphi_{ij}$.

Proof. Observe that $Y \times_X Y = \coprod_{i,j} X_i \times_X X_j$ and similarly for higher fibre products. Giving a morphism $V \rightarrow Y$ is exactly the same as giving a family $V_i \rightarrow X_i$. And giving a descent datum φ is exactly the same as giving a family φ_{ij} . \square

0ADL **Lemma 21.6.** *Pullback of descent data. Let S be a scheme.*

(1) *Let*

$$\begin{array}{ccc} Y' & \xrightarrow{\quad} & Y \\ a' \downarrow & \searrow f & \downarrow a \\ X' & \xrightarrow{h} & X \end{array}$$

be a commutative diagram of algebraic spaces over S . The construction

$$(V \rightarrow Y, \varphi) \longmapsto f^*(V \rightarrow Y, \varphi) = (V' \rightarrow Y', \varphi')$$

where $V' = Y' \times_Y V$ and where φ' is defined as the composition

$$\begin{aligned} V' \times_{X'} Y' & \longleftarrow (Y' \times_Y V) \times_{X'} Y' \longleftarrow (Y' \times_{X'} Y') \times_{Y \times_X Y} (V \times_X Y) \\ & \qquad \qquad \qquad \downarrow \text{id} \times \varphi \\ Y' \times_{X'} V' & \longleftarrow Y' \times_{X'} (Y' \times_Y V) \longleftarrow (Y' \times_X Y') \times_{Y \times_X Y} (Y \times_X V) \end{aligned}$$

defines a functor from the category of descent data relative to $Y \rightarrow X$ to the category of descent data relative to $Y' \rightarrow X'$.

- (2) Given two morphisms $f_i : Y' \rightarrow Y$, $i = 0, 1$ making the diagram commute the functors f_0^* and f_1^* are canonically isomorphic.

Proof. We omit the proof of (1), but we remark that the morphism φ' is the morphism $(f \times f)^*\varphi$ in the notation introduced in Remark 21.2. For (2) we indicate which morphism $f_0^*V \rightarrow f_1^*V$ gives the functorial isomorphism. Namely, since f_0 and f_1 both fit into the commutative diagram we see there is a unique morphism $r : Y' \rightarrow Y \times_X Y$ with $f_i = \text{pr}_i \circ r$. Then we take

$$\begin{aligned} f_0^*V &= Y' \times_{f_0, Y} V \\ &= Y' \times_{\text{pr}_0 \circ r, Y} V \\ &= Y' \times_{r, Y \times_X Y} (Y \times_X Y) \times_{\text{pr}_0, Y} V \\ &\xrightarrow{\varphi} Y' \times_{r, Y \times_X Y} (Y \times_X Y) \times_{\text{pr}_1, Y} V \\ &= Y' \times_{\text{pr}_1 \circ r, Y} V \\ &= Y' \times_{f_1, Y} V \\ &= f_1^*V \end{aligned}$$

We omit the verification that this works. \square

0ADM **Definition 21.7.** With $S, X, X', Y, Y', f, a, a', h$ as in Lemma 21.6 the functor

$$(V, \varphi) \mapsto f^*(V, \varphi)$$

constructed in that lemma is called the *pullback functor* on descent data.

0ADN **Lemma 21.8.** Let S be a scheme. Let $\mathcal{U}' = \{X'_i \rightarrow X'\}_{i \in I'}$ and $\mathcal{U} = \{X_j \rightarrow X\}_{j \in I}$ be families of morphisms with fixed target. Let $\alpha : I' \rightarrow I$, $g : X' \rightarrow X$ and $g_i : X'_i \rightarrow X_{\alpha(i)}$ be a morphism of families of maps with fixed target, see Sites, Definition 8.1.

- (1) Let (V_i, φ_{ij}) be a descent datum relative to the family \mathcal{U} . The system

$$(g_i^*V_{\alpha(i)}, (g_i \times g_j)^*\varphi_{\alpha(i)\alpha(j)})$$

(with notation as in Remark 21.4) is a descent datum relative to \mathcal{U}' .

- (2) This construction defines a functor between the category of descent data relative to \mathcal{U} and the category of descent data relative to \mathcal{U}' .
- (3) Given a second $\beta : I' \rightarrow I$, $h : X' \rightarrow X$ and $h'_i : X'_i \rightarrow X_{\beta(i)}$ morphism of families of maps with fixed target, then if $g = h$ the two resulting functors between descent data are canonically isomorphic.
- (4) These functors agree, via Lemma 21.5, with the pullback functors constructed in Lemma 21.6.

Proof. This follows from Lemma 21.6 via the correspondence of Lemma 21.5. \square

0ADP **Definition 21.9.** With $\mathcal{U}' = \{X'_i \rightarrow X'\}_{i \in I'}$, $\mathcal{U} = \{X_i \rightarrow X\}_{i \in I}$, $\alpha : I' \rightarrow I$, $g : X' \rightarrow X$, and $g_i : X'_i \rightarrow X_{\alpha(i)}$ as in Lemma 21.8 the functor

$$(V_i, \varphi_{ij}) \mapsto (g_i^*V_{\alpha(i)}, (g_i \times g_j)^*\varphi_{\alpha(i)\alpha(j)})$$

constructed in that lemma is called the *pullback functor* on descent data.

If \mathcal{U} and \mathcal{U}' have the same target X , and if \mathcal{U}' refines \mathcal{U} (see Sites, Definition 8.1) but no explicit pair (α, g_i) is given, then we can still talk about the pullback functor since we have seen in Lemma 21.8 that the choice of the pair does not matter (up to a canonical isomorphism).

0ADQ **Definition 21.10.** Let S be a scheme. Let $f : Y \rightarrow X$ be a morphism of algebraic spaces over S .

- (1) Given an algebraic space U over X we have the *trivial descent datum* of U relative to $\text{id} : X \rightarrow X$, namely the identity morphism on U .
- (2) By Lemma 21.6 we get a *canonical descent datum* on $Y \times_X U$ relative to $Y \rightarrow X$ by pulling back the trivial descent datum via f . We often denote $(Y \times_X U, \text{can})$ this descent datum.
- (3) A descent datum (V, φ) relative to Y/X is called *effective* if (V, φ) is isomorphic to the canonical descent datum $(Y \times_X U, \text{can})$ for some algebraic space U over X .

Thus being effective means there exists an algebraic space U over X and an isomorphism $\psi : V \rightarrow Y \times_X U$ over Y such that φ is equal to the composition

$$V \times_X Y \xrightarrow{\psi \times \text{id}_Y} Y \times_X U \times_S Y = Y \times_X Y \times_X U \xrightarrow{\text{id}_Y \times \psi^{-1}} Y \times_X U$$

There is a slight problem here which is that this definition (in spirit) conflicts with the definition given in Descent, Definition 31.10 in case Y and X are schemes. However, it will always be clear from context which version we mean.

0ADR **Definition 21.11.** Let S be a scheme. Let $\{X_i \rightarrow X\}$ be a family of morphisms of algebraic spaces over S with fixed target X .

- (1) Given an algebraic space U over X we have a *canonical descent datum* on the family of algebraic spaces $X_i \times_X U$ by pulling back the trivial descent datum for U relative to $\{\text{id} : S \rightarrow S\}$. We denote this descent datum $(X_i \times_X U, \text{can})$.
- (2) A descent datum (V_i, φ_{ij}) relative to $\{X_i \rightarrow X\}$ is called *effective* if there exists an algebraic space U over X such that (V_i, φ_{ij}) is isomorphic to $(X_i \times_X U, \text{can})$.

22. Descent data in terms of sheaves

0ADS This section is the analogue of Descent, Section 36. It is slightly different as algebraic spaces are already sheaves.

0ADT **Lemma 22.1.** *Let S be a scheme. Let $\{X_i \rightarrow X\}_{i \in I}$ be an fppf covering of algebraic spaces over S (Topologies on Spaces, Definition 7.1). There is an equivalence of categories*

$$\left\{ \begin{array}{l} \text{descent data } (V_i, \varphi_{ij}) \\ \text{relative to } \{X_i \rightarrow X\} \end{array} \right\} \leftrightarrow \left\{ \begin{array}{l} \text{sheaves } F \text{ on } (Sch/S)_{fppf} \text{ endowed} \\ \text{with a map } F \rightarrow X \text{ such that each} \\ X_i \times_X F \text{ is an algebraic space} \end{array} \right\}.$$

Moreover,

- (1) *the algebraic space $X_i \times_X F$ on the right hand side corresponds to V_i on the left hand side, and*
- (2) *the sheaf F is an algebraic space² if and only if the corresponding descent datum (X_i, φ_{ij}) is effective.*

²We will see later that this is always the case if I is not too large, see Bootstrap, Lemma 11.3.

Proof. Let us construct the functor from right to left. Let $F \rightarrow X$ be a map of sheaves on $(Sch/S)_{fppf}$ such that each $V_i = X_i \times_X F$ is an algebraic space. We have the projection $V_i \rightarrow X_i$. Then both $V_i \times_X X_j$ and $X_i \times_X V_j$ represent the sheaf $X_i \times_X F \times_X X_j$ and hence we obtain an isomorphism

$$\varphi_{ii'} : V_i \times_X X_j \rightarrow X_i \times_X V_j$$

It is straightforward to see that the maps φ_{ij} are morphisms over $X_i \times_X X_j$ and satisfy the cocycle condition. The functor from right to left is given by this construction $F \mapsto (V_i, \varphi_{ij})$.

Let us construct a functor from left to right. The isomorphisms φ_{ij} give isomorphisms

$$\varphi_{ij} : V_i \times_X X_j \longrightarrow X_i \times_X V_j$$

over $X_i \times X_j$. Set F equal to the coequalizer in the following diagram

$$\coprod_{i,i'} V_i \times_X X_j \begin{array}{c} \xrightarrow{\text{pr}_0} \\ \xrightarrow{\text{pr}_1 \circ \varphi_{ij}} \end{array} \coprod_i V_i \longrightarrow F$$

The cocycle condition guarantees that F comes with a map $F \rightarrow X$ and that $X_i \times_X F$ is isomorphic to V_i . The functor from left to right is given by this construction $(V_i, \varphi_{ij}) \mapsto F$.

We omit the verification that these constructions are mutually quasi-inverse functors. The final statements (1) and (2) follow from the constructions. \square

23. Other chapters

Preliminaries	(24) Hypercoverings
(1) Introduction	Schemes
(2) Conventions	(25) Schemes
(3) Set Theory	(26) Constructions of Schemes
(4) Categories	(27) Properties of Schemes
(5) Topology	(28) Morphisms of Schemes
(6) Sheaves on Spaces	(29) Cohomology of Schemes
(7) Sites and Sheaves	(30) Divisors
(8) Stacks	(31) Limits of Schemes
(9) Fields	(32) Varieties
(10) Commutative Algebra	(33) Topologies on Schemes
(11) Brauer Groups	(34) Descent
(12) Homological Algebra	(35) Derived Categories of Schemes
(13) Derived Categories	(36) More on Morphisms
(14) Simplicial Methods	(37) More on Flatness
(15) More on Algebra	(38) Groupoid Schemes
(16) Smoothing Ring Maps	(39) More on Groupoid Schemes
(17) Sheaves of Modules	(40) Étale Morphisms of Schemes
(18) Modules on Sites	Topics in Scheme Theory
(19) Injectives	(41) Chow Homology
(20) Cohomology of Sheaves	(42) Intersection Theory
(21) Cohomology on Sites	(43) Picard Schemes of Curves
(22) Differential Graded Algebra	(44) Adequate Modules
(23) Divided Power Algebra	

- | | |
|-------------------------------------|---------------------------------------|
| (45) Dualizing Complexes | (78) Restricted Power Series |
| (46) Duality for Schemes | (79) Resolution of Surfaces Revisited |
| (47) Discriminants and Differents | Deformation Theory |
| (48) Local Cohomology | (80) Formal Deformation Theory |
| (49) Algebraic Curves | (81) Deformation Theory |
| (50) Resolution of Surfaces | (82) The Cotangent Complex |
| (51) Semistable Reduction | (83) Deformation Problems |
| (52) Fundamental Groups of Schemes | Algebraic Stacks |
| (53) Étale Cohomology | (84) Algebraic Stacks |
| (54) Crystalline Cohomology | (85) Examples of Stacks |
| (55) Pro-étale Cohomology | (86) Sheaves on Algebraic Stacks |
| Algebraic Spaces | (87) Criteria for Representability |
| (56) Algebraic Spaces | (88) Artin's Axioms |
| (57) Properties of Algebraic Spaces | (89) Quot and Hilbert Spaces |
| (58) Morphisms of Algebraic Spaces | (90) Properties of Algebraic Stacks |
| (59) Decent Algebraic Spaces | (91) Morphisms of Algebraic Stacks |
| (60) Cohomology of Algebraic Spaces | (92) Limits of Algebraic Stacks |
| (61) Limits of Algebraic Spaces | (93) Cohomology of Algebraic Stacks |
| (62) Divisors on Algebraic Spaces | (94) Derived Categories of Stacks |
| (63) Algebraic Spaces over Fields | (95) Introducing Algebraic Stacks |
| (64) Topologies on Algebraic Spaces | (96) More on Morphisms of Stacks |
| (65) Descent and Algebraic Spaces | (97) The Geometry of Stacks |
| (66) Derived Categories of Spaces | Topics in Moduli Theory |
| (67) More on Morphisms of Spaces | (98) Moduli Stacks |
| (68) Flatness on Algebraic Spaces | (99) Moduli of Curves |
| (69) Groupoids in Algebraic Spaces | Miscellany |
| (70) More on Groupoids in Spaces | (100) Examples |
| (71) Bootstrap | (101) Exercises |
| (72) Pushouts of Algebraic Spaces | (102) Guide to Literature |
| Topics in Geometry | (103) Desirables |
| (73) Quotients of Groupoids | (104) Coding Style |
| (74) More on Cohomology of Spaces | (105) Obsolete |
| (75) Simplicial Spaces | (106) GNU Free Documentation License |
| (76) Duality for Spaces | (107) Auto Generated Index |
| (77) Formal Algebraic Spaces | |

References

- [Gro95a] Alexander Grothendieck, *Technique de descente et théorèmes d'existence en géométrie algébrique. I. Généralités. Descente par morphismes fidèlement plats*, Séminaire Bourbaki, Vol. 5, Soc. Math. France, Paris, 1995, pp. 299–327.
- [Gro95b] ———, *Technique de descente et théorèmes d'existence en géométrie algébrique. II. Le théorème d'existence en théorie formelle des modules*, Séminaire Bourbaki, Vol. 5, Soc. Math. France, Paris, 1995, pp. 369–390.
- [Gro95c] ———, *Technique de descente et théorèmes d'existence en géométrie algébrique. V. Les schémas de Picard: théorèmes d'existence*, Séminaire Bourbaki, Vol. 7, Soc. Math. France, Paris, 1995, pp. 143–161.
- [Gro95d] ———, *Technique de descente et théorèmes d'existence en géométrie algébrique. VI. Les schémas de Picard: propriétés générales*, Séminaire Bourbaki, Vol. 7, Soc. Math. France, Paris, 1995, pp. 221–243.

- [Gro95e] ———, *Techniques de construction et théorèmes d'existence en géométrie algébrique. III. Préschémas quotients*, Séminaire Bourbaki, Vol. 6, Soc. Math. France, Paris, 1995, pp. 99–118.
- [Gro95f] ———, *Techniques de construction et théorèmes d'existence en géométrie algébrique. IV. Les schémas de Hilbert*, Séminaire Bourbaki, Vol. 6, Soc. Math. France, Paris, 1995, pp. 249–276.