

ALGEBRAIC SPACES

025R

Contents

1. Introduction	1
2. General remarks	2
3. Representable morphisms of presheaves	2
4. Lists of useful properties of morphisms of schemes	4
5. Properties of representable morphisms of presheaves	5
6. Algebraic spaces	8
7. Fibre products of algebraic spaces	9
8. Glueing algebraic spaces	10
9. Presentations of algebraic spaces	12
10. Algebraic spaces and equivalence relations	12
11. Algebraic spaces, retrofitted	17
12. Immersions and Zariski coverings of algebraic spaces	19
13. Separation conditions on algebraic spaces	20
14. Examples of algebraic spaces	21
15. Change of big site	26
16. Change of base scheme	27
17. Other chapters	30
References	31

1. Introduction

025S Algebraic spaces were first introduced by Michael Artin, see [Art69b], [Art70], [Art73], [Art71b], [Art71a], [Art69a], [Art69c], and [Art74]. Some of the foundational material was developed jointly with Knutson, who produced the book [Knu71]. Artin defined (see [Art69c, Definition 1.3]) an algebraic space as a sheaf for the étale topology which is locally in the étale topology representable. In most of Artin's work the categories of schemes considered are schemes locally of finite type over a fixed excellent Noetherian base.

Our definition is slightly different. First of all we consider sheaves for the fppf topology. This is just a technical point and scarcely makes any difference. Second, we include the condition that the diagonal is representable.

After defining algebraic spaces we make some foundational observations. The main result in this chapter is that with our definitions an algebraic space is the same thing as an étale equivalence relation, see the discussion in Section 9 and Theorem 10.5. The analogue of this theorem in Artin's setting is [Art69c, Theorem 1.5], or [Knu71, Proposition II.1.7]. In other words, the sheaf defined by an étale equivalence relation

This is a chapter of the Stacks Project, version 74eb6f76, compiled on May 29, 2018.

has a representable diagonal. It follows that our definition agrees with Artin's original definition in a broad sense. It also means that one can give examples of algebraic spaces by simply writing down an étale equivalence relation.

In Section 13 we introduce various separation axioms on algebraic spaces that we have found in the literature. Finally in Section 14 we give some weird and not so weird examples of algebraic spaces.

2. General remarks

025T We work in a suitable big fppf site Sch_{fppf} as in Topologies, Definition 7.6. So, if not explicitly stated otherwise all schemes will be objects of Sch_{fppf} . In Section 15 we discuss what changes if you change the big fppf site.

We will always work relative to a base S contained in Sch_{fppf} . And we will then work with the big fppf site $(Sch/S)_{fppf}$, see Topologies, Definition 7.8. The absolute case can be recovered by taking $S = \text{Spec}(\mathbf{Z})$.

If U, T are schemes over S , then we denote $U(T)$ for the set of T -valued points over S . In a formula: $U(T) = \text{Mor}_S(T, U)$.

Note that any fpqc covering is a universal effective epimorphism, see Descent, Lemma 10.3. Hence the topology on Sch_{fppf} is weaker than the canonical topology and all representable presheaves are sheaves.

3. Representable morphisms of presheaves

025U Let S be a scheme contained in Sch_{fppf} . Let $F, G : (Sch/S)_{fppf}^{opp} \rightarrow \text{Sets}$. Let $a : F \rightarrow G$ be a representable transformation of functors, see Categories, Definition 8.2. This means that for every $U \in \text{Ob}((Sch/S)_{fppf})$ and any $\xi \in G(U)$ the fiber product $h_U \times_{\xi, G} F$ is representable. Choose a representing object V_ξ and an isomorphism $h_{V_\xi} \rightarrow h_U \times_G F$. By the Yoneda lemma, see Categories, Lemma 3.5, the projection $h_{V_\xi} \rightarrow h_U \times_G F \rightarrow h_U$ comes from a unique morphism of schemes $a_\xi : V_\xi \rightarrow U$. Suggestively we could represent this by the diagram

$$\begin{array}{ccccc} V_\xi & \xrightarrow{\sim} & h_{V_\xi} & \longrightarrow & F \\ a_\xi \downarrow & & \downarrow & & \downarrow a \\ U & \xrightarrow{\sim} & h_U & \xrightarrow{\xi} & G \end{array}$$

where the squiggly arrows represent the Yoneda embedding. Here are some lemmas about this notion that work in great generality.

02W9 **Lemma 3.1.** *Let S, X, Y be objects of Sch_{fppf} . Let $f : X \rightarrow Y$ be a morphism of schemes. Then*

$$h_f : h_X \longrightarrow h_Y$$

is a representable transformation of functors.

Proof. This is formal and relies only on the fact that the category $(Sch/S)_{fppf}$ has fibre products. \square

02WA **Lemma 3.2.** *Let S be a scheme contained in Sch_{fppf} . Let $F, G, H : (Sch/S)_{fppf}^{opp} \rightarrow \text{Sets}$. Let $a : F \rightarrow G, b : G \rightarrow H$ be representable transformations of functors. Then*

$$b \circ a : F \longrightarrow H$$

is a representable transformation of functors.

Proof. This is entirely formal and works in any category. \square

02WB **Lemma 3.3.** *Let S be a scheme contained in Sch_{fppf} . Let $F, G, H : (Sch/S)_{fppf}^{opp} \rightarrow Sets$. Let $a : F \rightarrow G$ be a representable transformation of functors. Let $b : H \rightarrow G$ be any transformation of functors. Consider the fibre product diagram*

$$\begin{array}{ccc} H \times_{b, G, a} F & \longrightarrow & F \\ a' \downarrow & & \downarrow a \\ H & \xrightarrow{b} & G \end{array}$$

Then the base change a' is a representable transformation of functors.

Proof. This is entirely formal and works in any category. \square

02WC **Lemma 3.4.** *Let S be a scheme contained in Sch_{fppf} . Let $F_i, G_i : (Sch/S)_{fppf}^{opp} \rightarrow Sets$, $i = 1, 2$. Let $a_i : F_i \rightarrow G_i$, $i = 1, 2$ be representable transformations of functors. Then*

$$a_1 \times a_2 : F_1 \times F_2 \longrightarrow G_1 \times G_2$$

is a representable transformation of functors.

Proof. Write $a_1 \times a_2$ as the composition $F_1 \times F_2 \rightarrow G_1 \times F_2 \rightarrow G_1 \times G_2$. The first arrow is the base change of a_1 by the map $G_1 \times F_2 \rightarrow G_1$, and the second arrow is the base change of a_2 by the map $G_1 \times G_2 \rightarrow G_2$. Hence this lemma is a formal consequence of Lemmas 3.2 and 3.3. \square

02WD **Lemma 3.5.** *Let S be a scheme contained in Sch_{fppf} . Let $F, G : (Sch/S)_{fppf}^{opp} \rightarrow Sets$. Let $a : F \rightarrow G$ be a representable transformation of functors. If G is a sheaf, then so is F .*

Proof. Let $\{\varphi_i : T_i \rightarrow T\}$ be a covering of the site $(Sch/S)_{fppf}$. Let $s_i \in F(T_i)$ which satisfy the sheaf condition. Then $\sigma_i = a(s_i) \in G(T_i)$ satisfy the sheaf condition also. Hence there exists a unique $\sigma \in G(T)$ such that $\sigma_i = \sigma|_{T_i}$. By assumption $F' = h_T \times_{\sigma, G, a} F$ is a representable presheaf and hence (see remarks in Section 2) a sheaf. Note that $(\varphi_i, s_i) \in F'(T_i)$ satisfy the sheaf condition also, and hence come from some unique $(id_T, s) \in F'(T)$. Clearly s is the section of F we are looking for. \square

05L9 **Lemma 3.6.** *Let S be a scheme contained in Sch_{fppf} . Let $F, G : (Sch/S)_{fppf}^{opp} \rightarrow Sets$. Let $a : F \rightarrow G$ be a representable transformation of functors. Then $\Delta_{F/G} : F \rightarrow F \times_G F$ is representable.*

Proof. Let $U \in \text{Ob}((Sch/S)_{fppf})$. Let $\xi = (\xi_1, \xi_2) \in (F \times_G F)(U)$. Set $\xi' = a(\xi_1) = a(\xi_2) \in G(U)$. By assumption there exist a scheme V and a morphism $V \rightarrow U$ representing the fibre product $h_U \times_{\xi', G} F$. In particular, the elements ξ_1, ξ_2 give morphisms $f_1, f_2 : U \rightarrow V$ over U . Because V represents the fibre product $h_U \times_{\xi', G} F$ and because $\xi' = a \circ \xi_1 = a \circ \xi_2$ we see that if $g : U' \rightarrow U$ is a morphism then

$$g^* \xi_1 = g^* \xi_2 \Leftrightarrow f_1 \circ g = f_2 \circ g.$$

In other words, we see that $h_U \times_{\xi, F \times_G F} F$ is represented by $V \times_{\Delta, V \times V, (f_1, f_2)} U$ which is a scheme. \square

4. Lists of useful properties of morphisms of schemes

02WE For ease of reference we list in the following remarks the properties of morphisms which possess some of the properties required of them in later results.

02WF **Remark 4.1.** Here is a list of properties/types of morphisms which are *stable under arbitrary base change*:

- (1) closed, open, and locally closed immersions, see Schemes, Lemma 18.2,
- (2) quasi-compact, see Schemes, Lemma 19.3,
- (3) universally closed, see Schemes, Definition 20.1,
- (4) (quasi-)separated, see Schemes, Lemma 21.13,
- (5) monomorphism, see Schemes, Lemma 23.5
- (6) surjective, see Morphisms, Lemma 9.4,
- (7) universally injective, see Morphisms, Lemma 10.2,
- (8) affine, see Morphisms, Lemma 11.8,
- (9) quasi-affine, see Morphisms, Lemma 12.5,
- (10) (locally) of finite type, see Morphisms, Lemma 14.4,
- (11) (locally) quasi-finite, see Morphisms, Lemma 19.13,
- (12) (locally) of finite presentation, see Morphisms, Lemma 20.4,
- (13) locally of finite type of relative dimension d , see Morphisms, Lemma 28.2,
- (14) universally open, see Morphisms, Definition 22.1,
- (15) flat, see Morphisms, Lemma 24.7,
- (16) syntomic, see Morphisms, Lemma 29.4,
- (17) smooth, see Morphisms, Lemma 32.5,
- (18) unramified (resp. G-unramified), see Morphisms, Lemma 33.5,
- (19) étale, see Morphisms, Lemma 34.4,
- (20) proper, see Morphisms, Lemma 39.5,
- (21) H-projective, see Morphisms, Lemma 41.8,
- (22) (locally) projective, see Morphisms, Lemma 41.9,
- (23) finite or integral, see Morphisms, Lemma 42.6,
- (24) finite locally free, see Morphisms, Lemma 45.4,
- (25) universally submersive, see Morphisms, Lemma 23.2,
- (26) universal homeomorphism, see Morphisms, Lemma 43.2.

Add more as needed.

02WG **Remark 4.2.** Of the properties of morphisms which are stable under base change (as listed in Remark 4.1) the following are also *stable under compositions*:

- (1) closed, open and locally closed immersions, see Schemes, Lemma 24.3,
- (2) quasi-compact, see Schemes, Lemma 19.4,
- (3) universally closed, see Morphisms, Lemma 39.4,
- (4) (quasi-)separated, see Schemes, Lemma 21.13,
- (5) monomorphism, see Schemes, Lemma 23.4,
- (6) surjective, see Morphisms, Lemma 9.2,
- (7) universally injective, see Morphisms, Lemma 10.5,
- (8) affine, see Morphisms, Lemma 11.7,
- (9) quasi-affine, see Morphisms, Lemma 12.4,
- (10) (locally) of finite type, see Morphisms, Lemma 14.3,
- (11) (locally) quasi-finite, see Morphisms, Lemma 19.12,
- (12) (locally) of finite presentation, see Morphisms, Lemma 20.3,
- (13) universally open, see Morphisms, Lemma 22.3,

- (14) flat, see Morphisms, Lemma 24.5,
- (15) syntomic, see Morphisms, Lemma 29.3,
- (16) smooth, see Morphisms, Lemma 32.4,
- (17) unramified (resp. G-unramified), see Morphisms, Lemma 33.4,
- (18) étale, see Morphisms, Lemma 34.3,
- (19) proper, see Morphisms, Lemma 39.4,
- (20) H-projective, see Morphisms, Lemma 41.7,
- (21) finite or integral, see Morphisms, Lemma 42.5,
- (22) finite locally free, see Morphisms, Lemma 45.3,
- (23) universally submersive, see Morphisms, Lemma 23.3,
- (24) universal homeomorphism, see Morphisms, Lemma 43.3.

Add more as needed.

02WH **Remark 4.3.** Of the properties mentioned which are stable under base change (as listed in Remark 4.1) the following are also *fqc local on the base* (and a fortiori *fppf local on the base*):

- (1) for immersions we have this for
 - (a) closed immersions, see Descent, Lemma 20.19,
 - (b) open immersions, see Descent, Lemma 20.16, and
 - (c) quasi-compact immersions, see Descent, Lemma 20.21,
- (2) quasi-compact, see Descent, Lemma 20.1,
- (3) universally closed, see Descent, Lemma 20.3,
- (4) (quasi-)separated, see Descent, Lemmas 20.2, and 20.6,
- (5) monomorphism, see Descent, Lemma 20.31,
- (6) surjective, see Descent, Lemma 20.7,
- (7) universally injective, see Descent, Lemma 20.8,
- (8) affine, see Descent, Lemma 20.18,
- (9) quasi-affine, see Descent, Lemma 20.20,
- (10) (locally) of finite type, see Descent, Lemmas 20.10, and 20.12,
- (11) (locally) quasi-finite, see Descent, Lemma 20.24,
- (12) (locally) of finite presentation, see Descent, Lemmas 20.11, and 20.13,
- (13) locally of finite type of relative dimension d , see Descent, Lemma 20.25,
- (14) universally open, see Descent, Lemma 20.4,
- (15) flat, see Descent, Lemma 20.15,
- (16) syntomic, see Descent, Lemma 20.26,
- (17) smooth, see Descent, Lemma 20.27,
- (18) unramified (resp. G-unramified), see Descent, Lemma 20.28,
- (19) étale, see Descent, Lemma 20.29,
- (20) proper, see Descent, Lemma 20.14,
- (21) finite or integral, see Descent, Lemma 20.23,
- (22) finite locally free, see Descent, Lemma 20.30,
- (23) universally submersive, see Descent, Lemma 20.5,
- (24) universal homeomorphism, see Descent, Lemma 20.9.

Note that the property of being an “immersion” may not be *fqc local on the base*, but in Descent, Lemma 21.1 we proved that it is *fppf local on the base*.

5. Properties of representable morphisms of presheaves

02WI Here is the definition that makes this work.

025V **Definition 5.1.** With S , and $a : F \rightarrow G$ representable as above. Let \mathcal{P} be a property of morphisms of schemes which

- (1) is preserved under any base change, see Schemes, Definition 18.3, and
- (2) is fppf local on the base, see Descent, Definition 19.1.

In this case we say that a has *property* \mathcal{P} if for every $U \in \text{Ob}((\text{Sch}/S)_{\text{fppf}})$ and any $\xi \in G(U)$ the resulting morphism of schemes $V_\xi \rightarrow U$ has property \mathcal{P} .

It is important to note that we will only use this definition for properties of morphisms that are stable under base change, and local in the fppf topology on the base. This is not because the definition doesn't make sense otherwise; rather it is because we may want to give a different definition which is better suited to the property we have in mind.

02YN **Remark 5.2.** Consider the property \mathcal{P} =“surjective”. In this case there could be some ambiguity if we say “let $F \rightarrow G$ be a surjective map”. Namely, we could mean the notion defined in Definition 5.1 above, or we could mean a surjective map of presheaves, see Sites, Definition 3.1, or, if both F and G are sheaves, we could mean a surjective map of sheaves, see Sites, Definition 11.1, If not mentioned otherwise when discussing morphisms of algebraic spaces we will always mean the first. See Lemma 5.9 for a case where surjectivity implies surjectivity as a map of sheaves.

Here is a sanity check.

02WJ **Lemma 5.3.** Let S, X, Y be objects of Sch_{fppf} . Let $f : X \rightarrow Y$ be a morphism of schemes. Let \mathcal{P} be as in Definition 5.1. Then $h_X \rightarrow h_Y$ has property \mathcal{P} if and only if f has property \mathcal{P} .

Proof. Note that the lemma makes sense by Lemma 3.1. Proof omitted. \square

02WK **Lemma 5.4.** Let S be a scheme contained in Sch_{fppf} . Let $F, G, H : (\text{Sch}/S)_{\text{fppf}}^{\text{opp}} \rightarrow \text{Sets}$. Let \mathcal{P} be a property as in Definition 5.1 which is stable under composition. Let $a : F \rightarrow G, b : G \rightarrow H$ be representable transformations of functors. If a and b have property \mathcal{P} so does $b \circ a : F \rightarrow H$.

Proof. Note that the lemma makes sense by Lemma 3.2. Proof omitted. \square

02WL **Lemma 5.5.** Let S be a scheme contained in Sch_{fppf} . Let $F, G, H : (\text{Sch}/S)_{\text{fppf}}^{\text{opp}} \rightarrow \text{Sets}$. Let \mathcal{P} be a property as in Definition 5.1. Let $a : F \rightarrow G$ be a representable transformation of functors. Let $b : H \rightarrow G$ be any transformation of functors. Consider the fibre product diagram

$$\begin{array}{ccc} H \times_{b,G,a} F & \longrightarrow & F \\ a' \downarrow & & \downarrow a \\ H & \xrightarrow{b} & G \end{array}$$

If a has property \mathcal{P} then also the base change a' has property \mathcal{P} .

Proof. Note that the lemma makes sense by Lemma 3.3. Proof omitted. \square

03KD **Lemma 5.6.** Let S be a scheme contained in Sch_{fppf} . Let $F, G, H : (\text{Sch}/S)_{\text{fppf}}^{\text{opp}} \rightarrow \text{Sets}$. Let \mathcal{P} be a property as in Definition 5.1. Let $a : F \rightarrow G$ be a representable

transformations of functors. Let $b : H \rightarrow G$ be any transformation of functors. Consider the fibre product diagram

$$\begin{array}{ccc} H \times_{b,G,a} F & \longrightarrow & F \\ a' \downarrow & & \downarrow a \\ H & \xrightarrow{b} & G \end{array}$$

Assume that b induces a surjective map of fppf sheaves $H^\# \rightarrow G^\#$. In this case, if a' has property \mathcal{P} , then also a has property \mathcal{P} .

Proof. First we remark that by Lemma 3.3 the transformation a' is representable. Let $U \in \text{Ob}((\text{Sch}/S)_{\text{fppf}})$, and let $\xi \in G(U)$. By assumption there exists an fppf covering $\{U_i \rightarrow U\}_{i \in I}$ and elements $\xi_i \in H(U_i)$ mapping to $\xi|_{U_i}$ via b . From general category theory it follows that for each i we have a fibre product diagram

$$\begin{array}{ccc} U_i \times_{\xi_i, H, a'} (H \times_{b, G, a} F) & \longrightarrow & U \times_{\xi, G, a} F \\ \downarrow & & \downarrow \\ U_i & \longrightarrow & U \end{array}$$

By assumption the left vertical arrow is a morphism of schemes which has property \mathcal{P} . Since \mathcal{P} is local in the fppf topology this implies that also the right vertical arrow has property \mathcal{P} as desired. \square

02WM **Lemma 5.7.** Let S be a scheme contained in Sch_{fppf} . Let $F_i, G_i : (\text{Sch}/S)_{\text{fppf}}^{\text{opp}} \rightarrow \text{Sets}$, $i = 1, 2$. Let $a_i : F_i \rightarrow G_i$, $i = 1, 2$ be representable transformations of functors. Let \mathcal{P} be a property as in Definition 5.1 which is stable under composition. If a_1 and a_2 have property \mathcal{P} so does $a_1 \times a_2 : F_1 \times F_2 \rightarrow G_1 \times G_2$.

Proof. Note that the lemma makes sense by Lemma 3.4. Proof omitted. \square

02YO **Lemma 5.8.** Let S be a scheme contained in Sch_{fppf} . Let $F, G : (\text{Sch}/S)_{\text{fppf}}^{\text{opp}} \rightarrow \text{Sets}$. Let $a : F \rightarrow G$ be a representable transformation of functors. Let $\mathcal{P}, \mathcal{P}'$ be properties as in Definition 5.1. Suppose that for any morphism of schemes $f : X \rightarrow Y$ we have $\mathcal{P}(f) \Rightarrow \mathcal{P}'(f)$. If a has property \mathcal{P} then a has property \mathcal{P}' .

Proof. Formal. \square

05VM **Lemma 5.9.** Let S be a scheme. Let $F, G : (\text{Sch}/S)_{\text{fppf}}^{\text{opp}} \rightarrow \text{Sets}$ be sheaves. Let $a : F \rightarrow G$ be representable, flat, locally of finite presentation, and surjective. Then $a : F \rightarrow G$ is surjective as a map of sheaves.

Proof. Let T be a scheme over S and let $g : T \rightarrow G$ be a T -valued point of G . By assumption $T' = F \times_G T$ is (representable by) a scheme and the morphism $T' \rightarrow T$ is a flat, locally of finite presentation, and surjective. Hence $\{T' \rightarrow T\}$ is an fppf covering such that $g|_{T'} \in G(T')$ comes from an element of $F(T')$, namely the map $T' \rightarrow F$. This proves the map is surjective as a map of sheaves, see Sites, Definition 11.1. \square

Here is a characterization of those functors for which the diagonal is representable.

025W **Lemma 5.10.** Let S be a scheme contained in Sch_{fppf} . Let F be a presheaf of sets on $(\text{Sch}/S)_{\text{fppf}}$. The following are equivalent:

- (1) the diagonal $F \rightarrow F \times F$ is representable,

- (2) for $U \in \text{Ob}((\text{Sch}/S)_{fppf})$ and any $a \in F(U)$ the map $a : h_U \rightarrow F$ is representable,
- (3) for every pair $U, V \in \text{Ob}((\text{Sch}/S)_{fppf})$ and any $a \in F(U), b \in F(V)$ the fibre product $h_U \times_{a,F,b} h_V$ is representable.

Proof. This is completely formal, see Categories, Lemma 8.4. It depends only on the fact that the category $(\text{Sch}/S)_{fppf}$ has products of pairs of objects and fibre products, see Topologies, Lemma 7.10. \square

In the situation of the lemma, for any morphism $\xi : h_U \rightarrow F$ as in the lemma, it makes sense to say that ξ has property \mathcal{P} , for any property as in Definition 5.1. In particular this holds for $\mathcal{P} = \text{“surjective”}$ and $\mathcal{P} = \text{“étale”}$, see Remark 4.3 above. We will use this remark in the definition of algebraic spaces below.

0CB7 **Lemma 5.11.** *Let S be a scheme contained in Sch_{fppf} . Let F be a presheaf of sets on $(\text{Sch}/S)_{fppf}$. Let \mathcal{P} be a property as in Definition 5.1. If for every $U, V \in \text{Ob}((\text{Sch}/S)_{fppf})$ and $a \in F(U), b \in F(V)$ we have*

- (1) $h_U \times_{a,F,b} h_V$ is representable, say by the scheme W , and
- (2) the morphism $W \rightarrow U \times_S V$ corresponding to $h_U \times_{a,F,b} h_V \rightarrow h_U \times h_V$ has property \mathcal{P} ,

then $\Delta : F \rightarrow F \times F$ is representable and has property \mathcal{P} .

Proof. Observe that Δ is representable by Lemma 5.10. We can formulate condition (2) as saying that the transformation $h_U \times_{a,F,b} h_V \rightarrow h_{U \times_S V}$ has property \mathcal{P} , see Lemma 5.3. Consider $T \in \text{Ob}((\text{Sch}/S)_{fppf})$ and $(a, b) \in (F \times F)(T)$. Observe that we have the commutative diagram

$$\begin{array}{ccc}
 F \times_{\Delta, F \times F, (a,b)} h_T & \longrightarrow & h_T \\
 \downarrow & & \downarrow \Delta_{T/S} \\
 h_T \times_{a,F,b} h_T & \longrightarrow & h_{T \times_S T} \\
 \downarrow & & \downarrow (a,b) \\
 F & \xrightarrow{\Delta} & F \times F
 \end{array}$$

both of whose squares are cartesian. In this way we see that the morphism $F \times_{F \times F} h_T \rightarrow h_T$ is the base change of a morphism having property \mathcal{P} by $\Delta_{T/S}$. Since \mathcal{P} is preserved under base change this finishes the proof. \square

6. Algebraic spaces

025X Here is the definition.

025Y **Definition 6.1.** Let S be a scheme contained in Sch_{fppf} . An algebraic space over S is a presheaf

$$F : (\text{Sch}/S)_{fppf}^{opp} \longrightarrow \text{Sets}$$

with the following properties

- (1) The presheaf F is a sheaf.
- (2) The diagonal morphism $F \rightarrow F \times F$ is representable.
- (3) There exists a scheme $U \in \text{Ob}((\text{Sch}/S)_{fppf})$ and a map $h_U \rightarrow F$ which is surjective, and étale.

There are two differences with the “usual” definition, for example the definition in Knutson’s book [Knu71].

The first is that we require F to be a sheaf in the fppf topology. One reason for doing this is that many natural examples of algebraic spaces satisfy the sheaf condition for the fppf coverings (and even for fpqc coverings). Also, one of the reasons that algebraic spaces have been so useful is via Michael Artin’s results on algebraic spaces. Built into his method is a condition which guarantees the result is locally of finite presentation over S . Combined it somehow seems to us that the fppf topology is the natural topology to work with. In the end the category of algebraic spaces ends up being the same. See Bootstrap, Section 12.

The second is that we only require the diagonal map for F to be representable, whereas in [Knu71] it is required that it also be quasi-compact. If $F = h_U$ for some scheme U over S this corresponds to the condition that U be quasi-separated. Our point of view is to try to prove a certain number of the results that follow only assuming that the diagonal of F be representable, and simply add an additional hypothesis wherever this is necessary. In any case it has the pleasing consequence that the following lemma is true.

025Z **Lemma 6.2.** *A scheme is an algebraic space. More precisely, given a scheme $T \in \text{Ob}((\text{Sch}/S)_{\text{fppf}})$ the representable functor h_T is an algebraic space.*

Proof. The functor h_T is a sheaf by our remarks in Section 2. The diagonal $h_T \rightarrow h_T \times h_T = h_{T \times T}$ is representable because $(\text{Sch}/S)_{\text{fppf}}$ has fibre products. The identity map $h_T \rightarrow h_T$ is surjective étale. \square

0260 **Definition 6.3.** *Let F, F' be algebraic spaces over S . A morphism $f : F \rightarrow F'$ of algebraic spaces over S is a transformation of functors from F to F' .*

The category of algebraic spaces over S contains the category $(\text{Sch}/S)_{\text{fppf}}$ as a full subcategory via the Yoneda embedding $T/S \mapsto h_T$. From now on we no longer distinguish between a scheme T/S and the algebraic space it represents. Thus when we say “Let $f : T \rightarrow F$ be a morphism from the scheme T to the algebraic space F ”, we mean that $T \in \text{Ob}((\text{Sch}/S)_{\text{fppf}})$, that F is an algebraic space over S , and that $f : h_T \rightarrow F$ is a morphism of algebraic spaces over S .

7. Fibre products of algebraic spaces

04T8 The category of algebraic spaces over S has both products and fibre products.

02X0 **Lemma 7.1.** *Let S be a scheme contained in Sch_{fppf} . Let F, G be algebraic spaces over S . Then $F \times G$ is an algebraic space, and is a product in the category of algebraic spaces over S .*

Proof. It is clear that $H = F \times G$ is a sheaf. The diagonal of H is simply the product of the diagonals of F and G . Hence it is representable by Lemma 3.4. Finally, if $U \rightarrow F$ and $V \rightarrow G$ are surjective étale morphisms, with $U, V \in \text{Ob}((\text{Sch}/S)_{\text{fppf}})$, then $U \times V \rightarrow F \times G$ is surjective étale by Lemma 5.7. \square

04T9 **Lemma 7.2.** *Let S be a scheme contained in Sch_{fppf} . Let H be a sheaf on $(\text{Sch}/S)_{\text{fppf}}$ whose diagonal is representable. Let F, G be algebraic spaces over S . Let $F \rightarrow H, G \rightarrow H$ be maps of sheaves. Then $F \times_H G$ is an algebraic space.*

Proof. We check the 3 conditions of Definition 6.1. A fibre product of sheaves is a sheaf, hence $F \times_H G$ is a sheaf. The diagonal of $F \times_H G$ is the left vertical arrow in

$$\begin{array}{ccc} F \times_H G & \longrightarrow & F \times G \\ \Delta \downarrow & & \downarrow \Delta_F \times \Delta_G \\ (F \times F) \times_{(H \times H)} (G \times G) & \longrightarrow & (F \times F) \times (G \times G) \end{array}$$

which is cartesian. Hence Δ is representable as the base change of the morphism on the right which is representable, see Lemmas 3.4 and 3.3. Finally, let $U, V \in \text{Ob}((Sch/S)_{fppf})$ and $a : U \rightarrow F, b : V \rightarrow G$ be surjective and étale. As Δ_H is representable, we see that $U \times_H V$ is a scheme. The morphism

$$U \times_H V \longrightarrow F \times_H G$$

is surjective and étale as a composition of the base changes $U \times_H V \rightarrow U \times_H G$ and $U \times_H G \rightarrow F \times_H G$ of the étale surjective morphisms $U \rightarrow F$ and $V \rightarrow G$, see Lemmas 3.2 and 3.3. This proves the last condition of Definition 6.1 holds and we conclude that $F \times_H G$ is an algebraic space. \square

02X2 **Lemma 7.3.** *Let S be a scheme contained in Sch_{fppf} . Let $F \rightarrow H, G \rightarrow H$ be morphisms of algebraic spaces over S . Then $F \times_H G$ is an algebraic space, and is a fibre product in the category of algebraic spaces over S .*

Proof. It follows from the stronger Lemma 7.2 that $F \times_H G$ is an algebraic space. It is clear that $F \times_H G$ is a fibre product in the category of algebraic spaces over S since that is a full subcategory of the category of (pre)sheaves of sets on $(Sch/S)_{fppf}$. \square

8. Glueing algebraic spaces

02WN In this section we really start abusing notation and not distinguish between schemes and the spaces they represent.

02WO **Lemma 8.1.** *Let $S \in \text{Ob}(Sch_{fppf})$. Let $U \in \text{Ob}((Sch/S)_{fppf})$. Given a set I and sheaves F_i on $\text{Ob}((Sch/S)_{fppf})$, if $U \cong \coprod_{i \in I} F_i$ as sheaves, then each F_i is representable by an open and closed subscheme U_i and $U \cong \coprod U_i$ as schemes.*

Proof. By assumption this means there exists an fppf covering $\{U_j \rightarrow U\}_{j \in J}$ such that each $U_j \rightarrow U$ factors through $F_{i(j)}$ for some $i(j) \in I$. Denote $V_j = \text{Im}(U_j \rightarrow U)$. This is an open of U by Morphisms, Lemma 24.9, and $\{U_j \rightarrow V_j\}$ is an fppf covering. Hence it follows that $V_j \rightarrow U$ factors through $F_{i(j)}$ since $F_{i(j)}$ is a subsheaf. It follows from $F_i \cap F_{i'} = \emptyset, i \neq i'$ that $V_j \cap V_{j'} = \emptyset$ unless $i(j) = i(j')$. Hence we can take $U_i = \bigcup_{j, i(j)=i} V_j$ and everything is clear. \square

02WP **Lemma 8.2.** *Let $S \in \text{Ob}(Sch_{fppf})$. Let F be an algebraic space over S . Given a set I and sheaves F_i on $\text{Ob}((Sch/S)_{fppf})$, if $F \cong \coprod_{i \in I} F_i$ as sheaves, then each F_i is an algebraic space over S .*

Proof. It follows directly from the representability of $F \rightarrow F \times F$ that each diagonal morphism $F_i \rightarrow F_i \times F_i$ is representable. Choose a scheme U in $(Sch/S)_{fppf}$ and a surjective étale morphism $U \rightarrow \coprod F_i$ (this exist by hypothesis). By considering the inverse image of F_i we get a decomposition of U (as a sheaf) into a coproduct of sheaves. By Lemma 8.1 we get correspondingly $U \cong \coprod U_i$. Then it follows

easily that $U_i \rightarrow F_i$ is surjective and étale (from the corresponding property of $U \rightarrow F$). \square

The condition on the size of I and the F_i in the following lemma may be ignored by those not worried about set theoretic questions.

02WQ **Lemma 8.3.** *Let $S \in \text{Ob}(\text{Sch}_{fppf})$. Suppose given a set I and algebraic spaces F_i , $i \in I$. Then $F = \coprod_{i \in I} F_i$ is an algebraic space provided I , and the F_i are not too “large”: for example if we can choose surjective étale morphisms $U_i \rightarrow F_i$ such that $\coprod_{i \in I} U_i$ is isomorphic to an object of $(\text{Sch}/S)_{fppf}$, then F is an algebraic space.*

Proof. By construction F is a sheaf. We omit the verification that the diagonal morphism of F is representable. Finally, if U is an object of $(\text{Sch}/S)_{fppf}$ isomorphic to $\coprod_{i \in I} U_i$ then it is straightforward to verify that the resulting map $U \rightarrow \coprod F_i$ is surjective and étale. \square

Here is the analogue of Schemes, Lemma 15.4.

02WR **Lemma 8.4.** *Let $S \in \text{Ob}(\text{Sch}_{fppf})$. Let F be a presheaf of sets on $(\text{Sch}/S)_{fppf}$. Assume*

- (1) F is a sheaf,
- (2) there exists an index set I and subfunctors $F_i \subset F$ such that
 - (a) each F_i is an algebraic space,
 - (b) each $F_i \rightarrow F$ is a representable,
 - (c) each $F_i \rightarrow F$ is an open immersion (see Definition 5.1),
 - (d) the map $\coprod F_i \rightarrow F$ is surjective as a map of sheaves, and
 - (e) $\coprod F_i$ is an algebraic space (set theoretic condition, see Lemma 8.3).

Then F is an algebraic space.

Proof. Let T be an object of $(\text{Sch}/S)_{fppf}$. Let $T \rightarrow F$ be a morphism. By assumption (2)(b) and (2)(c) the fibre product $F_i \times_F T$ is representable by an open subscheme $V_i \subset T$. It follows that $(\coprod F_i) \times_F T$ is represented by the scheme $\coprod V_i$ over T . By assumption (2)(d) there exists an fppf covering $\{T_j \rightarrow T\}_{j \in J}$ such that $T_j \rightarrow T \rightarrow F$ factors through F_i , $i = i(j)$. Hence $T_j \rightarrow T$ factors through the open subscheme $V_{i(j)} \subset T$. Since $\{T_j \rightarrow T\}$ is jointly surjective, it follows that $T = \bigcup V_i$ is an open covering. In particular, the transformation of functors $\coprod F_i \rightarrow F$ is representable and surjective in the sense of Definition 5.1 (see Remark 5.2 for a discussion).

Next, let $T' \rightarrow F$ be a second morphism from an object in $(\text{Sch}/S)_{fppf}$. Write as above $T' = \bigcup V'_i$ with $V'_i = T' \times_F F_i$. To show that the diagonal $F \rightarrow F \times F$ is representable we have to show that $G = T \times_F T'$ is representable, see Lemma 5.10. Consider the subfunctors $G_i = G \times_F F_i$. Note that $G_i = V_i \times_{F_i} V'_i$, and hence is representable as F_i is an algebraic space. By the above the G_i form a Zariski covering of F . Hence by Schemes, Lemma 15.4 we see G is representable.

Choose a scheme $U \in \text{Ob}((\text{Sch}/S)_{fppf})$ and a surjective étale morphism $U \rightarrow \coprod F_i$ (this exist by hypothesis). We may write $U = \coprod U_i$ with U_i the inverse image of F_i , see Lemma 8.1. We claim that $U \rightarrow F$ is surjective and étale. Surjectivity follows as $\coprod F_i \rightarrow F$ is surjective (see first paragraph of the proof) by applying Lemma 5.4. Consider the fibre product $U \times_F T$ where $T \rightarrow F$ is as above. We have to show that $U \times_F T \rightarrow T$ is étale. Since $U \times_F T = \coprod U_i \times_F T$ it suffices to show each $U_i \times_F T \rightarrow T$ is étale. Since $U_i \times_F T = U_i \times_{F_i} V_i$ this follows from the fact that

$U_i \rightarrow F_i$ is étale and $V_i \rightarrow T$ is an open immersion (and Morphisms, Lemmas 34.9 and 34.3). \square

9. Presentations of algebraic spaces

0261 Given an algebraic space we can find a “presentation” of it.

0262 **Lemma 9.1.** *Let F be an algebraic space over S . Let $f : U \rightarrow F$ be a surjective étale morphism from a scheme to F . Set $R = U \times_F U$. Then*

- (1) $j : R \rightarrow U \times_S U$ defines an equivalence relation on U over S (see Groupoids, Definition 3.1).
- (2) the morphisms $s, t : R \rightarrow U$ are étale, and
- (3) the diagram

$$R \begin{array}{c} \xrightarrow{\quad} \\ \xrightarrow{\quad} \end{array} U \longrightarrow F$$

is a coequalizer diagram in $Sh((Sch/S)_{fppf})$.

Proof. Let T/S be an object of $(Sch/S)_{fppf}$. Then $R(T) = \{(a, b) \in U(T) \times U(T) \mid f \circ a = f \circ b\}$ which clearly defines an equivalence relation on $U(T)$. The morphisms $s, t : R \rightarrow U$ are étale because the morphism $U \rightarrow F$ is étale.

To prove (3) we first show that $U \rightarrow F$ is a surjection of sheaves, see Sites, Definition 11.1. Let $\xi \in F(T)$ with T as above. Let $V = T \times_{\xi, F, f} U$. By assumption V is a scheme and $V \rightarrow T$ is surjective étale. Hence $\{V \rightarrow T\}$ is a covering for the fppf topology. Since $\xi|_V$ factors through U by construction we conclude $U \rightarrow F$ is surjective. Surjectivity implies that F is the coequalizer of the diagram by Sites, Lemma 11.3. \square

This lemma suggests the following definitions.

02WS **Definition 9.2.** Let S be a scheme. Let U be a scheme over S . An *étale equivalence relation* on U over S is an equivalence relation $j : R \rightarrow U \times_S U$ such that $s, t : R \rightarrow U$ are étale morphisms of schemes.

0263 **Definition 9.3.** Let F be an algebraic space over S . A *presentation* of F is given by a scheme U over S and an étale equivalence relation R on U over S , and a surjective étale morphism $U \rightarrow F$ such that $R = U \times_F U$.

Equivalently we could ask for the existence of an isomorphism

$$U/R \cong F$$

where the quotient U/R is as defined in Groupoids, Section 20. To construct algebraic spaces we will study the converse question, namely, for which equivalence relations the quotient sheaf U/R is an algebraic space. It will finally turn out this is always the case if R is an étale equivalence relation on U over S , see Theorem 10.5.

10. Algebraic spaces and equivalence relations

0264 Suppose given a scheme U over S and an étale equivalence relation R on U over S . We would like to show this defines an algebraic space. We will produce a series of lemmas that prove the quotient sheaf U/R (see Groupoids, Definition 20.1) has all the properties required of it in Definition 6.1.

02WT **Lemma 10.1.** *Let S be a scheme. Let U be a scheme over S . Let $j = (s, t) : R \rightarrow U \times_S U$ be an étale equivalence relation on U over S . Let $U' \rightarrow U$ be an étale morphism. Let R' be the restriction of R to U' , see Groupoids, Definition 3.3. Then $j' : R' \rightarrow U' \times_S U'$ is an étale equivalence relation also.*

Proof. It is clear from the description of s', t' in Groupoids, Lemma 18.1 that $s', t' : R' \rightarrow U'$ are étale as compositions of base changes of étale morphisms (see Morphisms, Lemma 34.4 and 34.3). \square

We will often use the following lemma to find open subspaces of algebraic spaces. A slight improvement (with more general hypotheses) of this lemma is Bootstrap, Lemma 7.1.

02WU **Lemma 10.2.** *Let S be a scheme. Let U be a scheme over S . Let $j = (s, t) : R \rightarrow U \times_S U$ be a pre-relation. Let $g : U' \rightarrow U$ be a morphism. Assume*

- (1) j is an equivalence relation,
- (2) $s, t : R \rightarrow U$ are surjective, flat and locally of finite presentation,
- (3) g is flat and locally of finite presentation.

Let $R' = R|_{U'}$ be the restriction of R to U' . Then $U'/R' \rightarrow U/R$ is representable, and is an open immersion.

Proof. By Groupoids, Lemma 3.2 the morphism $j' = (t', s') : R' \rightarrow U' \times_S U'$ defines an equivalence relation. Since g is flat and locally of finite presentation we see that g is universally open as well (Morphisms, Lemma 24.9). For the same reason s, t are universally open as well. Let $W^1 = g(U') \subset U$, and let $W = t(s^{-1}(W^1))$. Then W^1 and W are open in U . Moreover, as j is an equivalence relation we have $t(s^{-1}(W)) = W$ (see Groupoids, Lemma 19.2 for example).

By Groupoids, Lemma 20.5 the map of sheaves $F' = U'/R' \rightarrow F = U/R$ is injective. Let $a : T \rightarrow F$ be a morphism from a scheme into U/R . We have to show that $T \times_F F'$ is representable by an open subscheme of T .

The morphism a is given by the following data: an fppf covering $\{\varphi_j : T_j \rightarrow T\}_{j \in J}$ of T and morphisms $a_j : T_j \rightarrow U$ such that the maps

$$a_j \times a_{j'} : T_j \times_T T_{j'} \longrightarrow U \times_S U$$

factor through $j : R \rightarrow U \times_S U$ via some (unique) maps $r_{jj'} : T_j \times_T T_{j'} \rightarrow R$. The system (a_j) corresponds to a in the sense that the diagrams

$$\begin{array}{ccc} T_j & \xrightarrow{a_j} & U \\ \downarrow & & \downarrow \\ T & \xrightarrow{a} & F \end{array}$$

commute.

Consider the open subsets $W_j = a_j^{-1}(W) \subset T_j$. Since $t(s^{-1}(W)) = W$ we see that

$$W_j \times_T T_{j'} = r_{jj'}^{-1}(t^{-1}(W)) = r_{jj'}^{-1}(s^{-1}(W)) = T_j \times_T W_{j'}.$$

By Descent, Lemma 10.2 this means there exists an open $W_T \subset T$ such that $\varphi_j^{-1}(W_T) = W_j$ for all $j \in J$. We claim that $W_T \rightarrow T$ represents $T \times_F F' \rightarrow T$.

First, let us show that $W_T \rightarrow T \rightarrow F$ is an element of $F'(W_T)$. Since $\{W_j \rightarrow W_T\}_{j \in J}$ is an fppf covering of W_T , it is enough to show that each $W_j \rightarrow U \rightarrow F$

is an element of $F'(W_j)$ (as F' is a sheaf for the fppf topology). Consider the commutative diagram

$$\begin{array}{ccccc}
 W'_j & \xrightarrow{\quad} & U' & & \\
 \downarrow & \searrow & \downarrow g & & \\
 & & s^{-1}(W^1) & \xrightarrow{s} & W^1 \\
 & & \downarrow t & & \downarrow \\
 W_j & \xrightarrow{a_j|_{W_j}} & W & \longrightarrow & F
 \end{array}$$

where $W'_j = W_j \times_W s^{-1}(W^1) \times_{W^1} U'$. Since t and g are surjective, flat and locally of finite presentation, so is $W'_j \rightarrow W_j$. Hence the restriction of the element $W_j \rightarrow U \rightarrow F$ to W'_j is an element of F' as desired.

Suppose that $f : T' \rightarrow T$ is a morphism of schemes such that $a|_{T'} \in F'(T')$. We have to show that f factors through the open W_T . Since $\{T' \times_T T_j \rightarrow T'\}$ is an fppf covering of T' it is enough to show each $T' \times_T T_j \rightarrow T$ factors through W_T . Hence we may assume f factors as $\varphi_j \circ f_j : T' \rightarrow T_j \rightarrow T$ for some j . In this case the condition $a|_{T'} \in F'(T')$ means that there exists some fppf covering $\{\psi_i : T'_i \rightarrow T'\}_{i \in I}$ and some morphisms $b_i : T'_i \rightarrow U'$ such that

$$\begin{array}{ccccc}
 T'_i & \xrightarrow{b_i} & U' & \xrightarrow{g} & U \\
 f_j \circ \psi_i \downarrow & & & & \downarrow \\
 T_j & \xrightarrow{a_j} & U & \longrightarrow & F
 \end{array}$$

is commutative. This commutativity means that there exists a morphism $r'_i : T'_i \rightarrow R$ such that $t \circ r'_i = a_j \circ f_j \circ \psi_i$, and $s \circ r'_i = g \circ b_i$. This implies that $\text{Im}(f_j \circ \psi_i) \subset W_j$ and we win. \square

The following lemma is not completely trivial although it looks like it should be trivial.

02WV **Lemma 10.3.** *Let S be a scheme. Let U be a scheme over S . Let $j = (s, t) : R \rightarrow U \times_S U$ be an étale equivalence relation on U over S . If the quotient U/R is an algebraic space, then $U \rightarrow U/R$ is étale and surjective. Hence $(U, R, U \rightarrow U/R)$ is a presentation of the algebraic space U/R .*

Proof. Denote $c : U \rightarrow U/R$ the morphism in question. Let T be a scheme and let $a : T \rightarrow U/R$ be a morphism. We have to show that the morphism (of schemes) $\pi : T \times_{a, U/R, c} U \rightarrow T$ is étale and surjective. The morphism a corresponds to an fppf covering $\{\varphi_i : T_i \rightarrow T\}$ and morphisms $a_i : T_i \rightarrow U$ such that $a_i \times a_{i'} : T_i \times_T T_{i'} \rightarrow U \times_S U$ factors through R , and such that $c \circ a_i = a \circ \varphi_i$. Hence

$$T_i \times_{\varphi_i, T} T \times_{a, U/R, c} U = T_i \times_{c \circ a_i, U/R, c} U = T_i \times_{a_i, U} U \times_{c, U/R, c} U = T_i \times_{a_i, U, t} R.$$

Since t is étale and surjective we conclude that the base change of π to T_i is surjective and étale. Since the property of being surjective and étale is local on the base in the fpqc topology (see Remark 4.3) we win. \square

0265 **Lemma 10.4.** *Let S be a scheme. Let U be a scheme over S . Let $j = (s, t) : R \rightarrow U \times_S U$ be an étale equivalence relation on U over S . Assume that U is affine. Then the quotient $F = U/R$ is an algebraic space, and $U \rightarrow F$ is étale and surjective.*

Proof. Since $j : R \rightarrow U \times_S U$ is a monomorphism we see that j is separated (see Schemes, Lemma 23.3). Since U is affine we see that $U \times_S U$ (which comes equipped with a monomorphism into the affine scheme $U \times U$) is separated. Hence we see that R is separated. In particular the morphisms s, t are separated as well as étale.

Since the composition $R \rightarrow U \times_S U \rightarrow U$ is locally of finite type we conclude that j is locally of finite type (see Morphisms, Lemma 14.8). As j is also a monomorphism it has finite fibres and we see that j is locally quasi-finite by Morphisms, Lemma 19.7. Altogether we see that j is separated and locally quasi-finite.

Our first step is to show that the quotient map $c : U \rightarrow F$ is representable. Consider a scheme T and a morphism $a : T \rightarrow F$. We have to show that the sheaf $G = T \times_{a, F, c} U$ is representable. As seen in the proofs of Lemmas 10.2 and 10.3 there exists an fppf covering $\{\varphi_i : T_i \rightarrow T\}_{i \in I}$ and morphisms $a_i : T_i \rightarrow U$ such that $a_i \times a_{i'} : T_i \times_T T_{i'} \rightarrow U \times_S U$ factors through R , and such that $c \circ a_i = a \circ \varphi_i$. As in the proof of Lemma 10.3 we see that

$$\begin{aligned} T_i \times_{\varphi_i, T} G &= T_i \times_{\varphi_i, T} T \times_{a, U/R, c} U \\ &= T_i \times_{c \circ a_i, U/R, c} U \\ &= T_i \times_{a_i, U} U \times_{c, U/R, c} U \\ &= T_i \times_{a_i, U, t} R \end{aligned}$$

Since t is separated and étale, and in particular separated and locally quasi-finite (by Morphisms, Lemmas 33.10 and 34.16) we see that the restriction of G to each T_i is representable by a morphism of schemes $X_i \rightarrow T_i$ which is separated and locally quasi-finite. By Descent, Lemma 36.1 we obtain a descent datum $(X_i, \varphi_{ii'})$ relative to the fppf-covering $\{T_i \rightarrow T\}$. Since each $X_i \rightarrow T_i$ is separated and locally quasi-finite we see by More on Morphisms, Lemma 47.1 that this descent datum is effective. Hence by Descent, Lemma 36.1 (2) we conclude that G is representable as desired.

The second step of the proof is to show that $U \rightarrow F$ is surjective and étale. This is clear from the above since in the first step above we saw that $G = T \times_{a, F, c} U$ is a scheme over T which base changes to schemes $X_i \rightarrow T_i$ which are surjective and étale. Thus $G \rightarrow T$ is surjective and étale (see Remark 4.3). Alternatively one can reread the proof of Lemma 10.3 in the current situation.

The third and final step is to show that the diagonal map $F \rightarrow F \times F$ is representable. We first observe that the diagram

$$\begin{array}{ccc} R & \longrightarrow & F \\ j \downarrow & & \downarrow \Delta \\ U \times_S U & \longrightarrow & F \times F \end{array}$$

is a fibre product square. By Lemma 3.4 the morphism $U \times_S U \rightarrow F \times F$ is representable (note that $h_U \times h_U = h_{U \times_S U}$). Moreover, by Lemma 5.7 the morphism $U \times_S U \rightarrow F \times F$ is surjective and étale (note also that étale and surjective occur in the lists of Remarks 4.3 and 4.2). It follows either from Lemma 3.3 and the

diagram above, or by writing $R \rightarrow F$ as $R \rightarrow U \rightarrow F$ and Lemmas 3.1 and 3.2 that $R \rightarrow F$ is representable as well. Let T be a scheme and let $a : T \rightarrow F \times F$ be a morphism. We have to show that $G = T \times_{a, F \times F, \Delta} F$ is representable. By what was said above the morphism (of schemes)

$$T' = (U \times_S U) \times_{F \times F, a} T \longrightarrow T$$

is surjective and étale. Hence $\{T' \rightarrow T\}$ is an étale covering of T . Note also that

$$T' \times_T G = T' \times_{U \times_S U, j} R$$

as can be seen contemplating the following cube

$$\begin{array}{ccccc}
 & & R & \xrightarrow{\quad} & F \\
 & \nearrow & \downarrow & & \downarrow \\
 T' \times_T G & \xrightarrow{\quad} & G & \xrightarrow{\quad} & G \\
 \downarrow & & \downarrow & & \downarrow \\
 & \nearrow & U \times_S U & \xrightarrow{\quad} & F \times F \\
 & & \downarrow & & \downarrow \\
 T' & \xrightarrow{\quad} & T & & T
 \end{array}$$

Hence we see that the restriction of G to T' is representable by a scheme X , and moreover that the morphism $X \rightarrow T'$ is a base change of the morphism j . Hence $X \rightarrow T'$ is separated and locally quasi-finite (see second paragraph of the proof). By Descent, Lemma 36.1 we obtain a descent datum (X, φ) relative to the fppf-covering $\{T' \rightarrow T\}$. Since $X \rightarrow T$ is separated and locally quasi-finite we see by More on Morphisms, Lemma 47.1 that this descent datum is effective. Hence by Descent, Lemma 36.1 (2) we conclude that G is representable as desired. \square

02WW **Theorem 10.5.** *Let S be a scheme. Let U be a scheme over S . Let $j = (s, t) : R \rightarrow U \times_S U$ be an étale equivalence relation on U over S . Then the quotient U/R is an algebraic space, and $U \rightarrow U/R$ is étale and surjective, in other words $(U, R, U \rightarrow U/R)$ is a presentation of U/R .*

Proof. By Lemma 10.3 it suffices to prove that U/R is an algebraic space. Let $U' \rightarrow U$ be a surjective, étale morphism. Then $\{U' \rightarrow U\}$ is in particular an fppf covering. Let R' be the restriction of R to U' , see Groupoids, Definition 3.3. According to Groupoids, Lemma 20.6 we see that $U/R \cong U'/R'$. By Lemma 10.1 R' is an étale equivalence relation on U' . Thus we may replace U by U' .

We apply the previous remark to $U' = \coprod U_i$, where $U = \bigcup U_i$ is an affine open covering of S . Hence we may and do assume that $U = \coprod U_i$ where each U_i is an affine scheme.

Consider the restriction R_i of R to U_i . By Lemma 10.1 this is an étale equivalence relation. Set $F_i = U_i/R_i$ and $F = U/R$. It is clear that $\coprod F_i \rightarrow F$ is surjective. By Lemma 10.2 each $F_i \rightarrow F$ is representable, and an open immersion. By Lemma 10.4 applied to (U_i, R_i) we see that F_i is an algebraic space. Then by Lemma 10.3 we see that $U_i \rightarrow F_i$ is étale and surjective. From Lemma 8.3 it follows that $\coprod F_i$ is an algebraic space. Finally, we have verified all hypotheses of Lemma 8.4 and it follows that $F = U/R$ is an algebraic space. \square

11. Algebraic spaces, retrofitted

02WX We start building our arsenal of lemmas dealing with algebraic spaces. The first result says that in Definition 6.1 we can weaken the condition on the diagonal as follows.

0BGQ **Lemma 11.1.** *Let S be a scheme contained in Sch_{fppf} . Let F be a sheaf on $(Sch/S)_{fppf}$ such that there exists $U \in \text{Ob}((Sch/S)_{fppf})$ and a map $U \rightarrow F$ which is representable, surjective, and étale. Then F is an algebraic space.*

Proof. Set $R = U \times_F U$. This is a scheme as $U \rightarrow F$ is assumed representable. The projections $s, t : R \rightarrow U$ are étale as $U \rightarrow F$ is assumed étale. The map $j = (t, s) : R \rightarrow U \times_S U$ is a monomorphism and an equivalence relation as $R = U \times_F U$. By Theorem 10.5 the quotient sheaf $F' = U/R$ is an algebraic space and $U \rightarrow F'$ is surjective and étale. Again since $R = U \times_F U$ we obtain a canonical factorization $U \rightarrow F' \rightarrow F$ and $F' \rightarrow F$ is an injective map of sheaves. On the other hand, $U \rightarrow F$ is surjective as a map of sheaves by Lemma 5.9. Thus $F' \rightarrow F$ is also surjective and we conclude $F' = F$ is an algebraic space. \square

0BGR **Lemma 11.2.** *Let S be a scheme contained in Sch_{fppf} . Let G be an algebraic space over S , let F be a sheaf on $(Sch/S)_{fppf}$, and let $G \rightarrow F$ be a representable transformation of functors which is surjective and étale. Then F is an algebraic space.*

Proof. Pick a scheme U and a surjective étale morphism $U \rightarrow G$. Since G is an algebraic space $U \rightarrow G$ is representable. Hence the composition $U \rightarrow G \rightarrow F$ is representable, surjective, and étale. See Lemmas 3.2 and 5.4. Thus F is an algebraic space by Lemma 11.1. \square

02WY **Lemma 11.3.** *Let S be a scheme contained in Sch_{fppf} . Let F be an algebraic space over S . Let $G \rightarrow F$ be a representable transformation of functors. Then G is an algebraic space.*

Proof. By Lemma 3.5 we see that G is a sheaf. The diagram

$$\begin{array}{ccc} G \times_F G & \longrightarrow & F \\ \downarrow & & \downarrow \Delta_F \\ G \times G & \longrightarrow & F \times F \end{array}$$

is cartesian. Hence we see that $G \times_F G \rightarrow G \times G$ is representable by Lemma 3.3. By Lemma 3.6 we see that $G \rightarrow G \times_F G$ is representable. Hence $\Delta_G : G \rightarrow G \times G$ is representable as a composition of representable transformations, see Lemma 3.2. Finally, let U be an object of $(Sch/S)_{fppf}$ and let $U \rightarrow F$ be surjective and étale. By assumption $U \times_F G$ is representable by a scheme U' . By Lemma 5.5 the morphism $U' \rightarrow G$ is surjective and étale. This verifies the final condition of Definition 6.1 and we win. \square

02WZ **Lemma 11.4.** *Let S be a scheme contained in Sch_{fppf} . Let F, G be algebraic spaces over S . Let $G \rightarrow F$ be a representable morphism. Let $U \in \text{Ob}((Sch/S)_{fppf})$, and $q : U \rightarrow F$ surjective and étale. Set $V = G \times_F U$. Finally, let \mathcal{P} be a property of morphisms of schemes as in Definition 5.1. Then $G \rightarrow F$ has property \mathcal{P} if and only if $V \rightarrow U$ has property \mathcal{P} .*

Proof. (This lemma follows from Lemmas 5.5 and 5.6, but we give a direct proof here also.) It is clear from the definitions that if $G \rightarrow F$ has property \mathcal{P} , then $V \rightarrow U$ has property \mathcal{P} . Conversely, assume $V \rightarrow U$ has property \mathcal{P} . Let $T \rightarrow F$ be a morphism from a scheme to F . Let $T' = T \times_F G$ which is a scheme since $G \rightarrow F$ is representable. We have to show that $T' \rightarrow T$ has property \mathcal{P} . Consider the commutative diagram of schemes

$$\begin{array}{ccccc} V & \longleftarrow & T \times_F V & \longrightarrow & T \times_F G = T' \\ \downarrow & & \downarrow & & \downarrow \\ U & \longleftarrow & T \times_F U & \longrightarrow & T \end{array}$$

where both squares are fibre product squares. Hence we conclude the middle arrow has property \mathcal{P} as a base change of $V \rightarrow U$. Finally, $\{T \times_F U \rightarrow T\}$ is a fppf covering as it is surjective étale, and hence we conclude that $T' \rightarrow T$ has property \mathcal{P} as it is local on the base in the fppf topology. \square

03I2 **Lemma 11.5.** *Let S be a scheme contained in Sch_{fppf} . Let $G \rightarrow F$ be a transformation of presheaves on $(Sch/S)_{fppf}$. Let \mathcal{P} be a property of morphisms of schemes. Assume*

- (1) \mathcal{P} is preserved under any base change, fppf local on the base, and morphisms of type \mathcal{P} satisfy descent for fppf coverings, see Descent, Definition 33.1,
- (2) G is a sheaf,
- (3) F is an algebraic space,
- (4) there exists a $U \in \text{Ob}((Sch/S)_{fppf})$ and a surjective étale morphism $U \rightarrow F$ such that $V = G \times_F U$ is representable, and
- (5) $V \rightarrow U$ has \mathcal{P} .

Then G is an algebraic space, $G \rightarrow F$ is representable and has property \mathcal{P} .

Proof. Let $R = U \times_F U$, and denote $t, s : R \rightarrow U$ the projection morphisms as usual. Let T be a scheme and let $T \rightarrow F$ be a morphism. Then $U \times_F T \rightarrow T$ is surjective étale, hence $\{U \times_F T \rightarrow T\}$ is a covering for the étale topology. Consider

$$W = G \times_F (U \times_F T) = V \times_F T = V \times_U (U \times_F T).$$

It is a scheme since F is an algebraic space. The morphism $W \rightarrow U \times_F T$ has property \mathcal{P} since it is a base change of $V \rightarrow U$. There is an isomorphism

$$\begin{aligned} W \times_T (U \times_F T) &= (G \times_F (U \times_F T)) \times_T (U \times_F T) \\ &= (U \times_F T) \times_T (G \times_F (U \times_F T)) \\ &= (U \times_F T) \times_T W \end{aligned}$$

over $(U \times_F T) \times_T (U \times_F T)$. The middle equality maps $((g, (u_1, t)), (u_2, t))$ to $((u_1, t), (g, (u_2, t)))$. This defines a descent datum for $W/U \times_F T/T$, see Descent, Definition 31.1. This follows from Descent, Lemma 36.1. Namely we have a sheaf $G \times_F T$, whose base change to $U \times_F T$ is represented by W and the isomorphism above is the one from the proof of Descent, Lemma 36.1. By assumption on \mathcal{P} the descent datum above is representable. Hence by the last statement of Descent, Lemma 36.1 we see that $G \times_F T$ is representable. This proves that $G \rightarrow F$ is a representable transformation of functors.

As $G \rightarrow F$ is representable, we see that G is an algebraic space by Lemma 11.3. The fact that $G \rightarrow F$ has property \mathcal{P} now follows from Lemma 11.4. \square

02X1 **Lemma 11.6.** *Let S be a scheme contained in Sch_{fppf} . Let F, G be algebraic spaces over S . Let $a : F \rightarrow G$ be a morphism. Given any $V \in \text{Ob}((Sch/S)_{fppf})$ and a surjective étale morphism $q : V \rightarrow G$ there exists a $U \in \text{Ob}((Sch/S)_{fppf})$ and a commutative diagram*

$$\begin{array}{ccc} U & \xrightarrow{\alpha} & V \\ p \downarrow & & \downarrow q \\ F & \xrightarrow{a} & G \end{array}$$

with p surjective and étale.

Proof. First choose $W \in \text{Ob}((Sch/S)_{fppf})$ with surjective étale morphism $W \rightarrow F$. Next, put $U = W \times_G V$. Since G is an algebraic space we see that U is isomorphic to an object of $(Sch/S)_{fppf}$. As q is surjective étale, we see that $U \rightarrow W$ is surjective étale (see Lemma 5.5). Thus $U \rightarrow F$ is surjective étale as a composition of surjective étale morphisms (see Lemma 5.4). \square

12. Immersions and Zariski coverings of algebraic spaces

02YT At this point an interesting phenomenon occurs. We have already defined the notion of an open immersion of algebraic spaces (through Definition 5.1) but we have yet to define the notion of a *point*¹. Thus the *Zariski topology* of an algebraic space has already been defined, but there is no space yet!

Perhaps superfluously we formally introduce immersions as follows.

02YU **Definition 12.1.** Let $S \in \text{Ob}(Sch_{fppf})$ be a scheme. Let F be an algebraic space over S .

- (1) A morphism of algebraic spaces over S is called an *open immersion* if it is representable, and an open immersion in the sense of Definition 5.1.
- (2) An *open subspace* of F is a subfunctor $F' \subset F$ such that F' is an algebraic space and $F' \rightarrow F$ is an open immersion.
- (3) A morphism of algebraic spaces over S is called a *closed immersion* if it is representable, and a closed immersion in the sense of Definition 5.1.
- (4) A *closed subspace* of F is a subfunctor $F' \subset F$ such that F' is an algebraic space and $F' \rightarrow F$ is a closed immersion.
- (5) A morphism of algebraic spaces over S is called an *immersion* if it is representable, and an immersion in the sense of Definition 5.1.
- (6) A *locally closed subspace* of F is a subfunctor $F' \subset F$ such that F' is an algebraic space and $F' \rightarrow F$ is an immersion.

We note that these definitions make sense since an immersion is in particular a monomorphism (see Schemes, Lemma 23.8 and Lemma 5.8), and hence the image of an immersion $G \rightarrow F$ of algebraic spaces is a subfunctor $F' \subset F$ which is (canonically) isomorphic to G . Thus some of the discussion of Schemes, Section 10 carries over to the setting of algebraic spaces.

02YV **Lemma 12.2.** *Let $S \in \text{Ob}(Sch_{fppf})$ be a scheme. A composition of (closed, resp. open) immersions of algebraic spaces over S is a (closed, resp. open) immersion of algebraic spaces over S .*

¹We will associate a topological space to an algebraic space in Properties of Spaces, Section 4, and its opens will correspond exactly to the open subspaces defined below.

Proof. See Lemma 5.4 and Remarks 4.3 (see very last line of that remark) and 4.2. \square

02YW **Lemma 12.3.** *Let $S \in \text{Ob}(Sch_{fppf})$ be a scheme. A base change of a (closed, resp. open) immersion of algebraic spaces over S is a (closed, resp. open) immersion of algebraic spaces over S .*

Proof. See Lemma 5.5 and Remark 4.3 (see very last line of that remark). \square

02YX **Lemma 12.4.** *Let $S \in \text{Ob}(Sch_{fppf})$ be a scheme. Let F be an algebraic space over S . Let F_1, F_2 be locally closed subspaces of F . If $F_1 \subset F_2$ as subfunctors of F , then F_1 is a locally closed subspace of F_2 . Similarly for closed and open subspaces.*

Proof. Let $T \rightarrow F_2$ be a morphism with T a scheme. Since $F_2 \rightarrow F$ is a monomorphism, we see that $T \times_{F_2} F_1 = T \times_F F_1$. The lemma follows formally from this. \square

Let us formally define the notion of a Zariski open covering of algebraic spaces. Note that in Lemma 8.4 we have already encountered such open coverings as a method for constructing algebraic spaces.

02YY **Definition 12.5.** Let $S \in \text{Ob}(Sch_{fppf})$ be a scheme. Let F be an algebraic space over S . A Zariski covering $\{F_i \subset F\}_{i \in I}$ of F is given by a set I and a collection of open subspaces $F_i \subset F$ such that $\coprod F_i \rightarrow F$ is a surjective map of sheaves.

Note that if T is a schemes, and $a : T \rightarrow F$ is a morphism, then each of the fibre products $T \times_F F_i$ is identified with an open subscheme $T_i \subset T$. The final condition of the definition signifies exactly that $T = \bigcup_{i \in I} T_i$.

It is clear that the collection F_{Zar} of open subspaces of F is a set (as $(Sch/S)_{fppf}$ is a site, hence a set). Moreover, we can turn F_{Zar} into a category by letting the morphisms be inclusions of subfunctors (which are automatically open immersions by Lemma 12.4). Finally, Definition 12.5 provides the notion of a Zariski covering $\{F_i \rightarrow F\}_{i \in I}$ in the category F_{Zar} . Hence, just as in the case of a topological space (see Sites, Example 6.4) by suitably choosing a set of coverings we may obtain a Zariski site of the algebraic space F .

02YZ **Definition 12.6.** Let $S \in \text{Ob}(Sch_{fppf})$ be a scheme. Let F be an algebraic space over S . A small Zariski site F_{Zar} of an algebraic space F is one of the sites described above.

Hence this gives a notion of what it means for something to be true Zariski locally on an algebraic space, which is how we will use this notion. In general the Zariski topology is not fine enough for our purposes. For example we can consider the category of Zariski sheaves on an algebraic space. It will turn out that this is not the correct thing to consider, even for quasi-coherent sheaves. One only gets the desired result when using the étale or fppf site of F to define quasi-coherent sheaves.

13. Separation conditions on algebraic spaces

02X3 A separation condition on an algebraic space F is a condition on the diagonal morphism $F \rightarrow F \times F$. Let us first list the properties the diagonal has automatically. Since the diagonal is representable by definition the following lemma makes sense (through Definition 5.1).

02X4 **Lemma 13.1.** *Let S be a scheme contained in Sch_{fppf} . Let F be an algebraic space over S . Let $\Delta : F \rightarrow F \times F$ be the diagonal morphism. Then*

- (1) Δ is locally of finite type,
- (2) Δ is a monomorphism,
- (3) Δ is separated, and
- (4) Δ is locally quasi-finite.

Proof. Let $F = U/R$ be a presentation of F . As in the proof of Lemma 10.4 the diagram

$$\begin{array}{ccc} R & \longrightarrow & F \\ j \downarrow & & \downarrow \Delta \\ U \times_S U & \longrightarrow & F \times F \end{array}$$

is cartesian. Hence according to Lemma 11.4 it suffices to show that j has the properties listed in the lemma. (Note that each of the properties (1) – (4) occur in the lists of Remarks 4.1 and 4.3.) Since j is an equivalence relation it is a monomorphism. Hence it is separated by Schemes, Lemma 23.3. As R is an étale equivalence relation we see that $s, t : R \rightarrow U$ are étale. Hence s, t are locally of finite type. Then it follows from Morphisms, Lemma 14.8 that j is locally of finite type. Finally, as it is a monomorphism its fibres are finite. Thus we conclude that it is locally quasi-finite by Morphisms, Lemma 19.7. \square

Here are some common types of separation conditions, relative to the base scheme S . There is also an absolute notion of these conditions which we will discuss in Properties of Spaces, Section 3. Moreover, we will discuss separation conditions for a morphism of algebraic spaces in Morphisms of Spaces, Section 4.

02X5 **Definition 13.2.** *Let S be a scheme contained in Sch_{fppf} . Let F be an algebraic space over S . Let $\Delta : F \rightarrow F \times F$ be the diagonal morphism.*

- (1) We say F is *separated over S* if Δ is a closed immersion.
- (2) We say F is *locally separated over S^2* if Δ is an immersion.
- (3) We say F is *quasi-separated over S* if Δ is quasi-compact.
- (4) We say F is *Zariski locally quasi-separated over S^3* if there exists a Zariski covering $F = \bigcup_{i \in I} F_i$ such that each F_i is quasi-separated.

Note that if the diagonal is quasi-compact (when F is separated or quasi-separated) then the diagonal is actually quasi-finite and separated, hence quasi-affine (by More on Morphisms, Lemma 38.2).

14. Examples of algebraic spaces

02Z0 In this section we construct some examples of algebraic spaces. Some of these were suggested by B. Conrad. Since we do not yet have a lot of theory at our disposal the discussion is a bit awkward in some places.

02Z1 **Example 14.1.** Let k be a field of characteristic $\neq 2$. Let $U = \mathbf{A}_k^1$. Set

$$j : R = \Delta \amalg \Gamma \longrightarrow U \times_k U$$

²In the literature this often refers to quasi-separated and locally separated algebraic spaces.

³This definition was suggested by B. Conrad.

where $\Delta = \{(x, x) \mid x \in \mathbf{A}_k^1\}$ and $\Gamma = \{(x, -x) \mid x \in \mathbf{A}_k^1, x \neq 0\}$. It is clear that $s, t : R \rightarrow U$ are étale, and hence j is an étale equivalence relation. The quotient $X = U/R$ is an algebraic space by Theorem 10.5. Since R is quasi-compact we see that X is quasi-separated. On the other hand, X is not locally separated because the morphism j is not an immersion.

03FN **Example 14.2.** Let k be a field. Let k'/k be a degree 2 Galois extension with $\text{Gal}(k'/k) = \{1, \sigma\}$. Let $S = \text{Spec}(k[x])$ and $U = \text{Spec}(k'[x])$. Note that

$$U \times_S U = \text{Spec}((k' \otimes_k k')[x]) = \Delta(U) \amalg \Delta'(U)$$

where $\Delta' = (1, \sigma) : U \rightarrow U \times_S U$. Take

$$R = \Delta(U) \amalg \Delta'(U \setminus \{0_U\})$$

where $0_U \in U$ denotes the k' -rational point whose x -coordinate is zero. It is easy to see that R is an étale equivalence relation on U over S and hence $X = U/R$ is an algebraic space by Theorem 10.5. Here are some properties of X (some of which will not make sense until later):

- (1) $X \rightarrow S$ is an isomorphism over $S \setminus \{0_S\}$,
- (2) the morphism $X \rightarrow S$ is étale (see Properties of Spaces, Definition 16.2)
- (3) the fibre 0_X of $X \rightarrow S$ over 0_S is isomorphic to $\text{Spec}(k') = 0_U$,
- (4) X is not a scheme because if it were, then $\mathcal{O}_{X,0_X}$ would be a local domain $(\mathcal{O}, \mathfrak{m}, \kappa)$ with fraction field $k(x)$, with $x \in \mathfrak{m}$ and residue field $\kappa = k'$ which is impossible,
- (5) X is not separated, but it is locally separated and quasi-separated,
- (6) there exists a surjective, finite, étale morphism $S' \rightarrow S$ such that the base change $X' = S' \times_S X$ is a scheme (namely, if we base change to $S' = \text{Spec}(k'[x])$ then U splits into two copies of S' and X' becomes isomorphic to the affine line with 0 doubled, see Schemes, Example 14.3), and
- (7) if we think of X as a finite type algebraic space over $\text{Spec}(k)$, then similarly the base change $X_{k'}$ is a scheme but X is not a scheme.

In particular, this gives an example of a descent datum for schemes relative to the covering $\{\text{Spec}(k') \rightarrow \text{Spec}(k)\}$ which is not effective.

See also Examples, Lemma 58.1, which shows that descent data need not be effective even for a projective morphism of schemes. That example gives a smooth separated algebraic space of dimension 3 over \mathbf{C} which is not a scheme.

We will use the following lemma as a convenient way to construct algebraic spaces as quotients of schemes by free group actions.

02Z2 **Lemma 14.3.** *Let $U \rightarrow S$ be a morphism of Sch_{fppf} . Let G be an abstract group. Let $G \rightarrow \text{Aut}_S(U)$ be a group homomorphism. Assume*

- (*) *if $u \in U$ is a point, and $g(u) = u$ for some non-identity element $g \in G$, then g induces a nontrivial automorphism of $\kappa(u)$.*

Then

$$j : R = \coprod_{g \in G} U \longrightarrow U \times_S U, \quad (g, x) \longmapsto (g(x), x)$$

is an étale equivalence relation and hence

$$F = U/R$$

is an algebraic space by Theorem 10.5.

Proof. In the statement of the lemma the symbol $\text{Aut}_S(U)$ denotes the group of automorphisms of U over S . Assume $(*)$ holds. Let us show that

$$j : R = \coprod_{g \in G} U \longrightarrow U \times_S U, \quad (g, x) \longmapsto (g(x), x)$$

is a monomorphism. This signifies that if T is a nonempty scheme, and $h : T \rightarrow U$ is a T -valued point such that $g \circ h = g' \circ h$ then $g = g'$. Suppose $T \neq \emptyset$, $h : T \rightarrow U$ and $g \circ h = g' \circ h$. Let $t \in T$. Consider the composition $\text{Spec}(\kappa(t)) \rightarrow \text{Spec}(\kappa(h(t))) \rightarrow U$. Then we conclude that $g^{-1} \circ g'$ fixes $u = h(t)$ and acts as the identity on its residue field. Hence $g = g'$ by $(*)$.

Thus if $(*)$ holds we see that j is a relation (see Groupoids, Definition 3.1). Moreover, it is an equivalence relation since on T -valued points for a connected scheme T we see that $R(T) = G \times U(T) \rightarrow U(T) \times U(T)$ (recall that we always work over S). Moreover, the morphisms $s, t : R \rightarrow U$ are étale since R is a disjoint product of copies of U . This proves that $j : R \rightarrow U \times_S U$ is an étale equivalence relation. \square

Given a scheme U and an action of a group G on U we say the action of G on U is *free* if condition $(*)$ of Lemma 14.3 holds. This is equivalent to the notion of a free action of the constant group scheme G_S on U as defined in Groupoids, Definition 10.2. The lemma can be interpreted as saying that quotients of schemes by free actions of groups exist in the category of algebraic spaces.

02Z3 **Definition 14.4.** Notation $U \rightarrow S$, G , R as in Lemma 14.3. If the action of G on U satisfies $(*)$ we say G *acts freely* on the scheme U . In this case the algebraic space U/R is denoted U/G and is called the *quotient of U by G* .

This notation is consistent with the notation U/G introduced in Groupoids, Definition 20.1. We will later make sense of the quotient as an algebraic stack without any assumptions on the action whatsoever; when we do this we will use the notation $[U/G]$. Before we discuss the examples we prove some more lemmas to facilitate the discussion. Here is a lemma discussing the various separation conditions for this quotient when G is finite.

02Z4 **Lemma 14.5.** *Notation and assumptions as in Lemma 14.3. Assume G is finite. Then*

- (1) *if $U \rightarrow S$ is quasi-separated, then U/G is quasi-separated over S , and*
- (2) *if $U \rightarrow S$ is separated, then U/G is separated over S .*

Proof. In the proof of Lemma 13.1 we saw that it suffices to prove the corresponding properties for the morphism $j : R \rightarrow U \times_S U$. If $U \rightarrow S$ is quasi-separated, then for every affine open $V \subset U$ which maps into an affine of S the opens $g(V) \cap V$ are quasi-compact. It follows that j is quasi-compact. If $U \rightarrow S$ is separated, the diagonal $\Delta_{U/S}$ is a closed immersion. Hence $j : R \rightarrow U \times_S U$ is a finite coproduct of closed immersions with disjoint images. Hence j is a closed immersion. \square

02Z5 **Lemma 14.6.** *Notation and assumptions as in Lemma 14.3. If $\text{Spec}(k) \rightarrow U/G$ is a morphism, then there exist*

- (1) *a finite Galois extension k'/k ,*
- (2) *a finite subgroup $H \subset G$,*
- (3) *an isomorphism $H \rightarrow \text{Gal}(k'/k)$, and*
- (4) *an H -equivariant morphism $\text{Spec}(k') \rightarrow U$.*

Conversely, such data determine a morphism $\text{Spec}(k) \rightarrow U/G$.

Proof. Consider the fibre product $V = \text{Spec}(k) \times_{U/G} U$. Here is a diagram

$$\begin{array}{ccc} V & \longrightarrow & U \\ \downarrow & & \downarrow \\ \text{Spec}(k) & \longrightarrow & U/G \end{array}$$

Then V is a nonempty scheme étale over $\text{Spec}(k)$ and hence is a disjoint union $V = \coprod_{i \in I} \text{Spec}(k_i)$ of spectra of fields k_i finite separable over k (Morphisms, Lemma 34.7). We have

$$\begin{aligned} V \times_{\text{Spec}(k)} V &= (\text{Spec}(k) \times_{U/G} U) \times_{\text{Spec}(k)} (\text{Spec}(k) \times_{U/G} U) \\ &= \text{Spec}(k) \times_{U/G} U \times_{U/G} U \\ &= \text{Spec}(k) \times_{U/G} U \times G \\ &= V \times G \end{aligned}$$

The action of G on U induces an action of $a : G \times V \rightarrow V$. The displayed equality means that $G \times V \rightarrow V \times_{\text{Spec}(k)} V$, $(g, v) \mapsto (a(g, v), v)$ is an isomorphism. In particular we see that for every i we have an isomorphism $H_i \times \text{Spec}(k_i) \rightarrow \text{Spec}(k_i \otimes_k k_i)$ where $H_i \subset G$ is the subgroup of elements fixing $i \in I$. Thus H_i is finite and is the Galois group of k_i/k . We omit the converse construction. \square

It follows from this lemma for example that if k'/k is a finite Galois extension, then $\text{Spec}(k')/\text{Gal}(k'/k) \cong \text{Spec}(k)$. What happens if the extension is infinite? Here is an example.

02Z6 **Example 14.7.** Let $S = \text{Spec}(\mathbf{Q})$. Let $U = \text{Spec}(\overline{\mathbf{Q}})$. Let $G = \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ with obvious action on U . Then by construction property (*) of Lemma 14.3 holds and we obtain an algebraic space

$$X = \text{Spec}(\overline{\mathbf{Q}})/G \longrightarrow S = \text{Spec}(\mathbf{Q}).$$

Of course this is totally ridiculous as an approximation of S ! Namely, by the Artin-Schreier theorem, see [Jac64, Theorem 17, page 316], the only finite subgroups of $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ are $\{1\}$ and the conjugates of the order two group $\text{Gal}(\overline{\mathbf{Q}}/\overline{\mathbf{Q}} \cap \mathbf{R})$. Hence, if $\text{Spec}(k) \rightarrow X$ is a morphism with k algebraic over \mathbf{Q} , then it follows from Lemma 14.6 and the theorem just mentioned that either k is $\overline{\mathbf{Q}}$ or isomorphic to $\overline{\mathbf{Q}} \cap \mathbf{R}$.

What is wrong with the example above is that the Galois group comes equipped with a topology, and this should somehow be part of any construction of a quotient of $\text{Spec}(\overline{\mathbf{Q}})$. The following example is much more reasonable in my opinion and may actually occur in “nature”.

02Z7 **Example 14.8.** Let k be a field of characteristic zero. Let $U = \mathbf{A}_k^1$ and let $G = \mathbf{Z}$. As action we take $n(x) = x+n$, i.e., the action of \mathbf{Z} on the affine line by translation. The only fixed point is the generic point and it is clearly the case that \mathbf{Z} injects into the automorphism group of the field $k(x)$. (This is where we use the characteristic zero assumption.) Consider the morphism

$$\gamma : \text{Spec}(k(x)) \longrightarrow X = \mathbf{A}_k^1/\mathbf{Z}$$

of the generic point of the affine line into the quotient. We claim that this morphism does not factor through any monomorphism $\text{Spec}(L) \rightarrow X$ of the spectrum of a field to X . (Contrary to what happens for schemes, see Schemes, Section 13.) In fact,

since \mathbf{Z} does not have any nontrivial finite subgroups we see from Lemma 14.6 that for any such factorization $k(x) = L$. Finally, γ is not a monomorphism since

$$\mathrm{Spec}(k(x)) \times_{\gamma, X, \gamma} \mathrm{Spec}(k(x)) \cong \mathrm{Spec}(k(x)) \times \mathbf{Z}.$$

This example suggests that in order to define points of an algebraic space X we should consider equivalence classes of morphisms from spectra of fields into X and not the set of monomorphisms from spectra of fields.

We finish with a truly awful example.

02Z8 **Example 14.9.** Let k be a field. Let $A = \prod_{n \in \mathbf{N}} k$ be the infinite product. Set $U = \mathrm{Spec}(A)$ seen as a scheme over $S = \mathrm{Spec}(k)$. Note that the projection maps $\mathrm{pr}_n : A \rightarrow k$ define open and closed immersions $f_n : S \rightarrow U$. Set

$$R = U \amalg \coprod_{(n,m) \in \mathbf{N}^2, n \neq m} S$$

with morphism j equal to $\Delta_{U/S}$ on the component U and $j = (f_n, f_m)$ on the component S corresponding to (n, m) . It is clear from the remark above that s, t are étale. It is also clear that j is an equivalence relation. Hence we obtain an algebraic space

$$X = U/R.$$

To see what this means we specialize to the case where the field k is finite with q elements. Let us first discuss the topological space $|U|$ associated to the scheme U a little bit. All elements of A satisfy $x^q = x$. Hence every residue field of A is isomorphic to k , and all points of U are closed. But the topology on U isn't the discrete topology. Let $u_n \in |U|$ be the point corresponding to f_n . As mentioned above the points u_n are the open points (and hence isolated). This implies there have to be other points since we know U is quasi-compact, see Algebra, Lemma 16.10 (hence not equal to an infinite discrete set). Another way to see this is because the (proper) ideal

$$I = \{x = (x_n) \in A \mid \text{all but a finite number of } x_n \text{ are zero}\}$$

is contained in a maximal ideal. Note also that every element x of A is of the form $x = ue$ where u is a unit and e is an idempotent. Hence a basis for the topology of A consists of open and closed subsets (see Algebra, Lemma 20.1.) So the topology on $|U|$ is totally disconnected, but nontrivial. Finally, note that $\{u_n\}$ is dense in $|U|$.

We will later define a topological space $|X|$ associated to X , see Properties of Spaces, Section 4. What can we say about $|X|$? It turns out that the map $|U| \rightarrow |X|$ is surjective and continuous. All the points u_n map to the same point x_0 of $|X|$, and none of the other points get identified. Since $\{u_n\}$ is dense in $|U|$ we conclude that the closure of x_0 in $|X|$ is $|X|$. In other words $|X|$ is irreducible and x_0 is a generic point of $|X|$. This seems bizarre since also x_0 is the image of a section $S \rightarrow X$ of the structure morphism $X \rightarrow S$ (and in the case of schemes this would imply it was a closed point, see Morphisms, Lemma 19.2).

Whatever you think is actually going on in this example, it certainly shows that some care has to be exercised when defining irreducible components, connectedness, etc of algebraic spaces.

15. Change of big site

03FO In this section we briefly discuss what happens when we change big sites. The upshot is that we can always enlarge the big site at will, hence we may assume any set of schemes we want to consider is contained in the big fppf site over which we consider our algebraic space. Here is a precise statement of the result.

03FP **Lemma 15.1.** *Suppose given big sites Sch_{fppf} and Sch'_{fppf} . Assume that Sch_{fppf} is contained in Sch'_{fppf} , see Topologies, Section 11. Let S be an object of Sch_{fppf} . Let*

$$\begin{aligned} g &: Sh((Sch/S)_{fppf}) \longrightarrow Sh((Sch'/S)_{fppf}), \\ f &: Sh((Sch'/S)_{fppf}) \longrightarrow Sh((Sch/S)_{fppf}) \end{aligned}$$

be the morphisms of topoi of Topologies, Lemma 11.2. Let F be a sheaf of sets on $(Sch/S)_{fppf}$. Then

- (1) if F is representable by a scheme $X \in \text{Ob}((Sch/S)_{fppf})$ over S , then $f^{-1}F$ is representable too, in fact it is representable by the same scheme X , now viewed as an object of $(Sch'/S)_{fppf}$, and
- (2) if F is an algebraic space over S , then $f^{-1}F$ is an algebraic space over S also.

Proof. Let $X \in \text{Ob}((Sch/S)_{fppf})$. Let us write h_X for the representable sheaf on $(Sch/S)_{fppf}$ associated to X , and h'_X for the representable sheaf on $(Sch'/S)_{fppf}$ associated to X . By the description of f^{-1} in Topologies, Section 11 we see that $f^{-1}h_X = h'_X$. This proves (1).

Next, suppose that F is an algebraic space over S . By Lemma 9.1 this means that $F = h_U/h_R$ for some étale equivalence relation $R \rightarrow U \times_S U$ in $(Sch/S)_{fppf}$. Since f^{-1} is an exact functor we conclude that $f^{-1}F = h'_U/h'_R$. Hence $f^{-1}F$ is an algebraic space over S by Theorem 10.5. \square

Note that this lemma is purely set theoretical and has virtually no content. Moreover, it is not true (in general) that the restriction of an algebraic space over the bigger site is an algebraic space over the smaller site (simply by reasons of cardinality). Hence we can only ever use a simple lemma of this kind to enlarge the base category and never to shrink it.

04W1 **Lemma 15.2.** *Suppose Sch_{fppf} is contained in Sch'_{fppf} . Let S be an object of Sch_{fppf} . Denote Spaces/S the category of algebraic spaces over S defined using Sch_{fppf} . Similarly, denote Spaces'/S the category of algebraic spaces over S defined using Sch'_{fppf} . The construction of Lemma 15.1 defines a fully faithful functor*

$$\text{Spaces}/S \longrightarrow \text{Spaces}'/S$$

whose essential image consists of those $X' \in \text{Ob}(\text{Spaces}'/S)$ such that there exist $U, R \in \text{Ob}((Sch/S)_{fppf})$ ⁴ and morphisms

$$U \longrightarrow X' \quad \text{and} \quad R \longrightarrow U \times_{X'} U$$

⁴Requiring the existence of R is necessary because of our choice of the function *Bound* in Sets, Equation (9.1.1). The size of the fibre product $U \times_{X'} U$ can grow faster than *Bound* in terms of the size of U . We can illustrate this by setting $S = \text{Spec}(A)$, $U = \text{Spec}(A[x_i, i \in I])$ and $R = \coprod_{(\lambda_i) \in A^I} \text{Spec}(A[x_i, y_i]/(x_i - \lambda_i y_i))$. In this case the size of R grows like κ^κ where κ is the size of U .

in $Sh((Sch'/S)_{fppf})$ which are surjective as maps of sheaves (for example if the displayed morphisms are surjective and étale).

Proof. In Sites, Lemma 20.8 we have seen that the functor $f^{-1} : Sh((Sch/S)_{fppf}) \rightarrow Sh((Sch'/S)_{fppf})$ is fully faithful (see discussion in Topologies, Section 11). Hence we see that the displayed functor of the lemma is fully faithful.

Suppose that $X' \in \text{Ob}(Spaces'/S)$ such that there exists $U \in \text{Ob}((Sch/S)_{fppf})$ and a map $U \rightarrow X'$ in $Sh((Sch'/S)_{fppf})$ which is surjective as a map of sheaves. Let $U' \rightarrow X'$ be a surjective étale morphism with $U' \in \text{Ob}((Sch'/S)_{fppf})$. Let $\kappa = \text{size}(U)$, see Sets, Section 9. Then U has an affine open covering $U = \bigcup_{i \in I} U_i$ with $|I| \leq \kappa$. Observe that $U' \times_{X'} U \rightarrow U$ is étale and surjective. For each i we can pick a quasi-compact open $U'_i \subset U'$ such that $U'_i \times_{X'} U_i \rightarrow U_i$ is surjective (because the scheme $U' \times_{X'} U_i$ is the union of the Zariski opens $W \times_{X'} U_i$ for $W \subset U'$ affine and because $U' \times_{X'} U_i \rightarrow U_i$ is étale hence open). Then $\coprod_{i \in I} U'_i \rightarrow X$ is surjective étale because of our assumption that $U \rightarrow X$ and hence $\coprod U_i \rightarrow X$ is a surjection of sheaves (details omitted). Because $U'_i \times_{X'} U \rightarrow U'_i$ is a surjection of sheaves and because U'_i is quasi-compact, we can find a quasi-compact open $W_i \subset U'_i \times_{X'} U$ such that $W_i \rightarrow U'_i$ is surjective as a map of sheaves (details omitted). Then $W_i \rightarrow U$ is étale and we conclude that $\text{size}(W_i) \leq \text{size}(U)$, see Sets, Lemma 9.7. By Sets, Lemma 9.11 we conclude that $\text{size}(U'_i) \leq \text{size}(U)$. Hence $\coprod_{i \in I} U'_i$ is isomorphic to an object of $(Sch/S)_{fppf}$ by Sets, Lemma 9.5.

Now let $X', U \rightarrow X'$ and $R \rightarrow U \times_{X'} U$ be as in the statement of the lemma. In the previous paragraph we have seen that we can find $U' \in \text{Ob}((Sch/S)_{fppf})$ and a surjective étale morphism $U' \rightarrow X'$ in $Sh((Sch'/S)_{fppf})$. Then $U' \times_{X'} U \rightarrow U'$ is a surjection of sheaves, i.e., we can find an fppf covering $\{U'_i \rightarrow U'\}$ such that $U'_i \rightarrow U'$ factors through $U' \times_{X'} U \rightarrow U'$. By Sets, Lemma 9.12 we can find $\tilde{U} \rightarrow U'$ which is surjective, flat, and locally of finite presentation, with $\text{size}(\tilde{U}) \leq \text{size}(U')$, such that $\tilde{U} \rightarrow U'$ factors through $U' \times_{X'} U \rightarrow U'$. Then we consider

$$\begin{array}{ccccc} U' \times_{X'} U' & \longleftarrow & \tilde{U} \times_{X'} \tilde{U} & \longrightarrow & U \times_{X'} U \\ \downarrow & & \downarrow & & \downarrow \\ U' \times_S U' & \longleftarrow & \tilde{U} \times_S \tilde{U} & \longrightarrow & U \times_S U \end{array}$$

The squares are cartesian. We know the objects of the bottom row are represented by objects of $(Sch/S)_{fppf}$. By the result of the argument of the previous paragraph, the same is true for $U \times_{X'} U$ (as we have the surjection of sheaves $R \rightarrow U \times_{X'} U$ by assumption). Since $(Sch/S)_{fppf}$ is closed under fibre products (by construction), we see that $\tilde{U} \times_{X'} \tilde{U}$ is represented by an object of $(Sch/S)_{fppf}$. Finally, the map $\tilde{U} \times_{X'} \tilde{U} \rightarrow U' \times_{X'} U'$ is a surjection of fppf sheaves as $\tilde{U} \rightarrow U'$ is so. Thus we can once more apply the result of the previous paragraph to conclude that $R' = U' \times_{X'} U'$ is represented by an object of $(Sch/S)_{fppf}$. At this point Lemma 9.1 and Theorem 10.5 imply that $X = h_{U'}/h_{R'}$ is an object of $Spaces/S$ such that $f^{-1}X \cong X'$ as desired. \square

16. Change of base scheme

03I3 In this section we briefly discuss what happens when we change base schemes. The upshot is that given a morphism $S \rightarrow S'$ of base schemes, any algebraic space over S

can be viewed as an algebraic space over S' . And, given an algebraic space F' over S' there is a base change F'_S which is an algebraic space over S . We explain only what happens in case $S \rightarrow S'$ is a morphism of the big fppf site under consideration, if only S or S' is contained in the big site, then one first enlarges the big site as in Section 15.

03I4 **Lemma 16.1.** *Suppose given a big site Sch_{fppf} . Let $g : S \rightarrow S'$ be morphism of Sch_{fppf} . Let $j : (Sch/S)_{fppf} \rightarrow (Sch/S')_{fppf}$ be the corresponding localization functor. Let F be a sheaf of sets on $(Sch/S)_{fppf}$. Then*

- (1) *for a scheme T' over S' we have $j_!F(T'/S') = \coprod_{\varphi: T' \rightarrow S} F(T' \xrightarrow{\varphi} S)$,*
- (2) *if F is representable by a scheme $X \in \text{Ob}((Sch/S)_{fppf})$, then $j_!F$ is representable by $j(X)$ which is X viewed as a scheme over S' , and*
- (3) *if F is an algebraic space over S , then $j_!F$ is an algebraic space over S' , and if $F = U/R$ is a presentation, then $j_!F = j(U)/j(R)$ is a presentation.*

Let F' be a sheaf of sets on $(Sch/S')_{fppf}$. Then

- (4) *for a scheme T over S we have $j^{-1}F'(T/S) = F'(T/S')$,*
- (5) *if F' is representable by a scheme $X' \in \text{Ob}((Sch/S')_{fppf})$, then $j^{-1}F'$ is representable, namely by $X'_S = S \times_{S'} X'$, and*
- (6) *if F' is an algebraic space, then $j^{-1}F'$ is an algebraic space, and if $F' = U'/R'$ is a presentation, then $j^{-1}F' = U'_S/R'_S$ is a presentation.*

Proof. The functors $j_!$, j_* and j^{-1} are defined in Sites, Lemma 24.8 where it is also shown that $j = j_{S/S'}$ is the localization of $(Sch/S')_{fppf}$ at the object S/S' . Hence all of the material on localization functors is available for j . The formula in (1) is Sites, Lemma 26.1. By definition $j_!$ is the left adjoint to restriction j^{-1} , hence $j_!$ is right exact. By Sites, Lemma 24.5 it also commutes with fibre products and equalizers. By Sites, Lemma 24.3 we see that $j_!h_X = h_{j(X)}$ hence (2) holds. If F is an algebraic space over S , then we can write $F = U/R$ (Lemma 9.1) and we get

$$j_!F = j(U)/j(R)$$

because $j_!$ being right exact commutes with coequalizers, and moreover $j(R) = j(U) \times_{j_!F} j(U)$ as $j_!$ commutes with fibre products. Since the morphisms $j(s), j(t) : j(R) \rightarrow j(U)$ are simply the morphisms $s, t : R \rightarrow U$ (but viewed as morphisms of schemes over S'), they are still étale. Thus $(j(U), j(R), s, t)$ is an étale equivalence relation. Hence by Theorem 10.5 we conclude that $j_!F$ is an algebraic space.

Proof of (4), (5), and (6). The description of j^{-1} is in Sites, Section 24. The restriction of the representable sheaf associated to X'/S' is the representable sheaf associated to $X'_S = S \times_{S'} X'$ by Sites, Lemma 26.2. The restriction functor j^{-1} is exact, hence $j^{-1}F' = U'_S/R'_S$. Again by exactness the sheaf R'_S is still an equivalence relation on U'_S . Finally the two maps $R'_S \rightarrow U'_S$ are étale as base changes of the étale morphisms $R' \rightarrow U'$. Hence $j^{-1}F' = U'_S/R'_S$ is an algebraic space by Theorem 10.5 and we win. \square

Note how the presentation $j_!F = j(U)/j(R)$ is just the presentation of F but viewed as a presentation by schemes over S' . Hence the following definition makes sense.

03I5 **Definition 16.2.** Let Sch_{fppf} be a big fppf site. Let $S \rightarrow S'$ be a morphism of this site.

- (1) If F' is an algebraic space over S' , then the *base change of F' to S* is the algebraic space $j^{-1}F'$ described in Lemma 16.1. We denote it F'_S .
- (2) If F is an algebraic space over S , then F *viewed as an algebraic space over S'* is the algebraic space $j_!F$ over S' described in Lemma 16.1. We often simply denote this F ; if not then we will write $j_!F$.

The algebraic space $j_!F$ comes equipped with a canonical morphism $j_!F \rightarrow S$ of algebraic spaces over S' . This is true simply because the sheaf $j_!F$ maps to h_S (see for example the explicit description in Lemma 16.1). In fact, in Sites, Lemma 24.4 we have seen that the category of sheaves on $(Sch/S)_{fppf}$ is equivalent to the category of pairs $(\mathcal{F}, \mathcal{F}' \rightarrow h_S)$ consisting of a sheaf on $(Sch/S')_{fppf}$ and a map of sheaves $\mathcal{F}' \rightarrow h_S$. The equivalence assigns to the sheaf \mathcal{F} the pair $(j_!\mathcal{F}, j_!\mathcal{F} \rightarrow h_S)$. This, combined with the above, leads to the following result for categories of algebraic spaces.

04SG **Lemma 16.3.** *Let Sch_{fppf} be a big fppf site. Let $S \rightarrow S'$ be a morphism of this site. The construction above give an equivalence of categories*

$$\left\{ \begin{array}{l} \text{category of algebraic} \\ \text{spaces over } S \end{array} \right\} \leftrightarrow \left\{ \begin{array}{l} \text{category of pairs } (F', F' \rightarrow S) \text{ consisting} \\ \text{of an algebraic space } F' \text{ over } S' \text{ and a} \\ \text{morphism } F' \rightarrow S \text{ of algebraic spaces over } S' \end{array} \right\}$$

Proof. Let F be an algebraic space over S . The functor from left to right assigns the pair $(j_!F, j_!F \rightarrow S)$ of F which is an object of the right hand side by Lemma 16.1. Since this defines an equivalence of categories of sheaves by Sites, Lemma 24.4 to finish the proof it suffices to show: if F is a sheaf and $j_!F$ is an algebraic space, then F is an algebraic space. To do this, write $j_!F = U'/R'$ as in Lemma 9.1 with $U', R' \in \text{Ob}((Sch/S')_{fppf})$. Then the compositions $U' \rightarrow j_!F \rightarrow S$ and $R' \rightarrow j_!F \rightarrow S$ are morphisms of schemes over S' . Denote U, R the corresponding objects of $(Sch/S)_{fppf}$. The two morphisms $R' \rightarrow U'$ are morphisms over S and hence correspond to morphisms $R \rightarrow U$. Since these are simply the same morphisms (but viewed over S) we see that we get an étale equivalence relation over S . As $j_!$ defines an equivalence of categories of sheaves (see reference above) we see that $F = U/R$ and by Theorem 10.5 we see that F is an algebraic space. \square

The following lemma is a slight rephrasing of the above.

04SH **Lemma 16.4.** *Let Sch_{fppf} be a big fppf site. Let $S \rightarrow S'$ be a morphism of this site. Let F' be a sheaf on $(Sch/S')_{fppf}$. The following are equivalent:*

- (1) *The restriction $F'|_{(Sch/S)_{fppf}}$ is an algebraic space over S , and*
- (2) *the sheaf $h_S \times F'$ is an algebraic space over S' .*

Proof. The restriction and the product match under the equivalence of categories of Sites, Lemma 24.4 so that Lemma 16.3 above gives the result. \square

We finish this section with a lemma on a compatibility.

03I6 **Lemma 16.5.** *Let Sch_{fppf} be a big fppf site. Let $S \rightarrow S'$ be a morphism of this site. Let F be an algebraic space over S . Let T be a scheme over S and let $f : T \rightarrow F$ be a morphism over S . Let $f' : T' \rightarrow F'$ be the morphism over S' we get from f by applying the equivalence of categories described in Lemma 16.3. For any property \mathcal{P} as in Definition 5.1 we have $\mathcal{P}(f') \Leftrightarrow \mathcal{P}(f)$.*

Proof. Suppose that U is a scheme over S , and $U \rightarrow F$ is a surjective étale morphism. Denote U' the scheme U viewed as a scheme over S' . In Lemma 16.1 we have seen that $U' \rightarrow F'$ is surjective étale. Since

$$j(T \times_{f,F} U) = T' \times_{f',F'} U'$$

the morphism of schemes $T \times_{f,F} U \rightarrow U$ is identified with the morphism of schemes $T' \times_{f',F'} U' \rightarrow U'$. It is the same morphism, just viewed over different base schemes. Hence the lemma follows from Lemma 11.4. \square

17. Other chapters

- | | |
|------------------------------------|-------------------------------------|
| Preliminaries | (36) More on Morphisms |
| (1) Introduction | (37) More on Flatness |
| (2) Conventions | (38) Groupoid Schemes |
| (3) Set Theory | (39) More on Groupoid Schemes |
| (4) Categories | (40) Étale Morphisms of Schemes |
| (5) Topology | Topics in Scheme Theory |
| (6) Sheaves on Spaces | (41) Chow Homology |
| (7) Sites and Sheaves | (42) Intersection Theory |
| (8) Stacks | (43) Picard Schemes of Curves |
| (9) Fields | (44) Adequate Modules |
| (10) Commutative Algebra | (45) Dualizing Complexes |
| (11) Brauer Groups | (46) Duality for Schemes |
| (12) Homological Algebra | (47) Discriminants and Differents |
| (13) Derived Categories | (48) Local Cohomology |
| (14) Simplicial Methods | (49) Algebraic and Formal Geometry |
| (15) More on Algebra | (50) Algebraic Curves |
| (16) Smoothing Ring Maps | (51) Resolution of Surfaces |
| (17) Sheaves of Modules | (52) Semistable Reduction |
| (18) Modules on Sites | (53) Fundamental Groups of Schemes |
| (19) Injectives | (54) Étale Cohomology |
| (20) Cohomology of Sheaves | (55) Crystalline Cohomology |
| (21) Cohomology on Sites | (56) Pro-étale Cohomology |
| (22) Differential Graded Algebra | Algebraic Spaces |
| (23) Divided Power Algebra | (57) Algebraic Spaces |
| (24) Hypercoverings | (58) Properties of Algebraic Spaces |
| Schemes | (59) Morphisms of Algebraic Spaces |
| (25) Schemes | (60) Decent Algebraic Spaces |
| (26) Constructions of Schemes | (61) Cohomology of Algebraic Spaces |
| (27) Properties of Schemes | (62) Limits of Algebraic Spaces |
| (28) Morphisms of Schemes | (63) Divisors on Algebraic Spaces |
| (29) Cohomology of Schemes | (64) Algebraic Spaces over Fields |
| (30) Divisors | (65) Topologies on Algebraic Spaces |
| (31) Limits of Schemes | (66) Descent and Algebraic Spaces |
| (32) Varieties | (67) Derived Categories of Spaces |
| (33) Topologies on Schemes | (68) More on Morphisms of Spaces |
| (34) Descent | (69) Flatness on Algebraic Spaces |
| (35) Derived Categories of Schemes | (70) Groupoids in Algebraic Spaces |

- | | |
|---------------------------------------|--------------------------------------|
| (71) More on Groupoids in Spaces | (90) Artin's Axioms |
| (72) Bootstrap | (91) Quot and Hilbert Spaces |
| (73) Pushouts of Algebraic Spaces | (92) Properties of Algebraic Stacks |
| Topics in Geometry | (93) Morphisms of Algebraic Stacks |
| (74) Chow Groups of Spaces | (94) Limits of Algebraic Stacks |
| (75) Quotients of Groupoids | (95) Cohomology of Algebraic Stacks |
| (76) More on Cohomology of Spaces | (96) Derived Categories of Stacks |
| (77) Simplicial Spaces | (97) Introducing Algebraic Stacks |
| (78) Duality for Spaces | (98) More on Morphisms of Stacks |
| (79) Formal Algebraic Spaces | (99) The Geometry of Stacks |
| (80) Restricted Power Series | Topics in Moduli Theory |
| (81) Resolution of Surfaces Revisited | (100) Moduli Stacks |
| Deformation Theory | (101) Moduli of Curves |
| (82) Formal Deformation Theory | Miscellany |
| (83) Deformation Theory | (102) Examples |
| (84) The Cotangent Complex | (103) Exercises |
| (85) Deformation Problems | (104) Guide to Literature |
| Algebraic Stacks | (105) Desirables |
| (86) Algebraic Stacks | (106) Coding Style |
| (87) Examples of Stacks | (107) Obsolete |
| (88) Sheaves on Algebraic Stacks | (108) GNU Free Documentation License |
| (89) Criteria for Representability | (109) Auto Generated Index |

References

- [Art69a] Michael Artin, *Algebraic approximation of structures over complete local rings*, Inst. Hautes Études Sci. Publ. Math. (1969), no. 36, 23–58.
- [Art69b] ———, *Algebraization of formal moduli: I*, Global Analysis (Papers in Honor of K. Kodaira), Univ. Tokyo Press, Tokyo, 1969, pp. 21–71.
- [Art69c] ———, *The implicit function theorem in algebraic geometry*, Algebraic Geometry (Internat. Colloq., Tata Inst. Fund. Res., Bombay, 1968), Oxford Univ. Press, London, 1969, pp. 13–34.
- [Art70] ———, *Algebraization of formal moduli: II – existence of modifications*, Annals of Mathematics **91** (1970), 88–135.
- [Art71a] ———, *Algebraic spaces*, Yale University Press, New Haven, Conn., 1971, A James K. Whittemore Lecture in Mathematics given at Yale University, 1969, Yale Mathematical Monographs, 3.
- [Art71b] ———, *Construction techniques for algebraic spaces*, Actes du Congrès International des Mathématiciens (Nice, 1970), Tome 1, Gauthier-Villars, Paris, 1971, pp. 419–423.
- [Art73] ———, *Théorèmes de représentabilité pour les espaces algébriques*, Les Presses de l'Université de Montréal, Montreal, Que., 1973, En collaboration avec Alexandru Lascu et Jean-François Boutot, Séminaire de Mathématiques Supérieures, No. 44 (Été, 1970).
- [Art74] ———, *Versal deformations and algebraic stacks*, Inventiones Mathematics **27** (1974), 165–189.
- [Jac64] Nathan Jacobson, *Lectures in abstract algebra. Vol III: Theory of fields and Galois theory*, D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London-New York, 1964.
- [Knu71] Donald Knutson, *Algebraic spaces*, Lecture Notes in Mathematics, vol. 203, Springer-Verlag, 1971.